



Charm Spectroscopy from B factories



Jose Benitez (SLAC)

*Representing the **BABAR** Collaboration*

Charm 2010 Workshop

Beijing
October 23, 2010



Contact: benitezj@slac.stanford.edu

Production of Charm Mesons at B-factories

- *B* factories produce charm mesons directly from $c\bar{c}$ hadronization or from decays of *B* mesons.

Inclusive Production: $e^+e^- \rightarrow c\bar{c} \rightarrow D_{(s)}^{**} X$

Exclusive Production: $e^+e^- \rightarrow b\bar{b} \rightarrow BB, B \rightarrow D_{(s)}^{**} X$

where $D_{(s)}^{**}$ can be some excited charm meson.



$\mathcal{L} \sim 550 \text{ fb}^{-1}$

→ ~700 Million $c\bar{c}$

→ ~550 Million $b\bar{b}$



$\mathcal{L} \sim 1000 \text{ fb}^{-1}$


→ ~1300 Million $c\bar{c}$

→ ~1000 Million $b\bar{b}$

$e^+e^- \rightarrow$	Cross-section (nb)
$b\bar{b}$	1.05
$c\bar{c}$	1.30
$s\bar{s}$	0.35
$u\bar{u}$	1.39
$d\bar{d}$	0.35
$\tau^+\tau^-$	0.94
$\mu^+\mu^-$	1.16
e^+e^-	~ 40

Predictions for the D and D_s states

- Predictions of the D and D_s mass eigenstates were performed since 1985 using QCD potential models.
- Recently (2001) the D_s spectrum predictions have been updated.
- The predicted masses of the excited states are generally in qualitative agreement with observations, however, for some states large quantitative differences exist.




The History

First observations:

D		D_s	
□ $1^1S_0 : D$	Mark I 1975	□ $1^1S_0 : D_s$	CLEO 1983
□ $1^3S_1 : D^*$	Mark I 1977	□ $1^3S_1 : D_s^*$	TPC 1984
□ $1^3P_1 : D_1(2420)$	ARGUS1986	□ $1^1P_1 : D_{s1}(2536)$	ARGUS1989
□ $1^3P_2 : D_2^*(2460)$	TPS 1989	□ $1^3P_2 : D_{s2}^*(2573)$	CLEO2 1994
□ $1^3P_0 : D_0^*(2400)$	BELLE 2004	□ $1^3P_1 : D_{s1}(2460)$	CLEO2 2003
□ $1^1P_1 : D_1(2430)$	BELLE 2004	□ $1^3P_0 : D_{s0}^*(2317)$	BaBar 2003
□ $2^1S_0 : D(2550)$	BaBar 2010	□ $2^3S_1 : D_s^*(2710)$	BaBar 2006
□ $2^3S_1 : D^*(2600)$	BaBar 2010	□ $1^?D_? : D_s^*(2860)$	BaBar 2006
□ $1^?D_? : D(2750)$	BaBar 2010	□ $?^?D_? : D_s(3040)$	BaBar 2009
□ $1^?D_? : D^*(2760)$	BaBar 2010		


Recent studies of these states will be presented in this talk.

The D_{s0}^* (2317)




The $D_{s0}^*(2317)$

- In an inclusive study of the $D_s\pi^0$ system BaBar discovered the narrow $D_{s0}^*(2317)$ state (2003).
- The low unexpected mass of this state triggered many subsequent studies by both BaBar and BELLE including the following:
 - $B \rightarrow D_{s0}^*(2317) D$
[BELLE $\sim 110 \text{ fb}^{-1}$ Phys. Rev. Lett. 91, 262002 (2003)]
 - $D_{s0}^*(2317) \rightarrow D_s\pi^0$ inclusive
[BELLE 87 fb^{-1} Phys. Rev. Lett. 92, 012002 (2004)]
 - $B \rightarrow D_{s0}^*(2317) D^{(*)}$
[BaBar 113 fb^{-1} Phys. Rev. Lett. 93, 181801 (2004)]
 - $B \rightarrow D_{s0}^*(2317) K$
[BELLE 140 fb^{-1} Phys. Rev. Lett. 94, 061802 (2005)]



[BaBar 91 fb^{-1}
Phys. Rev. Lett. **90**, 202001 (2003)]

Precision $D_{s1}(2536)$ Parameters




Precision $D_{s1}(2536)$ Parameters

- Preliminary BaBar analysis using 384 fb^{-1} [presented at ICHEP 2010].
- Inclusive reconstruction of $D_{s1}(2536)^+ \rightarrow D^{*+} K_S$ where $D^{*+} \rightarrow D^0 \pi^+$ and $D^0 \rightarrow K^- \pi^+$ or $K^- \pi^+ \pi^- \pi^+$
- Parameters determined from the mass difference $\Delta m = m(D^* K_S) - m(D^*) - m(K_S)$ resolution is about 0.26 MeV.
- Preliminary results:

$$m(D_{s1}^+) = (2535.10 \pm 0.01 \pm 0.18) \text{ MeV}/c^2$$




$$\Gamma(D_{s1}^+) = (0.92 \pm 0.03 \pm 0.04) \text{ MeV}$$
- This is the first measurement of the $D_{s1}(2536)$ width.



$D_{s1}(2536)$ Angular Analysis


BaBar Preliminary

- Signal yield is extracted as a function of the helicity angle θ' .
- Angular distribution indicates un-natural spin-parity.
- Significant D -wave contribution is present in the decay.




J^P	$dN/d \cos \theta'$	$\chi^2/NDF(K4\pi)$	$\chi^2/NDF(K6\pi)$
0^+	forbidden	—	—
0^-	$\propto \cos^2 \theta'$	2142.7/19	2440.8/19
$1^-, 2^+, 3^-, \dots$	$\propto \sin^2 \theta'$	103.2/19	108.8/19
$1^+, 2^-, 3^+, \dots$ (S -wave only)	const	392.1/19	425.1/19
$1^+, 2^-, 3^+, \dots$ (S -, D -wave)	$\propto (\sin^2 \theta' + \beta \cos^2 \theta')$	24.9/18 ($\beta = 0.23 \pm 0.03$)	9.5/18 ($\beta = 0.24 \pm 0.03$)

Inclusive Study of DK and D^*K Systems



Inclusive Study of DK and D^*K Systems

- The following channels have been analyzed:
 - D^0K^+ using $D^0 \rightarrow K^-\pi^+$
 - D^+K_S using $D^+ \rightarrow K^-\pi^+\pi^+$
 - $D^{*+}K_S$ using $D^{*+} \rightarrow D^0\pi^+, D^+\pi^0$ ($D \rightarrow K^-\pi^+, K^-\pi^+\pi^-, \pi^+, D^+ \rightarrow K^-\pi^+\pi^+$)
 - $D^{*0}K^+$ using $D^{*0} \rightarrow D^0\pi^0$ ($D^0 \rightarrow K^-\pi^+$)



[BABAR (470 fb^{-1}) *Phys. Rev. D* **80**, 092003 (2009)]

Parameters of D_{sJ} Structures

- The mass values of the $D_s^*(2710)$ and $D_s^*(2860)$ are close to those of the first radial excitation of the D_s^* and $L=2$ excited states, respectively.

$$m(D_{s1}^*(2710)^+) = 2710 \pm 2_{\text{stat}} \left(\begin{smallmatrix} +12 \\ -7 \end{smallmatrix} \right)_{\text{syst}} \text{ MeV}/c^2,$$

$$\Gamma = 149 \pm 7_{\text{stat}} \left(\begin{smallmatrix} +39 \\ -52 \end{smallmatrix} \right)_{\text{syst}} \text{ MeV},$$

- The mass of the $D_s(3040)$ is close to that of the second radial excitation predicted in Ref. [T. Matsuki *etal.*, *Eur. Phys. J. A* 31, 701 (2007)]

$$m(D_{sJ}^*(2860)^+) = 2862 \pm 2_{\text{stat}} \left(\begin{smallmatrix} +5 \\ -2 \end{smallmatrix} \right)_{\text{syst}} \text{ MeV}/c^2,$$

$$\Gamma = 48 \pm 3_{\text{stat}} \pm 6_{\text{syst}} \text{ MeV},$$

- The ratios of the D^*K over DK branching fractions have been determined.

$$m(D_{sJ}(3040)) = 3044 \pm 8_{\text{stat}} \left(\begin{smallmatrix} +30 \\ -5 \end{smallmatrix} \right)_{\text{syst}} \text{ MeV}/c^2,$$

$$\Gamma = 239 \pm 35_{\text{stat}} \left(\begin{smallmatrix} +46 \\ -42 \end{smallmatrix} \right)_{\text{syst}} \text{ MeV}.$$


- For the $D_s^*(2710)$ the ratio is consistent with the value predicted for the first radial excitation. [P. Colangelo *etal.*, *Phys.Rev. D* 77, 014012 (2008)]

$$\frac{\mathcal{B}(D_{s1}^*(2710)^+ \rightarrow D^*K)}{\mathcal{B}(D_{s1}^*(2710)^+ \rightarrow DK)} = 0.91 \pm 0.13_{\text{stat}} \pm 0.12_{\text{syst}},$$

$$\frac{\mathcal{B}(D_{sJ}^*(2860)^+ \rightarrow D^*K)}{\mathcal{B}(D_{sJ}^*(2860)^+ \rightarrow DK)} = 1.10 \pm 0.15_{\text{stat}} \pm 0.19_{\text{syst}}.$$


Angular Analysis of the D^*K System

- The helicity angle θ_h defined by the Kaon and pion provides information about the quantum numbers of the resonances.
- For both the $D_s^*(2710)$ and $D_s^*(2860)$, the angular distribution is consistent with natural parity.
- For the $D_s(3040)$ the angular distribution is not conclusive.



[BABAR (470 fb^{-1}) *Phys. Rev. D* **80**, 092003 (2009)]

Exclusive Studies of Broad D States




Exclusive Studies of Broad D States

- Parameters of the $D_1(2430)$ determined from a Dalitz plot analysis of $B^+ \rightarrow D^{*-} \pi^+ \pi^+$:

$$M_{D_1^{*0}} = 2427 \pm 26 \pm 20 \pm 15 \text{ MeV}/c^2,$$

$$\Gamma_{D_1^{*0}} = 384_{-75}^{+107} \pm 24 \pm 70 \text{ MeV}.$$




[**BELLE 60 fb⁻¹** *Phys. Rev. D* **69**, 112002 (2004)]

- The $D_0^*(2400)$ was confirmed by BaBar and its parameters determined from a Dalitz plot analysis of $B^+ \rightarrow D^- \pi^+ \pi^+$:


$$m_{D_0^{*0}} = (2297 \pm 8 \pm 5 \pm 19) \text{ MeV}/c^2$$

$$\Gamma_{D_0^{*0}} = (273 \pm 12 \pm 17 \pm 45) \text{ MeV},$$



[**BaBar 350 fb⁻¹** *Phys. Rev. D* **79**, 112004 (2009)]

Inclusive Study of $D\pi$ and $D^*\pi$



Inclusive Study of $D\pi$ and $D^*\pi$


- Analysis of $D\pi$ and $D^*\pi$ systems produced from $c\bar{c}$ events:

$$e^+e^- \rightarrow c\bar{c} \rightarrow D^{**}X \rightarrow D^{(*)}\pi X$$

X represents any additional system.

- The following channels are reconstructed:

- $$\begin{aligned}
&\bullet \quad D^{**0} \rightarrow D^+ \pi^- \\
&\qquad \qquad \qquad \downarrow \\
&\qquad \qquad K^- \pi^+ \pi^+ \\
\\
&\bullet \quad D^{**+} \rightarrow D^0 \pi^+ \\
&\qquad \qquad \qquad \downarrow \\
&\qquad \qquad K^- \pi^+ \\
\\
&\bullet \quad D^{**0} \rightarrow D^{*+} \pi^- \\
&\qquad \qquad \qquad \downarrow \\
&\qquad \qquad D^0 \pi^+ \\
&\qquad \qquad \qquad \downarrow \\
&\qquad \qquad K^- \pi^+ \text{ or } K^- \pi^+ \pi^- \pi^+
\end{aligned}$$




- The data set corresponds to about **10 times** more signal events than the previous study by the CDF collaboration.

[BABAR (454 fb⁻¹) *arXiv:1009.2076, submitted to PRD-RC (2010)*]

The $D^+\pi^-$ System

BaBar Preliminary

- The mass distribution of the $D^+\pi^-$ final state presents a prominent signal from the $D_2^*(2460)$ as well as two new structures at 2.60 GeV and 2.76 GeV.
- The peaking background at 2.30 GeV is due to $D_2^*(2460)$ and $D_1(2420)$ decaying to $D^*\pi$ where the slow pion is missing.
- The broad $D_0^*(2400)$ improves the fit quality, its parameters are floated within 2σ from the known values.
- The χ^2/NDF of the fit is 281/242.




[BABAR (454 fb⁻¹)
arXiv:1009.2076, submitted to PRD-RC (2010)]

The $D^0\pi^+$ System

BaBar Preliminary

- To confirm the new signals, the $D^0\pi^+$ system is analyzed.
- In this channel the feed-down backgrounds are stronger and the signal statistics of this are smaller so the widths of all signals are fixed to the widths measured in the $D^+\pi^-$.
- The mass values obtained are a few MeV higher than in $D^+\pi^-$, consistent with being the isospin partners.
- The fit quality is $\chi^2/\text{NDF}=278/224$.




[BABAR (454 fb⁻¹)

arXiv:1009.2076, submitted to PRD-RC (2010)]

The $D^{*+}\pi^-$ System

BaBar Preliminary

- The $D^*\pi$ system shows prominent signals of $D_1(2420)$ and $D_2^*(2460)$.
- In addition, there are new structures in the higher mass region. The region at 2.60 GeV is populated by two signals, while at 2.75 GeV there is a signal similar to the $D^*(2760)$ from $D^+\pi^-$.
- The parameters of the $D_2^*(2460)$ and $D^*(2600)$ are fixed to the ones from the $D^+\pi^-$.
- The fit quality is $\chi^2/\text{NDF}=244/207$.




[BABAR (454 fb⁻¹)
arXiv:1009.2076, submitted to PRD-RC (2010)]

$D^{*+}\pi^{-}$ with $|\cos\theta_H| > 0.75$

BaBar Preliminary

- The selection $|\cos(\theta_H)| > 0.75$ is applied to suppress the resonances with natural spin-parity ($dN/d\cos\theta_H \sim \sin^2\theta_H$).
- This fit allows to determine the parameters of the $D(2550)$ under the assumption that the $D^*(2600)$ is the same signal observed in $D^+\pi^-$.
- The parameters of the $D_2^*(2460)$ and $D^*(2600)$ are fixed to the values from $D^+\pi^-$.
- This fit also determines the parameters of the $D_1(2420)$.
- The fit quality is $\chi^2/\text{NDF}=214/205$.




[BABAR (454 fb⁻¹)
arXiv:1009.2076, submitted to PRD-RC (2010)]

$D^{*+}\pi^{-}$ with $|\cos\theta_H|<0.5$

BaBar Preliminary

- The selection $|\cos\theta_H|<0.5$ is applied to favor the resonances with natural spin-parity.
- In this fit, the parameters of all signals, except the $D(2750)$, are fixed to the values from the previous fits.
- This fit allows to observe clearly the $D^*(2600)$ signal and shows consistency in the fit model.
- The fit quality is $\chi^2/\text{NDF}=210/209$.



[BABAR (454 fb⁻¹)
arXiv:1009.2076, submitted to PRD-RC (2010)]

Resonance Parameters:

- The uncertainties on most parameters are dominated by systematic uncertainties.
- The systematic uncertainty includes the following sources: bin size and mass range of the histogram, errors on parameters fixed in the fits, Breit-Wigner shape of the new signals, a possible contribution from the $D_1(2430)$, and background modeling.
- The significance of the new signals is estimated from the yield over its total uncertainty.




Resonance	Channel(Fit)	Efficiency (%)	Yield ($\times 10^3$)	Mass MeV/ c^2	Width MeV	Significance
$D_1(2420)^0$	$D^{*+}\pi^-$ (C)		$102.8 \pm 1.3 \pm 2.3$	$2420.1 \pm 0.1 \pm 0.8$	$31.4 \pm 0.5 \pm 1.3$	
	$D^{*+}\pi^-$ (E)	1.09 ± 0.03	$214.6 \pm 1.2 \pm 6.4$	2420.1(fixed)	31.4(fixed)	
$D_2^*(2460)^0$	$D^+\pi^-$ (A)	1.29 ± 0.03	$242.8 \pm 1.8 \pm 3.4$	$2462.2 \pm 0.1 \pm 0.8$	$50.5 \pm 0.6 \pm 0.7$	
	$D^{*+}\pi^-$ (E)	1.12 ± 0.04	$136 \pm 2 \pm 13$	2462.2(fixed)	50.5(fixed)	
$D(2550)^0$	$D^{*+}\pi^-$ (C)		$34.3 \pm 6.7 \pm 9.2$	$2539.4 \pm 4.5 \pm 6.8$	$130 \pm 12 \pm 13$	3.0σ
	$D^{*+}\pi^-$ (E)	1.14 ± 0.04	$98.4 \pm 8.2 \pm 38$	2539.4(fixed)	130(fixed)	
$D^*(2600)^0$	$D^+\pi^-$ (A)	1.35 ± 0.05	$26.0 \pm 1.4 \pm 6.6$	$2608.7 \pm 2.4 \pm 2.5$	$93 \pm 6 \pm 13$	3.9σ
	$D^{*+}\pi^-$ (D)		$50.2 \pm 3.0 \pm 6.7$	2608.7(fixed)	93(fixed)	7.3σ
	$D^{*+}\pi^-$ (E)	1.18 ± 0.05	$71.4 \pm 1.7 \pm 7.3$	2608.7(fixed)	93(fixed)	
$D(2750)^0$	$D^{*+}\pi^-$ (E)	1.23 ± 0.07	$23.5 \pm 2.1 \pm 5.2$	$2752.4 \pm 1.7 \pm 2.7$	$71 \pm 6 \pm 11$	4.2σ
$D^*(2760)^0$	$D^+\pi^-$ (A)	1.41 ± 0.09	$11.3 \pm 0.8 \pm 1.0$	$2763.3 \pm 2.3 \pm 2.3$	$60.9 \pm 5.1 \pm 3.6$	8.9σ
$D_2^*(2460)^+$	$D^0\pi^+$ (B)		$110.8 \pm 1.3 \pm 7.5$	$2465.4 \pm 0.2 \pm 1.1$	50.5(fixed)	
$D^*(2600)^+$	$D^0\pi^+$ (B)		$13.0 \pm 1.3 \pm 4.5$	$2621.3 \pm 3.7 \pm 4.2$	93(fixed)	2.8σ
$D^*(2760)^+$	$D^0\pi^+$ (B)		$5.7 \pm 0.7 \pm 1.5$	$2769.7 \pm 3.8 \pm 1.5$	60.9(fixed)	3.5σ

[BABAR (454 fb⁻¹) *arXiv:1009.2076, submitted to PRD-RC (2010)*]

Angular Analysis of $D^{*+}\pi^-$

BaBar Preliminary




- The signal yields have been extracted as a function of $\cos\theta_H$.
- The $D_1(2420)$ shows a $1+A\cos^2\theta_H$ distribution indicating unnatural spin-parity. The value for A indicates a significant S-wave contribution in the decay.
- The $D_2^*(2460)$ shows a $\sin^2\theta_H$ distribution consistent with the natural spin-parity assignment.



Angular Analysis of $D^{*+}\pi^-$


BaBar Preliminary

- For the signal $D(2550)$ a $\cos^2(\theta_H)$ distribution is obtained consistent with a $J^P=0^-$ value.
- For the signal $D^*(2600)$ a $\sin^2(\theta_H)$ distribution is obtained consistent with natural spin-parity.
- For the signal $D(2750)$ the interpretation of the distribution is not conclusive.



Interpretation of the $D\pi$ and $D^*\pi$ Results

- The $D^*(2600)$ signal observed in $D\pi$ and $D^*\pi$ has a mass value and helicity distribution consistent with the first radial excitation of the $D^*(2010)$.
- Likewise, the $D(2550)$ observed in $D^*\pi$ has a mass value and helicity distribution consistent with the first radial excitation of the D^0 .
- The $D(2750)$ observed in $D^*\pi$ has mass value lower than the $D^*(2760)$ observed in $D^+\pi^-$. The helicity distribution is not conclusive. These two signals may be due to the four L=2 excited states.



Conclusions

- The B -factories have large potential for advancing the understanding of the charmed hadron spectrum. In this talk, recent studies of the D and D_s mesons have been presented.
- The spectroscopy of charmed mesons has revived in recent years (2009-2010) with the observations by the BaBar of new structures in the DK , D^*K , $D\pi$, and $D^*\pi$ systems. These studies find candidates for the radial and $L=2$ excited states of the D_s and D mesons.
- Precision measurements of the narrow $L=1$ D_s mesons are possible from the large Data sets. A first measurement of the $D_{s1}(2536)$ decay width has been presented here.
- Charmed mesons obtained from B decays allow for the study of the broad $L=1$ states. However, updated studies of these decays are needed. In particular an analysis of $B^+ \rightarrow D^{*-}\pi^+\pi^+$ might provide much better parameter values for the $D_1(2430)$, and evidence for new the structures observed in the inclusive $D^{(*)}\pi$ analyses.