




# AGN Family

- Classification
- AGN Types
- Interpretation



Lectures available at:

[http://www.chara.gsu.edu/~crenshaw/beijing\\_agn.html](http://www.chara.gsu.edu/~crenshaw/beijing_agn.html)

# AGN Classification

AGN have been classified in the past on the basis of:

- Appearance of their optical spectra (Sey 1 vs. Sey 2)
- Luminosity (Seyfert  $\rightarrow$  Quasar)
- Radio Power (Radio Loud vs. Radio Quiet)
- Morphology (FR I vs. FR II; host galaxy type)
- A number of other properties

Ultimately, we want a more physical description, e.g.:

- Black hole mass ( $M$ )
- Luminosity ( $L$ )
- Eddington Ratio ( $L/L_{\text{Edd}} \sim L/M$ )
- Radio Power



# Seyfert Galaxies




- Nucleus - absolute blue magnitude:  $M_B > -21.5$  (to distinguish from quasars)
- $L_{\text{Bol}} = 10^{43} - 10^{45} \text{ ergs s}^{-1}$
- “Classic” Seyferts:  $z < 0.1$  (SDSS has many higher  $z$  Seys.)
- Broad permitted lines (FWHM = 800 - 8000  $\text{km s}^{-1}$ ) from BLR
- Narrow permitted and forbidden lines (FWHM = 200 - 500  $\text{km s}^{-1}$ ) from NLR
  - Seyfert 1: both BLR and NLR, strong nonstellar continuum
  - Seyfert 2: only NLR, weak continuum (mostly stellar)
- Spectropolarimetry (Antonucci 1985) shows hidden BLR in some Seyfert 2s:
  - Balmer lines scattered into the line of sight by electrons and/or dust  
→ Unified model

- Additional Osterbrock types:
  - Seyfert 1.5: narrow permitted components are easily seen
  - Seyfert 1.8: weak broad H $\alpha$  and H $\beta$
  - Seyfert 1.9: only weak broad H $\alpha$  detectable
  - Narrow-line Seyfert 1 galaxies (NLS1s) (not Seyfert 2s!)
    - FWHM (BLR) = 800 – 2000 km/sec
    - Strong Fe II (high density region like other BLRs)
    - strong excess below 1 – 2 keV and rapid X-ray variability
- Seyferts are weak radio sources (radio blobs rather than jets)
- Strong X-ray sources at  $E > 2$  keV
  - Seyfert 2 galaxies are often weak in soft X-rays ( $E < 2$  keV), due to absorption by a large column of gas (e.g., the torus)
- Seyfert host galaxies are almost always spirals

## Seyfert 1.5 - BLR+NLR




# Seyfert 1.9






# Quasars



- At redshifts  $z = 0.1$  to  $\sim 7$
- Higher luminosities than Seyferts:  $L = 10^{45} - 10^{47} \text{ ergs s}^{-1}$
- **Quasars** (quasi-stellar radio sources): discovered first by radio surveys, emission-line spectra revealed high redshifts
- **QSOs** (quasi-stellar objects): discovered optically from their strong blue continua, broad emission lines, X-ray flux, etc.
  - The terms “quasars” and “QSOs” have become interchangeable; now we use **radio-loud quasars (RLQ)** and **radio-quiet quasars (RQQ)**
  - Radio loud:  $\nu F_{\nu}(6 \text{ cm}) / \nu F_{\nu}(4400 \text{ \AA}) \geq 10$
  - Only 5 - 10% of all quasars are RLQ
- Quasars have spectra like Seyfert 1 galaxies, but
  - stellar absorption features not easily detected
  - narrow-lines tend to be weak
- Type 2 quasars (no broad lines) have also been detected

# Quasar Spectrum






# Radio Galaxies




- Low-luminosity analogs of RL quasars  
(Seyferts are low-luminosity analogs of RQ quasars)
- Characterized by compact radio core, lobes, and (often) jets
  1. FR I: lower luminosity; bright in center and weak toward edges
  2. FR II: high luminosity; brighter at edges
  3. Dividing line:  $L_v = 10^{32} \text{ ergs s}^{-1} \text{ Hz}^{-1}$
- Radio galaxies with emission lines are similar to Seyferts, but are typically found in giant ellipticals (E or cD)
- **Broad-line radio galaxies (BLRG):** similar to Seyferts 1s, but
  1. Balmer profiles are broader and more flat-topped
  2. Fe II emission is weaker
  3.  $H\alpha/H\beta$  ratios higher (steeper Balmer decrement)
- **Narrow-line radio galaxies (NLRG):** optical spectra are essentially identical to Seyfert 2s

# NLRG



# BLRG



# Blazars


- Defined by 1) strong variability (time scales one day or less) from radio to X-rays and high polarizations (1 – 4 %)
- Moderate to strong radio sources (radio loud)
- Two classes:
  - 1) **BL-Lac objects**: no strong emission or absorption lines
    - likely beamed FR Is
  - 2) **Optically-violent** variables (OVVs): highly polarized, variable, but have broad emission lines like quasars
    - likely beamed FR IIs
- Continuous spectra are less complicated than those of quasars – likely synchrotron radiation plus Compton “upscattering”
- Interpretation: relativistically beamed jets close to our line of sight (overwhelms other emission components)
- Two types of BL Lacs:
  - High-frequency BL Lacs (HBLs): synchrotron peak in X-rays (or XBLs)
  - Low-frequency BL Lacs (LBLs): synchrotron peak in radio (or RBLs)

# Blazar SEDs (Urry 1998)

Synchrotron



Inverse Compton




Inverse Compton Models:

- Synchrotron Self Compton (SSC)
- External Compton (EC): seed photons from accretion disk, BLR, Cosmic Microwave Background (CMB), etc.



# Radio Galaxy Unification




FR IIs:  
OVV, RLQSO,  
BLRG, NLRG

FR Is:  
BL Lac, WLRG,  
(weak-lined  
radio galaxy)

Barthel et  
al. (1989)




# LINERS



- Low-ionization nuclear emission-line regions (LINERs) (Heckman 1980)
  - Strong low-ionization lines like Seyferts: [O I], [S II], [N II]
  - However, high-ionization lines are weak e.g., ([O III]/H $\beta$  < 3)
- Lower luminosities than Seyferts:  $10^{39} - 10^{42}$  ergs s $^{-1}$ 
  - Difficult to detect against background of host galaxy
- Recent evidence shows that most LINERs are AGN (previous explanations include very hot stars and shock heating)
- About 1/3 of all luminous galaxies (including Ellipticals) contain LINERs!
- Broad Balmer emission detected in  $\sim 20\%$  (type 1 LINERs)
- LINERs are more radio-loud than Seyferts
- There are some transition objects, which may be a combination of starbursts (H II galaxies) and AGN

# LINER (NGC 1052)



normal galaxy

## AGN – Approximate Space Densities (Local)

Type of Object	# per Mpc <sup>3</sup>
Field galaxies	$10^{-1}$
Luminous spirals	$10^{-2}$
LINERs	$3 \times 10^{-3}$
Seyfert galaxies	$10^{-4}$ ( $\sim 1\%$ of spirals)
Radio galaxies	$10^{-6}$
Radio-quiet quasars (QSOs)	$10^{-7}$
Radio-loud quasars	$10^{-9}$

(Osterbrock, p. 310)