

# Recent developments on hadron interaction and dynamically generated resonances

E. Oset, A. Ramos, Xie Ju Jun, M. Albaladejo, R. Molina, J. Garzon, Chu Wen Xiao, J.J. Wu, B.S. Zou

Chiral dynamics and the local hidden gauge approach

Meson-meson interaction ( vector-vector interaction)

Vector baryon interaction

Vector- Vector and Vector-Baryon molecules

Evidence for a new  $h_1$  state around 1820 MeV

The  $\gamma p \rightarrow K^0 \Sigma^+$  and  $\gamma n \rightarrow K^0 \Sigma^0$  in the  $K^* \Lambda$  threshold  $\rightarrow$

Evidence for a new baryon resonance  $1/2^-$  around 2035 MeV

Recent developments in the charm and beauty sectors

# Hidden gauge formalism for vector mesons, pseudoscalars and photons

Bando et al. PRL, 112 (85); Phys. Rep. 164, 217 (88)

Meissner, U.G., Phys. Rep. 161,213 (88)

$$\mathcal{L} = \mathcal{L}^{(2)} + \mathcal{L}_{III} \quad (1)$$

with

$$\mathcal{L}^{(2)} = \frac{1}{4} f^2 \langle D_\mu U D^\mu U^\dagger + \chi U^\dagger + \chi^\dagger U \rangle \quad (2)$$

$$\mathcal{L}_{III} = -\frac{1}{4} \langle V_{\mu\nu} V^{\mu\nu} \rangle + \frac{1}{2} M_V^2 \langle [V_\mu - \frac{i}{g} \Gamma_\mu]^2 \rangle, \quad (3)$$

where  $\langle \dots \rangle$  represents a trace over  $SU(3)$  matrices. The covariant derivative is defined by

$$D_\mu U = \partial_\mu U - ieQ A_\mu U + ieU Q A_\mu, \quad (4)$$

with  $Q = \text{diag}(2, -1, -1)/3$ ,  $e = -|e|$  the electron charge, and  $A_\mu$  the photon field. The chiral matrix  $U$  is given by

$$U = e^{i\sqrt{2}\phi/f} \quad (5)$$

$$\phi \equiv \begin{pmatrix} \frac{1}{\sqrt{2}}\pi^0 + \frac{1}{\sqrt{6}}\eta_8 & \pi^+ & K^+ \\ \pi^- & -\frac{1}{\sqrt{2}}\pi^0 + \frac{1}{\sqrt{6}}\eta_8 & K^0 \\ K^- & \bar{K}^0 & -\frac{2}{\sqrt{6}}\eta_8 \end{pmatrix}, \quad V_\mu \equiv \begin{pmatrix} \frac{1}{\sqrt{2}}\rho^0 + \frac{1}{\sqrt{2}}\omega & \rho^+ & K^{*+} \\ \rho^- & -\frac{1}{\sqrt{2}}\rho^0 + \frac{1}{\sqrt{2}}\omega & K^{*0} \\ K^{*-} & \bar{K}^{*0} & \phi \end{pmatrix}_\mu. \quad (6)$$

In  $\mathcal{L}_{III}$ ,  $V_{\mu\nu}$  is defined as

$$V_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu - ig[V_\mu, V_\nu] \quad (9)$$

and

$$\Gamma_\mu = \frac{1}{2}[u^\dagger(\partial_\mu - ieQA_\mu)u + u(\partial_\mu - ieQA_\mu)u^\dagger] \quad (10)$$

with  $u^2 = U$ . The hidden gauge coupling constant  $g$  is related to  $f$  and the vector meson mass ( $M_V$ ) through

$$g = \frac{M_V}{2f}, \quad (11)$$




$$\mathcal{L}_{V\gamma} = -M_V^2 \frac{e}{g} A_\mu \langle V^\mu Q \rangle$$

$$\mathcal{L}_{V\gamma PP} = e \frac{M_V^2}{4gf^2} A_\mu \langle V^\mu (Q\phi^2 + \phi^2 Q - 2\phi Q\phi) \rangle$$

$$\mathcal{L}_{VPP} = -i \frac{M_V^2}{4gf^2} \langle V^\mu [\phi, \partial_\mu \phi] \rangle$$



$$\mathcal{L}_{III}^{(c)} = \frac{g^2}{2} \langle V_\mu V_\nu V^\mu V^\nu - V_\nu V_\mu V^\mu V^\nu \rangle ,$$

$$\mathcal{L}_{III}^{(3V)} = ig \langle (\partial_\mu V_\nu - \partial_\nu V_\mu) V^\mu V^\nu \rangle ,$$



# Rho-rho interaction in the hidden gauge approach

R.Molina, D. Nicmorus, E. O. PRD (08)



$$\mathcal{P}^{(0)} = \frac{1}{3} \epsilon_\mu \epsilon^\mu \epsilon_\nu \epsilon^\nu$$


$$\mathcal{P}^{(1)} = \frac{1}{2} (\epsilon_\mu \epsilon_\nu \epsilon^\mu \epsilon^\nu - \epsilon_\mu \epsilon_\nu \epsilon^\nu \epsilon^\mu)$$

$$\mathcal{P}^{(2)} = \left\{ \frac{1}{2} (\epsilon_\mu \epsilon_\nu \epsilon^\mu \epsilon^\nu + \epsilon_\mu \epsilon_\nu \epsilon^\nu \epsilon^\mu) - \frac{1}{3} \epsilon_\alpha \epsilon^\alpha \epsilon_\beta \epsilon^\beta \right\}$$


Spin projectors neglecting  $q/M_V$ ,  
in  $L=0$

Bethe Salpeter eqn.  $T = \frac{V}{1 - VG}$  G is the pp propagator

Squared amplitude for S=2 and  $q_{\max}=875$  MeV




Squared amplitude for S=2 and  $q_{\max}=1000$  MeV




Two  $l=0$  states generated  $f_0, f_2$  that we associate to  $f_0(1370)$  and  $f_2(1270)$

Squared amplitude for S=0 and  $q_{\max}=875$  MeV




Squared amplitude for S=0 and  $q_{\max}=1000$  MeV



Belle finds the  $f_0(1370)$  around 1470 MeV

Attraction found in many channels



The  $f_2(1270)$  is not changed by the addition of new channels, but a new resonance appears can be associated to  $f'_2(1525)$

Exp :  $\Gamma(f_2(1270)) = 185$  MeV

Exp:  $\Gamma(f'_2(1525)) = 76$  MeV


# Predicted meson states from V V interaction

$I^G(J^{PC})$	M , $\Gamma$ [MeV] Theory			PDG data		
	Pole position	Real axis		Name	Mass	Width
		$\Lambda_b = 1.4$ GeV	$\Lambda_b = 1.5$ GeV			
$0^+(0^{++})$	(1512,51)	(1523,257)	(1517,396)	$f_0(1370)$	1200~1500	200~500
$0^+(0^{++})$	(1726,28)	(1721,133)	(1717,151)	$f_0(1710)$	$1724 \pm 7$	$137 \pm 8$
$0^-(1^{+-})$	(1802,78)	(1802,49)		$h_1$		
$0^+(2^{++})$	(1275,2)	(1276,97)	(1275,111)	$f_2(1270)$	$1275.1 \pm 1.2$	$185.0^{+2.9}_{-2.4}$
$0^+(2^{++})$	(1525,6)	(1525,45)	(1525,51)	$f'_2(1525)$	$1525 \pm 5$	$73^{+6}_{-5}$
$1^-(0^{++})$	(1780,133)	(1777,148)	(1777,172)	$a_0$		
$1^+(1^{+-})$	(1679,235)	(1703,188)		$b_1$		
$1^-(2^{++})$	(1569,32)	(1567,47)	(1566,51)	$a_2(1700)??$	<b><math>a_2(1320)</math> Nagahiro PRD 11</b>	
$1/2(0^+)$	(1643,47)	(1639,139)	(1637,162)	$K_0^*$		
$1/2(1^+)$	(1737,165)	(1743,126)		$K_1(1650)?$		
$1/2(2^+)$	(1431,1)	(1431,56)	(1431,63)	$K_2^*(1430)$	$1429 \pm 1.4$	$104 \pm 4$



# Signature of an $h_1$ state in the $J/\psi \rightarrow \eta h_1 \rightarrow \eta K^{*0} \bar{K}^{*0}$ decay

Xie Ju Jun, M. Albaladejo and E. O., 2013



$I^G(J^{PC})$

$0^-(1^{+-})$	(1802,78)	(1802,49)	$h_1$
---------------	-----------	-----------	-------



Pole positions and residues in the strangeness=0 and isospin=0 channel. All quantities are in units of MeV.

(1802, -i39) [spin=1]					
	$K^* \bar{K}^*$	$\rho\rho$	$\omega\omega$	$\omega\phi$	$\phi\phi$
$g$	(8034, -i2542)	0	0	0	0

does not go to VV because of C-parity

It cannot go to PP, because  $J=1$  requires  $L=1$  in PP  $\rightarrow$  negative parity

Thus  $K^* \bar{K}^*$  is the only open channel



$$t = v + v\tilde{G}t = v(1 + \tilde{G}t) = (1 - v\tilde{G})^{-1}v = (v^{-1} - \tilde{G})^{-1} \quad v = \left(9 + b \left(1 - \frac{3M_{\text{inv}}^2}{4m_{K^*}^2}\right)\right) g^2$$

$$t_P = V_P \left(1 + \tilde{G}(M_{\text{inv}}^2)t(M_{\text{inv}}^2)\right) = V_P \frac{t(M_{\text{inv}}^2)}{v(M_{\text{inv}}^2)} \quad g = m_\rho/2f$$

$$G = \frac{1}{16\pi^2} \left( \alpha + \text{Log} \frac{m_1^2}{\mu^2} + \frac{m_2^2 - m_1^2 + s}{2s} \text{Log} \frac{m_2^2}{m_1^2} \right. \\ \left. + \frac{p}{\sqrt{s}} \left( \text{Log} \frac{s - m_2^2 + m_1^2 + 2p\sqrt{s}}{-s + m_2^2 - m_1^2 + 2p\sqrt{s}} + \text{Log} \frac{s + m_2^2 - m_1^2 + 2p\sqrt{s}}{-s - m_2^2 + m_1^2 + 2p\sqrt{s}} \right) \right)$$

$\mu = 1000 \text{ MeV}$   
 $a(\mu) = \alpha$

$$\frac{d\Gamma}{dM_{\text{inv}}} = \frac{C}{|v(M_{\text{inv}}^2)|^2} \frac{p_1 \tilde{p}_2}{M_{J/\psi}} |t(M_{\text{inv}}^2)|^2$$

A fit to data is made changing  $a(\mu)$

$$M_{h_1} = 1830 \pm 20 \text{ MeV} \text{ and } \Gamma_{h_1} = 110 \pm 10 \text{ MeV}$$



# Extension to the baryon sector

$$\mathcal{L}_{BBV} = -\frac{g}{2\sqrt{2}} (tr(\bar{B}\gamma_\mu[V^\mu, B]) + tr(\bar{B}\gamma_\mu B)tr(V^\mu))$$

Vector propagator  $1/(q^2-M_V^2)$

In the approximation  $q^2/M_V^2=0$  one recovers the chiral Lagrangians Weinberg-Tomozawa term. For consistency, for vectors we take  $\vec{q}/M_V=0$



> 200 works



A. Ramos, E. O. EPJA 10

Khemchandani, Hosaka, Kaneko, Martinez, Nagahiro, PRD 11

Kolomeitsev et al  
Sarkar et al



J. Vijande, P. Gonzalez. E.O  
PRC,2009

Sarkar, Vicente Vacas, B.X.Sun,  
E.O, EPJA 10

# Vector octet – baryon octet interaction

$$\begin{aligned}\mathcal{L}_{III}^{(3V)} &= ig \langle V^\nu \partial_\mu V_\nu V^\mu - \partial_\nu V_\mu V^\mu V^\nu \rangle \\ &= ig \langle V^\mu \partial_\nu V_\mu V^\nu - \partial_\nu V_\mu V^\mu V^\nu \rangle \\ &= ig \langle (V^\mu \partial_\nu V_\mu - \partial_\nu V_\mu V^\mu) V^\nu \rangle ,\end{aligned}$$

$$\mathcal{L}_{VPP} = -ig \operatorname{tr} ([P, \partial_\mu P] V^\mu) \quad \downarrow \quad B = \begin{pmatrix} \frac{1}{\sqrt{2}}\Sigma^0 + \frac{1}{\sqrt{6}}\Lambda & \Sigma^+ & p \\ \Sigma^- & -\frac{1}{\sqrt{2}}\Sigma^0 + \frac{1}{\sqrt{6}}\Lambda & n \\ \Xi^- & \Xi^0 & -\frac{2}{\sqrt{6}}\Lambda \end{pmatrix}$$

$V^\nu$  cannot correspond to an external vector.

Indeed, external vectors have only spatial components in the approximation of neglecting three momenta,  $\varepsilon^0 = k/M$  for longitudinal vectors,  $\varepsilon^0 = 0$  for transverse vectors. Then  $\partial_\nu$  becomes three momentum which is neglected.  $\rightarrow$

$V^\nu$  corresponds to the exchanged vector.  $\rightarrow$  complete analogy to VPP

Extra  $\varepsilon_\mu \varepsilon^\mu = -\varepsilon_i \varepsilon_i$  but the interaction is formally identical to the case of PB  $\rightarrow$  PB

In the same approximation only  $\gamma^0$  is kept for the baryons  $\rightarrow$  the spin dependence is only  $\varepsilon_i \varepsilon_i$  and the states are degenerate in spin 1/2 and 3/2

$$V_{ij} = -C_{ij} \frac{1}{4f^2} (k^0 + k'^0) \vec{\varepsilon} \vec{\varepsilon}'$$

$K^0$  energy of vector mesons

We solve the Bethe Salpeter equation in coupled channels Vector-Baryon octet.

$$T = (1 - GV)^{-1} V$$


with G the loop function of vector-baryon

Apart from the peaks, poles are searched in the second Riemann sheet and pole positions and residues are determined.

The G function takes into account the mass distribution of the vectors (width).


$$\begin{aligned} G_{(P,B)} &= i2M_B \int \frac{d^4q}{(2\pi)^4} \frac{1}{(P-q)^2 - M_B^2 + i\varepsilon} \frac{1}{q^2 - M_P^2 + i\varepsilon}, \\ &= \frac{2M_B}{16\pi^2} \left\{ a_\mu + \ln \frac{M_B^2}{\mu^2} + \frac{M_P^2 - M_B^2 + s}{2s} \ln \frac{M_P^2}{M_B^2} \right. \\ &\quad + \frac{\bar{q}}{\sqrt{s}} [\ln(s - (M_B^2 - M_P^2) + 2\bar{q}\sqrt{s}) + \ln(s + (M_B^2 - M_P^2) + 2\bar{q}\sqrt{s}) \\ &\quad \left. - \ln(-s - (M_B^2 - M_P^2) + 2\bar{q}\sqrt{s}) - \ln(-s + (M_B^2 - M_P^2) + 2\bar{q}\sqrt{s})] \right\} \end{aligned}$$

$a_\mu$  is a subtraction constant that regularizes the loop




$S, I$	Theory		PDG data				
	(real axis)		name	$J^P$	status	mass	width
	mass	width					
$0, 1/2$	1699	84	$N(1650)$	$1/2^-$	***	1645-1670	145-185
			$N(1700)$	$3/2^-$	***	1650-1750	50-150
	1967	82	$N(2080)$	$3/2^-$	**	$\approx 2080$	180-450
			$N(2090)$	$1/2^-$	*	$\approx 2090$	100-400
$-1, 0$	1783	8	$\Lambda(1690)$	$3/2^-$	***	1685-1695	50-70
			$\Lambda(1800)$	$3/2^-$	***	1720-1850	200-400
	1900	54	$\Lambda(2000)$	??	*	$\approx 2000$	73-240
	2158	20					
$-1, 1$	1830	44	$\Sigma(1750)$	$1/2^-$	***	1730-1800	60-160
	1985	244	$\Sigma(1940)$	$3/2^-$	***	1900-1950	150-300
			$\Sigma(2000)$	$1/2^-$	*	$\approx 2000$	100-450
$-2, 1/2$	2030	52	$\Xi(2030)$	??	***	$2025 \pm 5$	$21 \pm 6$
	2080	24	$\Xi(2120)$	??	*	$\approx 2120$	25

Table 1: The properties of the 9 dynamically generated resonances and their possible PDG counterparts.



Mixing of VB and PB channels



Note the degeneracy between  $J=1/2, 3/2$  is broken. The  $J=1/2$  moves to 1650 MeV, while  $J=3/2$  stays at 1700 MeV, as in experiment

## Other works that consider the mixing between PB and VB

Romanets, Tolos, Garcia Recio, Nieves, Salcedo, Timmermanns, 2012  
SU(6), SU(8) spin symmetry

Khemchandani, Martinez, Nagahiro, Kaneko, Hosaka, 2011  
Gauge terms from anomalous coupling of vector mesons to baryons


Kolomeitsev, Lutz some work in  $3/2^-$  sector

.....



# The role of vector-baryon channels and resonances in the $\gamma p \rightarrow K^0 \Sigma^+$ and $\gamma n \rightarrow K^0 \Sigma^0$ reactions near the $K^* \Lambda$ threshold.

A. Ramos and E. Oset, Phys. Lett B 2013




CBELSA/TAPS  
R. Ewald.....  
H. Schmieden.....  
PLB 2012

Sudden drop of cross section around  $K^* \Lambda$  threshold  
Angular dependence also becomes flat around this energy  
Hints at an important role of vector baryon interaction in  $L=0$

Standard models MAID, SAID fail badly to reproduce these features

We use the tools of the local hidden gauge formalism to describe the reaction




Tree level is null with final  $K^0$

$$\mathcal{L}_{PBB} = \frac{1}{2}(D + F)\langle \bar{B}\gamma^\mu\gamma^5 u_\mu B \rangle + \frac{1}{2}(D - F)\langle \bar{B}\gamma^\mu\gamma^5 B u_\mu \rangle$$


$$\gamma^\mu\gamma^5 u_\mu \rightarrow \frac{\sqrt{2}}{f}\sigma^i\partial_i\phi$$

$$-it_{\gamma N \rightarrow K^0 \Sigma}^{\pi\text{-pole}} = e \sum_{V=\rho^0, \omega, \phi} \mathcal{C}_{\gamma V} \sum_{V'B'} t_{VN \rightarrow V'B'} i \int \frac{d^4 q}{(2\pi)^4} \frac{1}{(q+k)^2 - M_{V'}^2 + i\varepsilon} \frac{1}{q^2 - m_\pi^2 + i\varepsilon} \frac{M_{B'}}{E_{B'}} \frac{1}{P^0 - q^0 - k^0 - E_{B'}(\vec{q} + \vec{k}) + i\varepsilon} (\vec{q} - \vec{k}) \vec{\epsilon}_\gamma \vec{\sigma} \vec{q} V_{Y, B'} F(q),$$

$$\mathcal{C}_{\gamma V} = \begin{cases} \frac{1}{\sqrt{2}} & \text{for } V = \rho \\ \frac{1}{3\sqrt{2}} & \text{for } V = \omega \\ -\frac{1}{3} & \text{for } V = \phi \end{cases}$$




$$\begin{aligned}
 -it_{\gamma N \rightarrow K^0 \Sigma}^{KR} &= e \sum_{V=\rho^0, \omega, \phi} \mathcal{C}_{\gamma V} \sum_{V' B'} t_{V N \rightarrow V' B'} i \int \frac{d^4 q}{(2\pi)^4} \frac{1}{(q+k)^2 - M_{V'}^2 + i\epsilon} \\
 &\quad \frac{M_{B'}}{E_{B'}} \frac{1}{P^0 - q^0 - k^0 - E_{B'}(\vec{q} + \vec{k}) + i\epsilon} \vec{\sigma} \vec{\epsilon}_\gamma V_{Y, B'} F(q) .
 \end{aligned}$$



(a)

+



(b)

$$\frac{d\sigma_{\gamma N \rightarrow K^0 \Sigma}}{d\Omega} = \frac{1}{16\pi^2} \frac{M_N M_\Sigma}{s} \frac{k}{p} \overline{\sum} \sum |t_{\gamma N \rightarrow K^0 \Sigma}^{KR}|^2 ,$$


$$\overline{\sum} \sum |t_{\gamma N \rightarrow K^0 \Sigma}^{KR}|^2 = \frac{1}{2} \left\{ \left[ |A|^2 \vec{k}^2 + 2 \operatorname{Re}(AB^*) \right] \vec{k}^2 \sin^2 \theta + 2 |B|^2 \right\}$$

A term comes from  $\pi$ -exchange. B term from Kroll Ruderman plus part of  $\pi$ -exchange.

Angular dependence is symmetrical with respect to  $\pi/2$ .

Cancellations between A and B parts weaken angular dependence

Background will change a bit the main behaviour.





**Spectacular difference:**  
 In  $\gamma p$  first loop contains  $K^{*+} \Sigma^0$   
 and  $K^{*+} \Lambda \rightarrow$  interfere destructively.

In  $\gamma n$  first loop only contains  $K^{*+} \Sigma^-$   
 and there is no interference  $\rightarrow$



This is the effect found by  
 Doring and Nakayama, PLB 2010  
 In  $\gamma p \rightarrow \eta p$  and  $\gamma n \rightarrow \eta n$  to interpret  
 second peak in  $\gamma n \rightarrow \eta n$

FIG. 4: Cross section for the  $\gamma p \rightarrow K^0 \Sigma^+$  (black line) and  $\gamma n \rightarrow K^0 \Sigma^0$  reactions, including only an intermediate  $K^{*+} \Sigma$  channel before the transition to the final  $K^0 \Sigma$  state.

**But we have a second loop for VB  $\rightarrow$  PB transition  $\rightarrow$  extra interference**




## Fine tuning to experiment changing a bit the subtraction constants



With the new parameters we solve again the BS equation and the resonance moves from  $M_R = 1972 \text{ MeV}$ ,  $\Gamma = 64 \text{ MeV} \rightarrow M_R = 2035 \text{ MeV}$ ,  $\Gamma = 125 \text{ MeV}$

Prediction for a resonance  $1/2^-$ , or  $3/2^-$  (degenerate in our model)

Important because resonances around this energy have been removed in the latest edition of the PDG.




Conclusion: Physical explanation of basic features of the reaction

Prediction of a new resonance around 2035 MeV

Importance of VB interaction and coupled channels

Prediction of spectacular differences in the shape of  $\sigma$  for  $\gamma n \rightarrow K^0 \Sigma^0$

J.J. Wu, R. Molina, E. O. and B. S. Zou, Phys Rev Lett 2010



$$V_{ab}(P_1 B_1 \rightarrow P_2 B_2) = \frac{C_{ab}}{4f^2} (E_{P_1} + E_{P_2}),$$

$$V_{ab}(V_1 B_1 \rightarrow V_2 B_2) = \frac{C_{ab}}{4f^2} (E_{V_1} + E_{V_2}) \vec{\epsilon}_1 \cdot \vec{\epsilon}_2$$

$$T = [1 - VG]^{-1}V$$

TABLE I: Coefficients  $C_{ab}$  in Eq. (2) for  $(I, S) = (1/2, 0)$

	$\bar{D}\Sigma_c$	$\bar{D}\Lambda_c^+$	$\eta_c N$	$\pi N$	$\eta N$	$\eta' N$	$K\Sigma$	$K\Lambda$
$\bar{D}\Sigma_c$	-1	0	$-\sqrt{3/2}$	-1/2	$-1/\sqrt{2}$	1/2	1	0
$\bar{D}\Lambda_c^+$		1	$\sqrt{3/2}$	-3/2	$1/\sqrt{2}$	-1/2	0	1

TABLE II: Coefficients  $C_{ab}$  in Eq. (2) for  $(I, S) = (0, -1)$

	$\bar{D}_s\Lambda_c^+$	$\bar{D}\Xi_c$	$\bar{D}\Xi'_c$	$\eta_c\Lambda$	$\pi\Sigma$	$\eta\Lambda$	$\eta'\Lambda$	$\bar{K}N$	$K\Xi$
$\bar{D}_s\Lambda_c^+$	0	$-\sqrt{2}$	0	1	0	$\sqrt{1/3}$	$\sqrt{2/3}$	$-\sqrt{3}$	0
$\bar{D}\Xi_c$		-1	0	$\sqrt{1/2}$	$-\frac{3}{2}$	$\sqrt{1/6}$	$-\sqrt{1/12}$	0	$\sqrt{3/2}$
$\bar{D}\Xi'_c$			-1	$-\sqrt{3/2}$	$\sqrt{3/4}$	$-\sqrt{1/2}$	$1/2$	0	$\sqrt{1/2}$
$\eta_c\Lambda$				0	0	0	0	0	0



$(I, S)$	$z_R$ (MeV)	$g_a$		
$(1/2, 0)$		$\bar{D}\Sigma_c$	$\bar{D}\Lambda_c^+$	
	4269	2.85	0	
$(0, -1)$		$\bar{D}_s\Lambda_c^+$	$\bar{D}\Xi_c$	$\bar{D}\Xi'_c$
	4213	1.37	3.25	0
	4403	0	0	2.64

TABLE III: Pole positions  $z_R$  and coupling constants  $g_a$  for the states from  $PB \rightarrow PB$ .

$(I, S)$	$z_R$ (MeV)	$g_a$		
$(1/2, 0)$		$\bar{D}^*\Sigma_c$	$\bar{D}^*\Lambda_c^+$	
	4418	2.75	0	
$(0, -1)$		$\bar{D}_s^*\Lambda_c^+$	$\bar{D}^*\Xi_c$	$\bar{D}^*\Xi'_c$
	4370	1.23	3.14	0
	4550	0	0	2.53

TABLE IV: Pole position and coupling constants for the bound states from  $VB \rightarrow VB$ .

$(I, S)$	$M$	$\Gamma$	$\Gamma_i$											
$(1/2, 0)$			$\pi N$	$\eta N$	$\eta' N$	$K\Sigma$	$\eta_c N$							
	4261	56.9	3.8	8.1	3.9	17.0	23.4							
$(0, -1)$			$KN$	$\pi\Sigma$	$\eta\Lambda$	$\eta'\Lambda$	$K\Xi$	$\eta_c\Lambda$						
	4209	32.4	15.8	2.9	3.2	1.7	2.4	5.8						
	4394	43.3	0	10.6	7.1	3.3	5.8	16.3						
$(I, S)$	$M$	$\Gamma$	$\Gamma_i$											
$(1/2, 0)$			$\rho N$	$\omega N$	$K^*\Sigma$									
	4412	47.3	3.2	10.4	13.7								19.2	
$(0, -1)$			$K^*N$	$\rho\Sigma$	$\omega\Lambda$	$\phi\Lambda$	$K^*\Xi$	$J/\psi\Lambda$						
	4368	28.0	13.9	3.1	0.3	4.0	1.8	5.4						
	4544	36.6	0	8.8	9.1	0	5.0	13.8						

# Prediction of super-heavy $N^*$ and $\Lambda^*$ resonances with hidden beauty

J. J. Wu, L. Zhao and B.S. Zou, Phys Lett B 2012

Pole positions  $z_R$  and coupling constants  $g_\alpha$  for the states in  $(I, S) = (1/2, 0)$  sector

$z_R$ (MeV)	$g_\alpha$	
	$B\Sigma_b$	$B\Lambda_b$
11052	2.05	0
$z_R$ (MeV)	$g_\alpha$	
	$B^*\Sigma_b$	$B^*\Lambda_b$
11100	2.02	0

Pole positions  $z_R$  and coupling constants  $g_\alpha$  for the states in  $(I, S) = (0, -1)$  sector

$z_R$ (MeV)	$g_\alpha$		
	$B_s\Lambda_b$	$B\Xi_b$	$B\Xi'_b$
11021 - 0.59i	0.14 - 0.11i	2.27 + 0.004i	0
11191	0	0	1.92
$z_R$ (MeV)	$g_\alpha$		
	$B_s^*\Lambda_b$	$B^*\Xi_b$	$B^*\Xi'_b$
11069 - 0.59i	0.14 - 0.12i	2.24 + 0.005i	0
11238	0	0	1.89

The search for molecules in the charm and beauty sectors has experienced a recent boom.

Nieves, Hidalgo, Pavon, Guo, Garcia, Salcedo, Romanets, Tolos, Ramos implementing heavy quark spin symmetry (HQSS)

S. L. Zhu with dynamics of meson exchange

Fernandez –Carames et al, Gutsche et al, Ding et al, Bondar et al., Cleven et al., Li et al,


Guo, Hanhart, Meissner, Sun, Liu, Zhu, Wang, Zhao, Dong, Zhang ....

Xiao, Nieves, Ozpineci, E. O. local hidden gauge (exchange of vector mesons) with HQSS.

# Heavy quark spin symmetry (HQSS) and the local hidden gauge approach (LHGA):

- 1) Dominant term of LHGA comes from exchange of light vector mesons  
→ The heavy quarks are spectators → interaction is independent of spin and flavour of heavy quarks (HQSS)  
→ The LHGA automatically implements HQSS
- 2) Take  $I=1$  in meson-meson as in figure: the light exchange involves  $u$   $u$ bar from upper vertex and  $d$   $d$ bar from lower vertex → OZI forbidden:  
 $\rho$  and  $\omega$  exchange cancel.  
But so does pseudoscalar exchange if masses of nonet are taken equal.

$I=1$  becomes subdominant



# Conclusions

Chiral dynamics or its extension with the LHGA is a good tool to deal with hadron interaction.

Its combination with nonperturbative unitary techniques allows to study the interaction of hadrons. Poles in amplitudes correspond to dynamically generated resonances. Many known resonances can be described in this way.

The interaction of vector mesons with other mesons or baryons plays an important role in many hadronic reactions.

We analyzed a recent BES reaction on  $J/\psi \rightarrow \eta K^{*0} \bar{K}^{*0}$  and interpreted it as showing evidence for a new  $h_1$  state around 1830 MeV predicted from the VV interaction.

The dynamics of vector baryon interaction allowed us to interpret the experimental results of the  $\gamma p \rightarrow K^0 \Sigma^+$  reaction and make interesting and unexpected predictions for  $\gamma n \rightarrow K^0 \Sigma^0$ .

Plus the prediction of a  $N^* \ 1/2^-, 3/2^-$  resonance around 2035 MeV.

Extension to the heavy quark sector is proving fruitful. More data beyond spectra needed.