

A test of CPT symmetry in K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ decays^{*}

XING Zhi-Zhong(邢志忠)^{1,2;1)}

¹ Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

² Theoretical Physics Center for Science Facilities, Chinese Academy of Sciences, Beijing 100049, China

Abstract I show that the CP -violating asymmetry in K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ decays differs from that in $K_L \rightarrow \pi^+\pi^-$, $K_L \rightarrow \pi^0\pi^0$ or the semileptonic K_L transitions, if there exists CPT violation in K^0 - \bar{K}^0 mixing. A delicate measurement of this difference at a super flavor factory (e.g., the ϕ factory) will provide us with a robust test of CPT symmetry in the neutral kaon system.

Key words K^0 - \bar{K}^0 mixing, CPT violation

PACS 11.30.Er, 13.25.Es, 14.40.Aq

1 Introduction

The CPT theorem claims that a Lorentz-invariant local quantum field theory with a Hermitian Hamiltonian must have CPT symmetry [1]. It is so far so good, because there is no convincing experimental hint at CPT violation [2]. The breaking of CPT symmetry, as expected in some “exotic” scenarios of new physics beyond the standard model (e.g., string theory) [3], would be a big deal. In any case, much more experimental tests of this theorem are desirable.

The K^0 - \bar{K}^0 mixing system has been playing an important role in particle physics for testing fundamental symmetries (such as CP , T and CPT) and examining conservation laws (such as $\Delta S = \Delta Q$). The existing experimental evidence for CPT invariance in the mixing and decays of neutral kaon mesons remains rather poor [2]: it is not excluded that the strength of CPT -violating interactions could be as large as about ten percentage of that of CP -violating interactions. This unsatisfactory situation will be improved in the near future, in particular after a variety of more delicate measurements are carried out at a super flavor factory [4] (e.g., the ϕ factory [5]).

There are several possibilities of probing CPT violation in K^0 - \bar{K}^0 mixing with the decays of K_S and K_L mesons into the two-pion and (or) the semileptonic states [2]. A different approach towards testing

CPT symmetry, with the help of neutral kaon decays into the three-pion states, has also been pointed out in Ref. [6]. The idea is simply that the CP -violating effect induced by K^0 - \bar{K}^0 mixing in K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ transitions should not be identical to that in $K_L \rightarrow \pi^+\pi^-$, $K_L \rightarrow \pi^0\pi^0$ or the semileptonic K_L decays, if CPT symmetry is broken. Thus a careful comparison between these two types of CP -violating effects may provide us with a robust test of CPT invariance in K^0 - \bar{K}^0 mixing.

An unfortunate fact is that no attention has so far been paid to the method advocated in Ref. [6]. In this talk, which is more or less an advertisement, I shall explain why a test of CPT symmetry is possible by measuring the time-dependent CP -violating asymmetry between $K^0(t) \rightarrow \pi^+\pi^-\pi^0$ and $\bar{K}^0(t) \rightarrow \pi^+\pi^-\pi^0$ decays. My result is hopefully useful for the upcoming experiments of kaon physics.

2 The idea

Let me outline the main idea. The mass eigenstates of K^0 and \bar{K}^0 can in general be written as

$$\begin{aligned}
 |K_S\rangle &= \frac{1}{\sqrt{|p_1|^2 + |q_1|^2}} (p_1|K^0\rangle + q_1|\bar{K}^0\rangle), \\
 |K_L\rangle &= \frac{1}{\sqrt{|p_2|^2 + |q_2|^2}} (p_2|K^0\rangle - q_2|\bar{K}^0\rangle), \quad (1)
 \end{aligned}$$

Received 25 January 2010

^{*} Supported by National Natural Science Foundation of China (10425522, 10875131) and Ministry of Science and Technology of China (2009CB825207)

1) E-mail: xingzz@ihep.ac.cn

©2009 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

in which p_i and q_i (for $i = 1, 2$) are complex mixing parameters. Note that $p_1 = p_2$ and $q_1 = q_2$ follow from CPT invariance [7]. The traditional characteristic quantities of CP violation in the K^0 - \bar{K}^0 mixing system [2], η_{+-} , η_{00} and δ_L , are all related to K_L decays and thus the (p_2, q_2) parameters. For example,

$$\delta_L \equiv \frac{|p_2|^2 - |q_2|^2}{|p_2|^2 + |q_2|^2} \quad (2)$$

in the absence of $\Delta S = -\Delta Q$ interactions. A measurement of CP violation associated with

$$\delta_S = \frac{|p_1|^2 - |q_1|^2}{|p_1|^2 + |q_1|^2} \quad (3)$$

has been assumed to be extremely difficult, if not impossible, due to the rapid decay of the K_S meson to the two-pion state or the semileptonic state. Nevertheless, I shall show that δ_S can be measured from the rate asymmetry of K^0 and \bar{K}^0 mesons decaying into the three-pion state $\pi^+\pi^-\pi^0$. The difference between δ_S and δ_L signifies CPT violation in K^0 - \bar{K}^0 mixing. This point can be seen more clearly if one adopts the popular (ϵ, δ) parameters to describe CP - and CPT -violating effects in the K^0 - \bar{K}^0 mixing system [2]:

$$\begin{aligned} p_1 &= 1 + \epsilon + \delta, \\ p_2 &= 1 + \epsilon - \delta, \\ q_1 &= 1 - \epsilon - \delta, \\ q_2 &= 1 - \epsilon + \delta. \end{aligned} \quad (4)$$

Then

$$\begin{aligned} \delta_L &= 2(\text{Re}\epsilon - \text{Re}\delta), \\ \delta_S &= 2(\text{Re}\epsilon + \text{Re}\delta). \end{aligned} \quad (5)$$

It turns out that $\delta_S - \delta_L = 4\text{Re}\delta$ is a clear signature of CPT violation [6].

Let me quote two typical experimental constraints on the CPT -violating parameter δ in K^0 - \bar{K}^0 mixing: $\text{Re}\delta = (2.9 \pm 2.6_{\text{stat}} \pm 0.6_{\text{sys}}) \times 10^{-4}$ obtained by the CPLEAR Collaboration [8] and $\text{Im}\delta = (0.4 \pm 2.1) \times 10^{-5}$ obtained by the KLOE Collaboration [9]. A systematic analysis of the CP - and CPT -violating parameter space has already been done by the Particle Data Group in Ref. [2].

3 The approach

The CP eigenvalue for the $\pi^+\pi^-\pi^0$ final state is given by $(-1)^{l+1}$, where l is the relative angular momentum between π^+ and π^- . Since the sum of the masses of three pions is close to the kaon mass, the pions have quite low kinetic energy $E_{\text{CM}}(\pi)$ in

the kaon rest-frame, and the states with $l > 0$ are suppressed by the centrifugal barrier [10]. Thus the K_L meson decays dominantly into the kinematics-favored ($l = 0$) and CP -allowed ($CP = -1$) $\pi^+\pi^-\pi^0$ state. The decay amplitude of $K_S \rightarrow \pi^+\pi^-\pi^0$ consists of both the kinematics-suppressed ($l = 1$) but CP -allowed ($CP = +1$) component, and the kinematics-favored ($l = 0$) but CP -forbidden ($CP = -1$) component. This implies an interesting Dalitz-plot distribution for the $K_S \rightarrow \pi^+\pi^-\pi^0$ transition: it is symmetric with respect to π^+ and π^- for the CP -violating amplitude, but anti-symmetric for the CP -conserving amplitude. Let the ratio of K_S and K_L decay amplitudes be

$$\eta_{+-0} = \frac{A(K_S \rightarrow \pi^+\pi^-\pi^0)}{A(K_L \rightarrow \pi^+\pi^-\pi^0)}. \quad (6)$$

It is clear that η_{+-0} depends only upon the CP -violating component of $A(K_S \rightarrow \pi^+\pi^-\pi^0)$, if data are integrated over the whole Dalitz plot [10, 11]. The time-dependent rates for the initially pure K^0 and \bar{K}^0 states decaying into $\pi^+\pi^-\pi^0$, denoted by $\mathcal{R}(t)$ and $\bar{\mathcal{R}}(t)$ respectively, can be calculated with the help of Eqs. (1) and (6). I arrive at [6]

$$\begin{aligned} \mathcal{R}(t) &\propto \left[|q_1|^2 + |q_2|^2 |\eta_{+-0}|^2 e^{-\Delta\Gamma t} + \right. \\ &\quad \left. 2\text{Re}(q_1^* q_2 \eta_{+-0} e^{i\Delta m t}) e^{-\Delta\Gamma t/2} \right], \\ \bar{\mathcal{R}}(t) &\propto \left[|p_1|^2 + |p_2|^2 |\eta_{+-0}|^2 e^{-\Delta\Gamma t} - \right. \\ &\quad \left. 2\text{Re}(p_1^* p_2 \eta_{+-0} e^{i\Delta m t}) e^{-\Delta\Gamma t/2} \right], \end{aligned} \quad (7)$$

where $\Delta m > 0$ and $\Delta\Gamma > 0$ denote the mass difference and the width difference of K_S and K_L mesons, respectively. To a good degree of accuracy, I obtain the following CP -violating asymmetry:

$$\begin{aligned} \mathcal{A}(t) &\equiv \frac{\bar{\mathcal{R}}(t) - \mathcal{R}(t)}{\bar{\mathcal{R}}(t) + \mathcal{R}(t)} = \delta_S - \\ &2e^{-\Delta\Gamma t/2} [\text{Re}\eta_{+-0} \cos(\Delta m t) - \text{Im}\eta_{+-0} \sin(\Delta m t)] \xi - \\ &2e^{-\Delta\Gamma t/2} [\text{Re}\eta_{+-0} \sin(\Delta m t) + \text{Im}\eta_{+-0} \cos(\Delta m t)] \zeta, \end{aligned} \quad (8)$$

in which

$$\begin{aligned} \xi &= \frac{\text{Re}(p_1 p_2^* + q_1 q_2^*)}{|p_1|^2 + |q_1|^2} = \\ &1 + \mathcal{O}(|\epsilon|^2) + \mathcal{O}(|\delta|^2) + \mathcal{O}(\text{Re}(\epsilon\delta^*)), \\ \zeta &= \frac{\text{Im}(p_1 p_2^* + q_1 q_2^*)}{|p_1|^2 + |q_1|^2} = \mathcal{O}(\text{Im}(\epsilon\delta^*)). \end{aligned} \quad (9)$$

It is obvious that δ_S can be determined through the measurement of $\mathcal{A}(t)$. In particular, the relationship $\lim_{t \rightarrow \infty} \mathcal{A}(t) = \delta_S$ holds.

As I have emphasized, the difference between δ_S and δ_L hints at CPT violation in K^0 - \bar{K}^0 mixing. If $|\text{Re}\delta|/\text{Re}\epsilon \sim 0.1$, then the difference $\delta_S - \delta_L = 4\text{Re}\delta$ can be as large as $0.4\text{Re}\epsilon \sim 6.6 \times 10^{-4}$ in magnitude, where the experimental value $\text{Re}\epsilon \approx 1.65 \times 10^{-3}$ has been used [2]. Since both ϵ and δ are small quantities, it turns out that $\xi \approx 1$ and $\zeta \approx 0$ are good approximations. Eq. (8) is therefore simplified to

$$\mathcal{A}(t) = \delta_S - 2e^{-\Delta\Gamma t/2} \left[\text{Re}\eta_{+-0} \cos(\Delta mt) - \text{Im}\eta_{+-0} \sin(\Delta mt) \right]. \quad (10)$$

In the neglect of CPT violation, namely, $\delta_S = 2\text{Re}\epsilon$, Eq. (10) can simply reproduce the result obtained in Ref. [10]. For illustration, I plot the behavior of $\mathcal{A}(t)$ in Fig. 1, in which $\delta_S = 3 \times 10^{-3}$ and $|\eta_{+-0}| = 5 \times 10^{-3}$ have typically been input. One may observe that $\mathcal{A}(t)$ approaches δ_S for $t \geq 5\tau_S$ and reaches δ_S if $t \geq 10\tau_S$, where τ_S is the mean lifetime of the K_S meson. This implies a certain feasibility to determine δ_S from the time-dependent CP -violating asymmetry $\mathcal{A}(t)$.




Fig. 1. An illustrative plot for the CP -violating asymmetry $\mathcal{A}(t)$ with the typical inputs $\delta_S = 3 \times 10^{-3}$ and $|\eta_{+-0}| = 5 \times 10^{-3}$ [6].

4 The discussion

In the above analysis I have taken an integration over the whole Dalitz plot, such that η_{+-0} solely contains the CP -violating part of $A(K_S \rightarrow \pi^+\pi^-\pi^0)$. To look at the CP -conserving component of $A(K_S \rightarrow \pi^+\pi^-\pi^0)$, one may study the phase-space regions $E_{\text{CM}}(\pi^+) > E_{\text{CM}}(\pi^-)$ and $E_{\text{CM}}(\pi^+) < E_{\text{CM}}(\pi^-)$ separately [10]. In this case the corresponding CP -violating asymmetries between $\bar{\mathcal{R}}(t)$ and $\mathcal{R}(t)$ take the same form as $\mathcal{A}(t)$ in Eq. (8) or Eq. (10), but η_{+-0} should be replaced by $(\eta_{+-0} \pm \lambda)$, where λ denotes the CP -conserving contribution to the ratio of K_S and K_L decay amplitudes [10]. Certainly, the CP -violating parameter δ_S can still be determined from

measuring the time dependence of the relevant decay rate asymmetries.

An accurate measurement of δ_S from K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ should be feasible at the ϕ factory, where a huge amount of $K^0\bar{K}^0$ events can be coherently produced [5]. Choosing the semileptonic decay of one kaon to tag the flavor of the other kaon decaying into $\pi^+\pi^-\pi^0$ on the ϕ resonance, one should be able to construct the time-dependent rate asymmetry between $K^0(t) \rightarrow \pi^+\pi^-\pi^0$ and $\bar{K}^0(t) \rightarrow \pi^+\pi^-\pi^0$ decays in a way similar to Eq. (8). It is also expected that other super flavor factories may measure δ_S and δ_L to a good degree of accuracy.

Note that Lorentz invariance has been taken for granted in what I have discussed. As pointed out by Greenberg [12], “If CPT invariance is violated in an interacting quantum field theory, then that theory also violates Lorentz invariance”. In my discussions, the dependence of the CPT -violating parameter δ on the sidereal time should in general be considered, since CPT violation may simultaneously imply a violation of Lorentz symmetry in the neutral kaon system. For simplicity, here I take δ to be a constant by assuming that the boost parameters of both K^0 and \bar{K}^0 are small and the corresponding Lorentz-violating effect is rotationally invariant in the laboratory frame [13]. In this approximation, my results are essentially valid as the averages over the sidereal time, such that the effect of Lorentz violation due to the direction of motion is negligible.

Finally, I like to mention that different approaches have been discussed to test CPT symmetry in D^0 - \bar{D}^0 , B_d^0 - \bar{B}_d^0 or B_s^0 - \bar{B}_s^0 mixing [14]. The idea presented here cannot directly be applied to those heavy neutral-meson systems. In this sense, it represents a unique way applicable in the K^0 - \bar{K}^0 mixing system to test the CPT theorem.

5 The conclusion

To conclude, the CP -violating effect induced by K^0 - \bar{K}^0 mixing in K^0 vs $\bar{K}^0 \rightarrow \pi^+\pi^-\pi^0$ decays is possible to deviate to some extent from that in $K_L \rightarrow \pi\pi$ or the semileptonic K_L transitions due to the violation of CPT symmetry. Measuring or constraining this tiny difference may serve as a robust test of CPT invariance in the neutral kaon system.

I would like to thank Changzheng Yuan and the organizing committee of PHIPSI09 for giving me this opportunity to present an “old” idea for the future.

References

- 1 Schwinger J. Phys. Rev., 1951, **82**: 914; Lüders G. Kgl. Dansk. Vidensk. Selsk. Mat. Fys. Medd., 1954, **28**: 1; Pauli W. In: Niels Bohr and the Development of Physics. New York: Pergamon Press. 1955. 30
- 2 Particle Data Group, Amsler C et al. Phys. Lett. B, 2008, **667**: 1
- 3 Kostelecky V A. hep-ph/0104227
- 4 Hitlin D. In these proceedings
- 5 The DAΦNE Physics Handbook. Edited by Maiani L, Panzeri G, Paver N. Frascati. 1992.
- 6 XING Z Z. Phys. Rev. D, 2000, **62**: 097901
- 7 Lee T D, WU C S. Ann. Rev. Nucl. Part. Sci., 1966, **16**: 511
- 8 CPLEAR collaboration, Angelopoulos A et al. Phys. Lett. B, 1998, **444**: 52
- 9 KLOE collaboration, Ambrosino F et al. JHEP, 2006, **0612**: 011
- 10 CPLEAR collaboration, Adler R et al. Phys. Lett. B, 1995, **364**: 239; 1996, **370**: 167; 1996, **374**: 313; 1997, **407**: 193
- 11 Nakada T, Wolfenstein L. Eur. Phys. J. C, 1998, **3**: 457
- 12 Greenberg O W. Phys. Rev. Lett., 2002, **89**: 231602
- 13 Kostelecky V A. Phys. Rev. Lett., 1998, **80**: 1818; Phys. Rev. D, 2001, **64**: 076001
- 14 XING Z Z. Phys. Rev. D, 1994, **50**: 2957; Phys. Rev. D, 1997, **55**: 196; Phys. Lett. B, 1999, **450**: 202; Phys. Lett. B, 2000, **487**: 327; Ren P, XING Z Z. Phys. Rev. D, 2007, **76**: 116001