

Future Z factory as an intensity frontier for exploring dark sector

(Based on the preliminary result)

With Lian-Tao Wang, Xiao-Ping Wang, Wei Xue

Jia Liu

JGU Mainz

CEPC Theory Group Workshop, Beijing, 13 July, 2017



Cluster of Excellence

PRISMA

Precision Physics,
Fundamental Interactions
and Structure of Matter



JOHANNES GUTENBERG
UNIVERSITÄT MAINZ



Motivation: dark sector probes at FCC-ee

- How to related with new physics
 - LEP Shines Light on DM (Fox et al 1103.0240)
 - Electroweak Precision Observable (Linear collider report 1504.01726)
 - Z invisible width (Carena et al hep-ph/0308053)
 - Rare Z decay in SM (Huang et al, Durieux et al)
 - DM physics at CEPC (see Peng-fei's talk, also mono object searches at ee collider with XJ Bi et al)
 - Lepton flavor violation (see FuSheng's talk)
- Dark sector at low energy ee collider e.g. BES-III (JL, PF Yin, SH Zhu;, HB Li, T. Luo, V. Prasad, HB Li, XC Lou)



Motivation: dark sector probes at FCC-ee

- LEP at Z pole
 - $Z \rightarrow \gamma\gamma\gamma$ by L3
 - $Z \rightarrow \gamma + \text{inv}$ by L3, OPAL
 - $Z \rightarrow hZ^*$ by L3
 - $Z \rightarrow H_1H_2$ by L3
- LEP at higher energy
 - MSSM by LEP-II
 - Multiple photon + MET
 - Hadronic event with MET
 - Invisible Higgs
 - ...
- How to related with new physics
 - LEP Shines Light on DM (Fox et al 1103.0240)
 - Electroweak Precision Observable (Linear collider report 1504.01726)
 - Z invisible width (Carena et al hep-ph/0308053)
 - Rare Z decay in SM (Huang et al, Durieux et al)
 - DM physics at CEPC (see Peng-fei's talk, also mono object searches at ee collider with XJ Bi et al)
 - Lepton flavor violation (see FuSheng's talk)
- Dark sector at low energy ee collider e.g. BES-III (JL, PF Yin, SH Zhu;, HB Li, T. Luo, V. Prasad, HB Li, XC Lou)




Motivation: dark sector probes at FCC-ee

- Future Z factory (Giga/Tera)
 - Resonant prod $\sim 59000\text{pb}$, $10^9/10^{12}$ Z
 - Direct search on dark sector via exotic Z decay

- LEP at Z pole
 - $Z \rightarrow \gamma\gamma\gamma$ by L3
 - $Z \rightarrow \gamma + \text{inv}$ by L3, OPAL
 - $Z \rightarrow hZ^*$ by L3
 - $Z \rightarrow H_1 H_2$ by L3

- LEP at higher energy
 - MSSM by LEP-II
 - Multiple photon + MET
 - Hadronic event with MET
 - Invisible Higgs
 - ...





Exotic Z decay at FCC-ee

Dark Sectors 2016 Workshop: Community Report

1. Dark sector models

- Higgs portal + DM
- Vector portal + DM
- Axion-like particle
- Higher dimensional Operator
 - Comparing with existing collider and DM limits

2. Exotic Z decay topologies

- Motivating topologies
- Classifying by final state and resonances
 - Limits on BR

3. FCC-ee specialty: focus on invisible final state

$$\mathcal{L} \supset \begin{cases} -\frac{\epsilon}{2 \cos \theta_W} B_{\mu\nu} F'^{\mu\nu}, & \text{vector portal} \\ (\mu\phi + \lambda\phi^2) H^\dagger H, & \text{Higgs portal} \\ y_n L H N, & \text{neutrino portal} \\ \frac{a}{f_a} F_{\mu\nu} \tilde{F}^{\mu\nu}, & \text{axion portal.} \end{cases}$$

see Oliver's talk



Higgs portal + DM

- Lagrangian

$$\mathcal{L} = \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{\mu_S^2}{2} S^2 - \frac{\lambda_3}{6} S^3 - \frac{\lambda_4}{24} S^4 - \lambda_1 (H^\dagger H) S - \lambda_2 (H^\dagger H) S^2 + \bar{\chi} (i\not{\partial} - m_\chi^0) \chi - y_\chi S \bar{\chi} \chi + |D_\mu H|^2 - \mu_H^2 (H^\dagger H) - \lambda_H (H^\dagger H)^2.$$

- Scalar mixing

$$H = \frac{1}{\sqrt{2}} (v_H + h), \quad S = v_S + s \quad \begin{pmatrix} \tilde{h} \\ \tilde{s} \end{pmatrix} = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} h \\ s \end{pmatrix}$$

- Interactions and decays

- s interacts with SM from scalar mixing


$$\Gamma(\tilde{s} \rightarrow \bar{\chi} \chi) = \frac{y_\chi^2 \cos^2 \alpha}{8\pi} m_{\tilde{s}} \left(1 - \frac{4m_\chi^2}{m_{\tilde{s}}^2}\right)^{3/2}$$

$$\Gamma(\tilde{h} \rightarrow \tilde{s} \tilde{s}) = \frac{\sin^2 \alpha \cos^2 \alpha}{32\pi} \sqrt{1 - \frac{4m_{\tilde{s}}^2}{m_{\tilde{h}}^2}} \left(1 + 2 \frac{m_{\tilde{s}}^2}{m_{\tilde{h}}^2}\right)^2 \frac{m_{\tilde{h}}^3 (\cos \alpha v_H - \sin \alpha v_S)^2}{v_H^2 v_S^2}$$




Higgs portal + DM

- Limit from exotic Z decay $Z \rightarrow \tilde{s}Z^* \rightarrow (\bar{\chi}\chi) + l^+l^-$




- Various constraints







Higgs portal + DM

- Limit from exotic Z decay $Z \rightarrow \tilde{s}Z^* \rightarrow (\bar{\chi}\chi) + l^+l^-$



- Various constraints





Vector portal + DM

- Vector Lagrangian

$$\mathcal{L}_{\text{boson}} = -\frac{1}{4}B_{\mu\nu}B^{\mu\nu} - \frac{1}{4}W_{\mu\nu}^i W^{i\mu\nu} - \frac{1}{4}K_{\mu\nu}K^{\mu\nu} + \frac{\epsilon}{2c_W}B_{\mu\nu}K^{\mu\nu}$$

- Vector mixing

$$\begin{pmatrix} Z_{\mu, \text{SM}} \\ A_{\mu, \text{SM}} \\ K_{\mu} \end{pmatrix} = \begin{pmatrix} 1 + \frac{\epsilon^2 t_W^2}{2} \frac{m_{Z, \text{SM}}^2 (m_{Z, \text{SM}}^2 - 2m_K^2)}{(m_K^2 - m_{Z, \text{SM}}^2)^2} & 0 & \frac{m_K^2}{-m_K^2 + m_{Z, \text{SM}}^2} \epsilon t_W \\ \epsilon^2 t_W \frac{m_{Z, \text{SM}}^2}{m_K^2 - m_{Z, \text{SM}}^2} & 1 & \epsilon \\ \frac{m_{Z, \text{SM}}^2}{m_K^2 - m_{Z, \text{SM}}^2} \epsilon t_W & 0 & 1 + \frac{\epsilon^2}{2c_W^2} \frac{(m_{Z, \text{SM}}^4 c_W^2 - 2m_K^2 m_{Z, \text{SM}}^2 + m_K^4)}{(m_K^2 - m_{Z, \text{SM}}^2)^2} \end{pmatrix} \begin{pmatrix} \tilde{Z}_{\mu} \\ \tilde{A}_{\mu} \\ \tilde{K}_{\mu} \end{pmatrix}$$

- Interactions and decays

$$\mathcal{L}_{\text{int}} = \tilde{Z}_{\mu} \left(g J_Z^{\mu} - g_D \frac{m_{Z, \text{SM}}^2 t_W}{m_{Z, \text{SM}}^2 - m_K^2} \epsilon J_D^{\mu} + g \frac{m_{Z, \text{SM}}^2 (m_{Z, \text{SM}}^2 - 2m_K^2) t_W^2}{2(m_K^2 - m_{Z, \text{SM}}^2)^2} \epsilon^2 J_Z^{\mu} - e \frac{m_{Z, \text{SM}}^2 t_W}{m_{Z, \text{SM}}^2 - m_K^2} \epsilon^2 J_{\text{em}}^{\mu} \right) \\ + \tilde{K}_{\mu} \left(g_D J_D^{\mu} + g \frac{m_K^2 t_W}{m_{Z, \text{SM}}^2 - m_K^2} \epsilon J_Z^{\mu} + e \epsilon J_{\text{em}}^{\mu} + g_D \frac{(m_{Z, \text{SM}}^4 c_W^2 - 2m_K^2 m_{Z, \text{SM}}^2 + m_K^4) c_W^{-2}}{2(m_{Z, \text{SM}}^2 - m_K^2)^2} \epsilon^2 J_D^{\mu} \right)$$



Vector portal + DM



- Complex Scalar DM Lagrangian

$$\mathcal{L}_S = (\partial_\mu S + ig_D K_\mu S)^* (\partial^\mu S + ig_D K^\mu S) - m_S^2 S^* S$$

- Connection to Z

$$\mathcal{L}_S \supset g_D^2 S^* S \left(\tilde{K}_\mu + t_W \epsilon \frac{m_{Z, SM}^2}{(m_K^2 - m_{Z, SM}^2)} \tilde{Z}_\mu \right)^2$$

- Various constraints





Vector portal + DM



- Complex Scalar DM Lagrangian

$$\mathcal{L}_S = (\partial_\mu S + ig_D K_\mu S)^* (\partial^\mu S + ig_D K^\mu S) - m_S^2 S^* S + \lambda_1 S^* S \Phi^* \Phi + \lambda_2 S^* S H^\dagger H + (\mu_1 \Phi^* S S + h.c.) .$$

- Connection to Z

$$\mathcal{L}_S \supset g_D^2 S^* S \left(\tilde{K}_\mu + t_W \epsilon \frac{m_{Z, SM}^2}{(m_K^2 - m_{Z, SM}^2)} \tilde{Z}_\mu \right)^2$$

- Various constraints





Axion-like particle


- Lagrangian

$$\mathcal{L}_{\text{ALP}} = \frac{1}{4\Lambda_{aBB}} a B_{\mu\nu} \tilde{B}^{\mu\nu}$$

- Interactions and decays

$$\Gamma(Z \rightarrow \gamma a) = \frac{1}{64\pi} \frac{1}{\Lambda_{aBB}^2} \cos^2 \theta_w \sin^2 \theta_w m_a^3$$

$$\Gamma(a \rightarrow \gamma\gamma) = \frac{1}{64\pi} \frac{1}{\Lambda_{aBB}^2} \cos^4 \theta_w m_a^3$$





Axion-like particle

- Lagrangian



$$\mathcal{L}_{ALP} = \frac{1}{4\Lambda_{aBB}} a B_{\mu\nu} \tilde{B}^{\mu\nu}$$

- Interactions and decays

$$\Gamma(Z \rightarrow \gamma a) = \frac{1}{64\pi} \frac{1}{\Lambda_{aBB}^2} \cos^2 \theta_w \sin^2 \theta_w m_a^3$$

$$\Gamma(a \rightarrow \gamma\gamma) = \frac{1}{64\pi} \frac{1}{\Lambda_{aBB}^2} \cos^4 \theta_w m_a^3$$

- Various constraints








Higher dimensional operator

- UV model

$$\mathcal{L} = \bar{\chi}(i\not{D} - m_\chi)\chi - \frac{1}{2}\delta m\bar{\chi}^c\chi + \bar{\psi}(i\not{D} - M_\psi)\psi + (D^\mu\phi)^\dagger(D_\mu\phi) - M_\phi^2\phi^\dagger\phi + (\lambda\bar{\psi}\chi\phi + h.c.),$$



- Interactions: **Magnetic inelastic DM and Rayleigh DM** (Weiner, Yavin)

$$O_{\text{MIDM}} = \frac{1}{\Lambda_{\text{MIDM}}}\bar{\chi}_2\sigma^{\mu\nu}\chi_1 B_{\mu\nu} + h.c., \quad O_{\text{RayDM}} = \frac{1}{\Lambda_{\text{RayDM}}^3}\bar{\chi}_1\chi_1 B^{\mu\nu} B_{\mu\nu}.$$



- Scale from mass in the loop





$$\frac{1}{\Lambda_{\text{MIDM}}} \approx \frac{\lambda^2 g_Y}{32\pi^2 M_\psi}, \quad \frac{1}{\Lambda_{\text{RayDM}}^3} \approx \frac{\lambda^2 g_Y^2}{24\pi^2 M_\psi^3},$$





Higher dimensional operator

- Magnetic inelastic DM and Rayleigh DM





Exotic Z decay topologies

- Exotic Z decay final states

$$Z \rightarrow n_{\cancel{E}} \cancel{E} + n_{\gamma} \gamma + n_{\ell^+ \ell^-} \ell^+ \ell^- + n_{\bar{q}q} \bar{q}q$$

exotic decay	topologies	n_{res}	models
$Z \rightarrow \cancel{E} + \gamma$	$Z \rightarrow \chi_1 \chi_2, \chi_2 \rightarrow \chi_1 \gamma$	0	2A: $\frac{1}{\Lambda} \bar{\chi}_2 \sigma^{\mu\nu} \chi_1 B_{\mu\nu}$ (MIDM)
	$Z \rightarrow \chi \bar{\chi} \gamma$	0	2B: $\frac{1}{\Lambda^3} \bar{\chi} \chi B_{\mu\nu} B^{\mu\nu}$ (RayDM)
	$Z \rightarrow a \gamma \rightarrow (\cancel{E}) \gamma$	1	2C: $\frac{1}{\Lambda_{2C}} a B_{\mu\nu} \tilde{B}^{\mu\nu}$ (long-lived ALP)
	$Z \rightarrow A' \gamma \rightarrow (\bar{\chi} \chi) \gamma$	1	2D: $\epsilon^{\mu\nu\rho\sigma} A'_\mu B_\nu \partial_\rho B_\sigma$ (WZ terms)
$Z \rightarrow \cancel{E} + \gamma\gamma$	$Z \rightarrow \phi_d A', \phi_d \rightarrow (\gamma\gamma), A' \rightarrow (\bar{\chi} \chi)$	2	3A: Vector portal
	$Z \rightarrow \phi_H \phi_A, \phi_H \rightarrow (\gamma\gamma), \phi_A \rightarrow (\bar{\chi} \chi)$	2	3B: 2HDM extension
	$Z \rightarrow \chi_2 \chi_1, \chi_2 \rightarrow \chi_1 \phi, \phi \rightarrow (\gamma\gamma)$	1	3C: Inelastic DM
	$Z \rightarrow \chi_2 \chi_2, \chi_2 \rightarrow \gamma \chi_1$	0	3D: MIDM
$Z \rightarrow \cancel{E} + \ell^+ \ell^-$	$Z \rightarrow \phi_d A', A' \rightarrow (\ell^+ \ell^-), \phi_d \rightarrow (\bar{\chi} \chi)$	2	4A: Vector portal
	$Z \rightarrow A' S S \rightarrow (\ell\ell) S S$	1	4B: Vector portal
	$Z \rightarrow \phi(Z^*/\gamma^*) \rightarrow \phi \ell^+ \ell^-$	1	4C: Long-lived ALP, Higgs portal
	$Z \rightarrow \chi_2 \chi_1 \rightarrow \chi_1 A' \chi_1 \rightarrow (\ell^+ \ell^-) \cancel{E}$	1	4D: Vector portal and Inelastic DM
	$Z \rightarrow \chi_2 \chi_1, \chi_2 \rightarrow \chi_1 \ell^+ \ell^-$	0	4E: MIDM, SUSY
	$Z \rightarrow \bar{\chi} \chi \ell^+ \ell^-$	0	4F: RayDM, slepton, heavy lepton mixing
$Z \rightarrow \cancel{E} + J J$	similar to $\cancel{E} + \ell^+ \ell^-$		15



Exotic Z decay topologies

- Exotic Z decay final states

$$Z \rightarrow n_{\cancel{E}} \cancel{E} + n_{\gamma} \gamma + n_{\ell^+ \ell^-} \ell^+ \ell^- + n_{\bar{q}q} \bar{q}q$$

exotic decay	topologies	n_{res}	models
$Z \rightarrow \cancel{E} + \gamma$	$Z \rightarrow \chi_1 \chi_2, \chi_2 \rightarrow \chi_1 \gamma$	0	2A: $\frac{1}{\Lambda} \bar{\chi}_2 \sigma^{\mu\nu} \chi_1 B_{\mu\nu}$ (MIDM)
	$Z \rightarrow \chi \bar{\chi} \gamma$	0	2B: $\frac{1}{\Lambda^3} \bar{\chi} \chi B_{\mu\nu} B^{\mu\nu}$ (RayDM)
	$Z \rightarrow a \gamma \rightarrow (\cancel{E}) \gamma$	1	2C: $\frac{1}{\Lambda_{2C}} a B_{\mu\nu} \tilde{B}^{\mu\nu}$ (long-lived ALP)
	$Z \rightarrow A' \gamma \rightarrow (\bar{\chi} \chi) \gamma$	1	2D: $\frac{1}{\Lambda} \bar{\chi} \chi B_{\mu\nu} B^{\mu\nu}$ (MIDM)
$Z \rightarrow \cancel{E} + \gamma\gamma$	$Z \rightarrow \phi_d A', \phi_d \rightarrow (\gamma\gamma), A' \rightarrow (\bar{\chi} \chi)$	2	3A: Vector portal $Z \rightarrow \phi_d A' \rightarrow (JJ)(JJ)$
	$Z \rightarrow \phi_H \phi_A, \phi_H \rightarrow (\gamma\gamma), \phi_A \rightarrow (\bar{\chi} \chi)$	2	3B: 2HDM extension $Z \rightarrow \phi_H \phi_A \rightarrow (JJ)(JJ)$
	$Z \rightarrow \chi_2 \chi_1, \chi_2 \rightarrow \chi_1 \phi, \phi \rightarrow (\gamma\gamma)$	1	3C: Inelastic DM $Z \rightarrow \chi_2 \chi_1 \rightarrow \chi_1 \phi \rightarrow (\gamma\gamma) \cancel{E}$
	$Z \rightarrow \chi_2 \chi_2, \chi_2 \rightarrow \gamma \chi_1$	0	3D: MIDM
	$Z \rightarrow \phi \gamma \rightarrow (\gamma\gamma) \gamma$	1	3E: ALP $Z \rightarrow \phi \gamma \rightarrow (\gamma\gamma) \gamma$
$Z \rightarrow \cancel{E} + \ell^+ \ell^-$	$Z \rightarrow \phi_d A', A' \rightarrow (\ell^+ \ell^-), \phi_d \rightarrow (\bar{\chi} \chi)$	2	4A: Vector portal
	$Z \rightarrow A' S S \rightarrow (\ell\ell) S S$	1	4B: Vector portal
	$Z \rightarrow \phi (Z^*/\gamma^*) \rightarrow \phi \ell^+ \ell^-$	1	4C: Long-lived ALP, Higgs portal
	$Z \rightarrow \chi_2 \chi_1 \rightarrow \chi_1 A' \chi_1 \rightarrow (\ell^+ \ell^-) \cancel{E}$	1	4D: Vector portal and Inelastic DM
	$Z \rightarrow \chi_2 \chi_1, \chi_2 \rightarrow \chi_1 \ell^+ \ell^-$	0	4E: MIDM, SUSY
	$Z \rightarrow \bar{\chi} \chi \ell^+ \ell^-$	0	4F: RayDM, slepton, heavy lepton mixing
$Z \rightarrow \cancel{E} + JJ$	similar to $\cancel{E} + \ell^+ \ell^-$		

$Z \rightarrow \phi_d A' \rightarrow (JJ)(JJ)$	2	6A: Vector portal
$Z \rightarrow \phi_H \phi_A \rightarrow (JJ)(JJ)$	2	6B: 2HDM
$Z \rightarrow G_0 G_0, G_0 \rightarrow (bb)$	2	6C: Hidden Valley
$Z \rightarrow \phi \gamma \rightarrow (\gamma\gamma) \gamma$	1	7A: ALP, Higgs portal



CEPC collider setup

- Detector performance

$$\text{Photon energy resolution: } \frac{\delta E_\gamma}{E_\gamma} = \frac{0.16}{\sqrt{E_\gamma/\text{GeV}}} \oplus 0.01,$$

$$\text{Lepton momentum resolution: } \Delta \frac{\text{GeV}}{p_T^\ell} = 2 \times 10^{-5} \oplus \frac{10^{-3}\text{GeV}}{p_T^\ell \sin \theta},$$

$$\text{Jet energy resolution: } \frac{\delta E_j}{E_j} = \frac{0.3}{\sqrt{E_j/\text{GeV}}} \oplus 0.02.$$

- Preliminary cuts

$$|\eta| < 2.3, E_\gamma > 10\text{GeV}, E_\ell > 5\text{GeV}, E_j > 10\text{GeV}, E_{(\text{MET})} > 10\text{GeV},$$
$$y_{ij} \equiv \frac{2\text{Min}(E_i^2, E_j^2) (1 - \cos \theta_{ij})}{E_{vis}^2} \geq 0.001$$

- SM background including one photon from ISR



Limits on exotic Z decay BR

- Mono-photon from MIDM

$$Z \rightarrow \chi_2 \chi_1 \rightarrow \chi_1 \chi_1 + \gamma$$

- Specific cuts


$$2A : E_{\gamma}^{\min,2A} < E_{\gamma} < E_{\gamma}^{\max,2A}, \quad m_{\text{inv}} \geq 2m_{\chi_1}$$

$$E_{\gamma}^{\max,2A} = \frac{m_2^2 - m_1^2 s + m_2^2 - m_1^2 + \sqrt{s^2 + (m_2^2 - m_1^2)^2 - 2s(m_2^2 + m_1^2)}}{4m_2^2 \sqrt{s}}$$

$$E_{\gamma}^{\min,2A} = \frac{m_2^2 - m_1^2 s + m_2^2 - m_1^2 - \sqrt{s^2 + (m_2^2 - m_1^2)^2 - 2s(m_2^2 + m_1^2)}}{4m_2^2 \sqrt{s}}$$

- Luminosity scaling

$$\frac{S}{\sqrt{B}} \sim \sqrt{L} = 10^{1.5}$$







Limits on exotic Z decay BR

- Mono-photon

$Z \rightarrow \chi\bar{\chi}\gamma$	0	2B: $\frac{1}{\Lambda^3} \bar{\chi}\chi B_{\mu\nu} B^{\mu\nu}$ (RayDM)
$Z \rightarrow a\gamma \rightarrow (\cancel{E})\gamma$	1	2C: $\frac{1}{\Lambda_{2C}} a B_{\mu\nu} \tilde{B}^{\mu\nu}$ (long-lived ALP)
$Z \rightarrow A'\gamma \rightarrow (\bar{\chi}\chi)\gamma$	1	2D: $\epsilon^{\mu\nu\rho\sigma} A'_\mu B_\nu \partial_\rho B_\sigma$ (WZ terms)







Limits on exotic Z decay BR

- Di-lepton + MET

$Z \rightarrow \phi_d A', A' \rightarrow (\ell^+ \ell^-), \phi_d \rightarrow (\bar{\chi} \chi)$	2	4A: Vector portal
$Z \rightarrow A' S S \rightarrow (\ell \ell) S S$	1	4B: Vector portal
$Z \rightarrow \phi(Z^*/\gamma^*) \rightarrow \phi \ell^+ \ell^-$	1	4C: Long-lived ALP, Higgs portal
$Z \rightarrow \chi_2 \chi_1 \rightarrow \chi_1 A' \chi_1 \rightarrow (\ell^+ \ell^-) \cancel{E}$	1	4D: Vector portal and Inelastic DM
$Z \rightarrow \chi_2 \chi_1, \chi_2 \rightarrow \chi_1 \ell^+ \ell^-$	0	4E: MIDM, SUSY
$Z \rightarrow \bar{\chi} \chi \ell^+ \ell^-$	0	4F: RayDM, slepton, heavy lepton mixing





Summary

- Dark sector models
 - Higgs portal, Vector portal,
 - Axion-like Particle, Higher dimension Operator
 - Can provide **leading and complementary** constraints comparing to current collider limits and DM limits
- Exotic Z decay topologies
 - Giga Z limit BR $10^{-6} \sim 10^{-8.5}$
 - Tera Z limit BR $10^{-7.5} \sim 10^{-11}$
 - Sensitivities on BR
$$l^+l^- + \text{MET} > \gamma\gamma + \text{MET} > \text{JJ} + \text{MET} > \gamma + \text{MET}$$



Thank you!



Backup slides






Limits on exotic Z decay BR

- Di-photon + MET

3C : $|m_{\gamma\gamma} - m_{\phi_d}| < 2.5\text{GeV}, \quad m_{\text{inv}} > 2m_{\chi_1}$

3D : $E_{\gamma}^{\text{max},3D} > E_{\gamma} > E_{\gamma}^{\text{min},3D}, \quad m_{\text{inv}} > 2m_{\chi_1}$

$Z \rightarrow \phi_d A', \phi_d \rightarrow (\gamma\gamma), A' \rightarrow (\bar{\chi}\chi)$	2	3A: Vector portal
$Z \rightarrow \phi_H \phi_A, \phi_H \rightarrow (\gamma\gamma), \phi_A \rightarrow (\bar{\chi}\chi)$	2	3B: 2HDM extension
$Z \rightarrow \chi_2 \chi_1, \chi_2 \rightarrow \chi_1 \phi, \phi \rightarrow (\gamma\gamma)$	1	3C: Inelastic DM
$Z \rightarrow \chi_2 \chi_2, \chi_2 \rightarrow \gamma \chi_1$	0	3D: MIDM







Limits on exotic Z decay BR

- Di-lepton + MET




$Z \rightarrow \phi_d A', A' \rightarrow (\ell^+ \ell^-), \phi_d \rightarrow (\bar{\chi} \chi)$	2	4A: Vector portal
$Z \rightarrow A' S S \rightarrow (\ell \ell) S S$	1	4B: Vector portal
$Z \rightarrow \phi (Z^* / \gamma^*) \rightarrow \phi \ell^+ \ell^-$	1	4C: Long-lived ALP, Higgs portal
$Z \rightarrow \chi_2 \chi_1 \rightarrow \chi_1 A' \chi_1 \rightarrow (\ell^+ \ell^-) \cancel{E}$	1	4D: Vector portal and Inelastic DM
$Z \rightarrow \chi_2 \chi_1, \chi_2 \rightarrow \chi_1 \ell^+ \ell^-$	0	4E: MIDM, SUSY
$Z \rightarrow \bar{\chi} \chi \ell^+ \ell^-$	0	4F: RayDM, slepton, heavy lepton mixing





Limits on exotic Z decay BR




- Di-jet + MET similar to Di-lepton + MET





Limits on exotic Z decay BR

- Z decay to (JJ)(JJ)





Limits on exotic Z decay BR

- Z decay to (AA)A

