



Hyperons at BESIII


平荣刚 (Ping Ronggang)

高能物理研究所(IHEP)

pingrg@ihep.ac.cn

中国科学院粒子物理前沿卓越创新中心第七次全体会议, 2019-12-06

BESIII $J/\psi, \psi'$ data sets



The analysis based on
2009+2012 J/ψ data:

~1.31 billion decays


J/ψ : Total 10.047 billion J/ψ decays

ψ' : 448 million decays

Continuum: 12/fb ($\sqrt{s} > 3.8$ GeV)


Hyperon pair production at BESIII

- $e^+e^- \rightarrow \gamma^* \rightarrow \Lambda\bar{\Lambda}, \Sigma\bar{\Sigma}, \Xi\bar{\Xi}, \Omega\bar{\Omega}, \Lambda_c^+\bar{\Lambda}_c^-,$ @ $\sqrt{s} = 2.0 \sim 4.6$ GeV, or update for $\Lambda_c^+\bar{\Sigma}_c^-, \Sigma_c\bar{\Sigma}_c$



- Threshold enhancement
- Form factor
- Excited hyperon
-

- $J/\psi, \psi' \rightarrow \Lambda\bar{\Lambda}, \Sigma\bar{\Sigma}, \Xi\bar{\Xi}, \Omega\bar{\Omega}$




10 billion J/ψ :

16.1×10^6	$\Lambda\bar{\Lambda}$
12.0×10^6	$\Xi^0\bar{\Xi}^0$
12.9×10^6	$\Sigma^0\bar{\Sigma}^0$
8.6×10^6	$\Xi^-\bar{\Xi}^+$

- **Hyperon polarization**
- CP test
- Semileptonic decay
- Radiative decay
- Rare decay

C,P-transformation in $\Lambda \rightarrow p\pi^-$



$$\alpha = \frac{|B_+|^2 - |B_-|^2}{|B_+|^2 + |B_-|^2},$$

$$\bar{\alpha} = \frac{|\bar{B}_+|^2 - |\bar{B}_-|^2}{|\bar{B}_+|^2 + |\bar{B}_-|^2}$$

CP invariance :

$$\bar{B}_{-\lambda_p} = \eta_\Lambda \eta_p \eta_\pi (-1)^{s_\Lambda - s_p - s_\pi} B_{\lambda_p}$$

$$= -B_{\lambda_p}$$

$$\alpha = -\bar{\alpha}$$

CP odd-variable:

$$\Delta = \frac{\Gamma - \bar{\Gamma}}{\Gamma + \bar{\Gamma}}, A = \frac{\alpha + \bar{\alpha}}{\alpha - \bar{\alpha}}, B = \frac{\beta + \bar{\beta}}{\beta - \bar{\beta}}, B' = \frac{\beta + \bar{\beta}}{\alpha - \bar{\alpha}}$$

Ancient history

Phys. Rev. Lett. 24, 843 (1970)

ASYMMETRY PARAMETER FOR $\Lambda^0 \rightarrow \pi\pi^0 \uparrow^*$

S. Olsen, L. Pondrom, R. Handler, and P. Limon†
University of Wisconsin, Madison, Wisconsin 53706

and

J. A. Smith and O. E. Overseth
University of Michigan, Ann Arbor, Michigan 48104
(Received 18 February 1970)

The asymmetry parameter α_0 for $\Lambda^0 \rightarrow \pi\pi^0$ has been measured relative to α_- by comparing the neutron distribution with the proton distribution from the decay $\Lambda^0 \rightarrow p\pi^-$ for polarized Λ^0 hyperons. A sample of 4760 neutron decay events and 8500 proton decay events gave $\alpha_0/\alpha_- = 1.000 \pm 0.068$ in good agreement with the $|\Delta I| = \frac{1}{2}$ rule.

Phys. Rev. Lett. 24, 165 (1970)

PROTON POLARIZATION IN $\Sigma^+ \rightarrow p\pi^0 \uparrow^*$

F. Harris and O. E. Overseth
University of Michigan, Ann Arbor, Michigan 48104

and

L. Pondrom and E. Dettmann
University of Wisconsin, Madison, Wisconsin 53706
(Received 10 November 1969)

The polarization of protons from the decay of polarized Σ^+ hyperons has been measured by scattering the protons in a carbon-plate spark chamber. A sample of 1335 useful scatters gave $\alpha_0 = -0.98 \pm 0.05$ and $\varphi_0 = 22^\circ \pm 90^\circ$, where $\tan\varphi_0 = \beta/\gamma_0$. Using the data on $\Sigma^+ \rightarrow p\pi^0$ and $\Sigma^- \rightarrow n\pi^-$ and fitting to the $|\Delta I| = \frac{1}{2}$ rule gave $\chi^2 = 0.3$ for 2 degrees of freedom.

Phys. Rev. 184, 1663 (1969)

Final-State Interactions in Nonleptonic Hyperon Decay

O. E. OVERSETH*

The University of Michigan, Ann Arbor, Michigan 48104


AND

S. PAKVASA†

University of Hawaii, Honolulu, Hawaii 96822

(Received 1 April 1969)

We discuss the consequences of including the final-state interactions in the analysis of Λ^0 , Σ , and Ξ nonleptonic decays. Emphasis is on the role that the final-state interactions play in tests for T invariance, in tests of the $\Delta I = \frac{1}{2}$ rule (including the resolution of sign ambiguities), and in the determination of the decay amplitudes for these processes.




Observation of Λ polarization and entanglement in


 **Highlights!**

$$J/\psi \rightarrow \Lambda \bar{\Lambda}$$


Published in Nature Physics, 15, 631 (2019)



Helicity system



Spin entanglement



Event display in BESIII detector

➤ Joint angular distribution

$$\begin{aligned} \mathcal{W}(\xi; \alpha_\psi, \Delta\Phi, \alpha_-, \alpha_+) &= 1 + \alpha_\psi \cos^2 \theta_\Lambda + \alpha_- \alpha_+ \sin^2 \theta_\Lambda (n_{1,x} n_{2,x} - \alpha_\psi n_{1,y} n_{2,y}) \\ &+ (\cos^2 \theta_\Lambda + \alpha_\psi) n_{1,z} n_{2,z} \quad \text{Entanglement term} \\ &+ \alpha_- \alpha_+ \sqrt{1 - \alpha_\psi^2} \cos(\Delta\Phi) \sin \theta_\Lambda \cos \theta_\Lambda (n_{1,x} n_{2,z} + n_{1,z} n_{2,x}) \\ &+ \sqrt{1 - \alpha_\psi^2} \sin(\Delta\Phi) \sin \theta_\Lambda \cos \theta_\Lambda (\alpha_- n_{1,y} + \alpha_+ n_{2,y}) \end{aligned}$$

Transverse polarization

➤ Transverse polarization

$$P_y(\cos \theta_\Lambda) = \frac{\sqrt{1 - \alpha_\psi^2} \sin(\Delta\Phi) \cos \theta_\Lambda \sin \theta_\Lambda}{1 + \alpha_\psi \cos^2 \theta_\Lambda}$$

Observation of Λ polarization and entanglement in





$$J/\psi \rightarrow \Lambda \bar{\Lambda}$$

Published in Nature Physics, 15, 631 (2019)

J/ψ data sample: 1.31×10^9

➤ Observation of $\Lambda / \bar{\Lambda}$ transverse polarization



• spin observable

$$\mu(\cos \theta_\Lambda) = \frac{m}{N} \sum_{i=1}^{N_k} (n_{1,y}^{(i)} - n_{2,y}^{(i)})$$

- Maximum $\Lambda / \bar{\Lambda}$ polarization: $\sim 25\%$
- Net polarization vanishing

Observation of Λ polarization and entanglement in



$$J/\psi \rightarrow \Lambda \bar{\Lambda}$$

Published in Nature Physics, 15, 631 (2019)

J/ψ data sample: 1.31×10^9

First observation of the $\Lambda/\bar{\Lambda}$ transverse polarization

Table 1 | Summary of the results

Parameters	This work	Previous results
α_ψ	$0.461 \pm 0.006 \pm 0.007$	0.469 ± 0.027 (ref. ¹⁴)
$\Delta\Phi$	$42.4 \pm 0.6 \pm 0.5^\circ$	-
α_-	$0.750 \pm 0.009 \pm 0.004$	0.642 ± 0.013 (ref. ⁶)
α_+	$-0.758 \pm 0.010 \pm 0.007$	-0.71 ± 0.08 (ref. ⁶)
$\bar{\alpha}_0$	$-0.692 \pm 0.016 \pm 0.006$	-
A_{CP}	$-0.006 \pm 0.012 \pm 0.007$	0.006 ± 0.021 (ref. ⁶)
$\bar{\alpha}_0/\alpha_+$	$0.913 \pm 0.028 \pm 0.012$	-

Higher than PDG value 17%, deviation with sig. $> 5\sigma$

CP odd observable:

$$A_{CP} = \frac{\alpha_\Lambda - \alpha_{\bar{\Lambda}}}{\alpha_\Lambda + \alpha_{\bar{\Lambda}}}$$

Test selection rule $\Delta I = \frac{1}{2}$

- Verification $\alpha_\Lambda/\alpha_{\bar{\Lambda}} : J/\psi \rightarrow \Xi^- \bar{\Xi}^+ \rightarrow (\Lambda \pi^-)(\bar{\Lambda} \pi^+)$
- Transverse polarization allows to improve CP test precision over previous measurements.
- Standard model precision $A_{CP} \sim 10^{-4}$.

Observation of Λ polarization and entanglement in



Highlights!

$$J/\psi \rightarrow \Lambda \bar{\Lambda}$$

α_- FOR $\Lambda \rightarrow p\pi^-$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.750 \pm 0.009 \pm 0.004$	420k	ABLIKIM 2018AG	BES3	J/ψ to $\Lambda \bar{\Lambda}$
••• We do not use the following data for averages, fits, limits, etc. •••				
0.584 ± 0.046	8500	ASTBURY 1975	SPEC	
0.649 ± 0.023	10325	CLELAND 1972	OSPK	
0.67 ± 0.06	3520	DAUBER 1969	HBC	From E decay
0.645 ± 0.017	10130	OVERSETH 1967	OSPK	Λ from $\pi^- p$
0.62 ± 0.07	1156	CRONIN 1963	CNTR	Λ from $\pi^- p$

α_+ FOR $\bar{\Lambda} \rightarrow \bar{p}\pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$-0.758 \pm 0.010 \pm 0.007$	420k	ABLIKIM 2018AG	BES3	J/ψ to $\Lambda \bar{\Lambda}$
••• We do not use the following data for averages, fits, limits, etc. •••				
$-0.755 \pm 0.083 \pm 0.063$	$\approx 8.7k$	ABLIKIM 2010	BES	$J/\psi \rightarrow \Lambda \bar{\Lambda}$
-0.63 ± 0.13	770	TIXIER 1988	DM2	$J/\psi \rightarrow \Lambda \bar{\Lambda}$

Where does the TP come from?

□ From the e^+ / e^- beam ?

✗ No, BEPC beams unpolarized

□ From the e^+ / e^- natural polarization when circulating in the BEPCII storage ring ?

✗ Sokolov-Ternov effects: 4.3 hrs @ ψ' peak, but beam lifetime ~ 2.0 hrs

□ From the J/ψ spin transfer ?


✓ Yes, it does from the J/ψ tensor polarization

J/ψ polarization: $\mathcal{P}_z = 0$, $T_{zz} = \frac{1}{\sqrt{6}}$


Λ transverse polarization:

$$\mathcal{P}_y = \sqrt{6} \frac{T_{zz} \sin \theta \cos \theta \sin \Delta \sqrt{1 - \alpha_\psi^2}}{1 + \alpha_\psi \left[\frac{1}{3} + \frac{1}{\sqrt{6}} T_{zz} (1 + 3 \cos 2\theta) \right]}$$

\mathcal{P}_y manifest if $\sin \Delta \neq 0$



$\Lambda / \bar{\Lambda}$ polarization in continuum production



data set: 2.396 GeV, $L=66.9 \text{ pb}^{-1}$



$$e^+e^- \rightarrow \gamma^* \rightarrow \Lambda\bar{\Lambda}$$

555 candidate events

$\Delta\Phi = 37^\circ \pm 12^\circ (\text{stat}) \pm 6^\circ (\text{syst})$





Maximum polarization degree is $\sim 30\%$

$$P_y = \frac{\sqrt{1-\eta^2} \sin\theta \cos\theta}{1+\eta\cos^2\theta} \sin(\Delta\Phi).$$



Polarization puzzle in charmonium decays

Angular distribution: $\frac{dN}{d\cos\theta} \propto (1 + \alpha \cos^2 \theta)$



Branching ratios

$$\frac{\mathcal{B}(\psi(3686) \rightarrow \Lambda \bar{\Lambda})}{\mathcal{B}(J/\psi \rightarrow \Lambda \bar{\Lambda})} = (20.43 \pm 0.11 \pm 0.58)\%$$

$$\frac{\mathcal{B}(\psi(3686) \rightarrow \Sigma^0 \bar{\Sigma}^0)}{\mathcal{B}(J/\psi \rightarrow \Sigma^0 \bar{\Sigma}^0)} = (20.96 \pm 0.27 \pm 0.92)\%$$

Angular distribution

$$\alpha = \frac{|H_{\uparrow\downarrow}|^2 - 2|H_{\uparrow\uparrow}|^2}{|H_{\uparrow\downarrow}|^2 + 2|H_{\uparrow\uparrow}|^2}$$


Helicity selection rule

$$|H_{\uparrow\uparrow}|/|H_{\uparrow\downarrow}| \sim 0$$

Measurements:

$$|H_{\uparrow\uparrow}|/|H_{\uparrow\downarrow}| = \begin{aligned} &\sim 1.3 (J/\psi \rightarrow \Sigma^0 \bar{\Sigma}^0) \\ &\sim 0.08 (\psi' \rightarrow \Sigma^0 \bar{\Sigma}^0) \end{aligned}$$

Λ_c spin and decay asymmetry parameter



Λ_c^+

udc


$I(J^P) = 0(\frac{1}{2}^+)$ Status: ****

The parity of the Λ_c^+ is defined to be positive (as are the parities of the proton, neutron, and Λ). The quark content is udc . Results of an analysis of $pK^-\pi^+$ decays (JEZABEK 92) are consistent with $J = 1/2$. Nobody doubts that the spin is indeed $1/2$.


We have omitted some results that have been superseded by later experiments. The omitted results may be found in earlier editions.

Predictions on Λ_c asymmetry parameters

- W -exchange complicate $\Lambda_c \rightarrow B M$





$\Lambda_c^+ \rightarrow BP$ decay asymmetry



Λ_c decay asymmetry parameter

- $L = 567\text{pb}^{-1}$ @ $\sqrt{s} = 4.6$ GeV with single tag



Phys. Rev. D 100, 072004 (2019)

- Significance for Λ_c transverse polarization: 2.1σ


CPV with $J/\psi \rightarrow \Xi^- (\Lambda\pi^-) \bar{\Xi}^+ (\Lambda\pi^+) + \text{c.c.}$

缺点:

complicated topology: 9-dimensions

$\theta_{\Xi}, \theta_{\Lambda}, \phi_{\Lambda}, \theta_{\bar{\Lambda}}, \phi_{\bar{\Lambda}}, \theta_p, \phi_p, \theta_{\bar{p}}, \phi_{\bar{p}}$

72 terms, 8 parameters to determine



Low rate compared to $\Lambda\bar{\Lambda}$

1.3B J/ψ : 420K $\Lambda(p\pi^-)\bar{\Lambda}(\bar{p}\pi^+)$ evts

61K $\Xi(\Lambda\pi^-)\bar{\Xi}(\Lambda\pi^+)$

优点:

- $\Lambda\bar{\Lambda}$ polarizations are measurable via their parity-violating $p\pi^-$ ($\bar{p}\pi^+$) decays;
- β_- and β_0 parameters can be determined.
- Preliminary results indicate that the Ξ s are even more polarized than the Λ s.

Ongoing polarization analyses at BESIII

- $J/\psi, \psi' \rightarrow \Sigma^- \bar{\Sigma}^+$
- $J/\psi \rightarrow \Xi^- \bar{\Xi}^+$ (Ξ, Λ asymmetry par., CP test)
- $\psi' \rightarrow \Omega^- \bar{\Omega}^+$ (polarization analysis)
- $\eta_c \rightarrow \Lambda \bar{\Lambda}$ (QM nonlocality & reality)
- $e^+ e^- \rightarrow \Lambda_c^+ \bar{\Lambda}_c^-$
 - $\Lambda_c^+ \rightarrow \Lambda \pi^+, \Sigma^+ \pi^0, \Sigma^0 \pi^+, p K_S$, to be improved
 - Λ_c spin
- future charmed baryon program
 - Update E_{cms} up to 4.9 GeV
 - Approved $\sim 10 \text{ fb}^{-1}$ data taking around $\Lambda_c \bar{\Lambda}_c$ threshold

Summary

- $\Lambda/\bar{\Lambda}$ transverse polarization significantly observed at BESIII in J/ψ or continuum processes
- BESIII 10 billion J/ψ data provides us chances to access hyperon physics.
- Extension study to charmed hyperon are ongoing.
- Polarized beam in the future super-tau charm facility (STCF) help to improve the precision.

敬请指教！

backup

Previous Measurements

2018 PDG list

α_- FOR $\Lambda \rightarrow p\pi^-$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.642 ± 0.013	OUR AVERAGE			
0.584 ± 0.046	8500	ASTBURY	1975	SPEC
0.649 ± 0.023	10325	CLELAND	1972	OSPK
0.67 ± 0.06	3520	DAUBER	1969	HBC From Ξ decay
0.645 ± 0.017	10130	OVERSETH	1967	OSPK Λ from $\pi^- p$
0.62 ± 0.07	1156	CRONIN	1963	CNTR Λ from $\pi^- p$



α_+ FOR $\bar{\Lambda} \rightarrow \bar{p}\pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.71 ± 0.08	OUR AVERAGE			
$-0.755 \pm 0.083 \pm 0.063$	$\approx 8.7k$	ABLIKIM	2010	BES $J/\psi \rightarrow \Lambda\bar{\Lambda}$
-0.63 ± 0.13	770	TIXIER	1988	DM2 $J/\psi \rightarrow \Lambda\bar{\Lambda}$

Most earlier measurement on α

- CNTR exp., $\pi^- + p \rightarrow \Lambda + K^0$
- Unpolarized $\Lambda \rightarrow p\pi^-$
- Proton polarization measured with carbon-plate spark chamber

Phys.Rev. 129 (1963) 1795-1807



$$W(\psi) = 1 + \alpha S \sin \epsilon \cos \psi$$




FIG. 1. Schematic diagram showing arrangement of apparatus. An example of an event has been sketched in.

$$\alpha = - \frac{2}{\pi} \frac{1}{\langle S \rangle \langle \sin \epsilon \rangle} \frac{N_+ - N_-}{N_+ + N_-}$$

1156 events

$$\langle S \rangle = 0.565$$

$$\langle \sin \epsilon \rangle = 0.84,$$


$$\alpha = 0.62.$$

Role of polarization physics


- Probing of spin degree freedom
 - Test the dynamic of SM and low energy hadron interaction
 - Existant exp. : RHIC, Jlab, GRAAL, CERN and DESY
 - Spin observable, spin-dependent structure function and parton distribution
 - Spin crisis at eighties
- BEPCII/BESIII, unpolarized beam, inaccessible polarization of final state by BESIII
 - Polarized beam for post-BEPCII options, CPV in tau decay, Hyperon weak decay,.....
 - Useful tool: transverse polarization of hyperon, spontaneous production at e^+e^- collision
 - $\Lambda \rightarrow p\pi^-$ decay plays important role in particle physics

Transverse polarization (TP) of baryons in e^+e^- collisions

FFs in continuum



J/ψ decays



Time likespin $\frac{1}{2}$ baryon FFs:

Dubnickova, Dubnicka, Rekaló

Nuovo Cim. A109 (1996) 241

W. Lu, et.al., Phys.Lett., B368, 261 (1996)

Gakh, Tomasi-Gustafsson Nucl.Phys. A771 (2006) 169

Czyz, Grzelinska, Kuhn PRD75 (2007) 074026

Fäldt EPJ A51 (2015) 74; EPJ A52 (2016)141


Fäldt, G. & Kupsc, A, Phys. Lett. B 772 (2017) 16

$$\Gamma_{\mu}^e(k_1, k_2) = -ie_{\psi} \gamma_{\mu}$$


$$\Gamma_{\mu}^{\Lambda}(p_1, p_2) =$$

$$-ie_g \left[G_M^{\psi} \gamma_{\mu} - \frac{2M}{Q^2} (G_M^{\psi} - G_E^{\psi}) Q_{\mu} \right]$$

Transverse polarization of baryons in e^+e^- collisions




Unpolarized e^+e^- beam \rightarrow
transverse polarization baryon



$$P_Y = \frac{\sqrt{1 - \alpha_\psi^2} \cos \theta_0 \sin \theta_0}{1 + \alpha_\psi \cos^2 \theta_0} \sin(\Delta\Phi)$$

P_Y along $\mathbf{k}_{e^+} \times \mathbf{p}_\Lambda$

$\Lambda \rightarrow p\pi^-$ as polarimeter



$$\frac{dN}{d\Omega} = \frac{1}{4\pi} (1 + \alpha_{\Lambda} \vec{P} \cdot \hat{q}) = \frac{1}{4\pi} (1 + \alpha_{\Lambda} P_{\Lambda} \cos\theta_p)$$

Lee-Yang parameters:

$$\alpha = \frac{2 \operatorname{Re}(S^* P)}{|S|^2 + |P|^2}, \quad \beta = \frac{2 \operatorname{Im}(S^* P)}{|S|^2 + |P|^2}, \quad \gamma = \frac{|S|^2 - |P|^2}{|S|^2 + |P|^2}$$

Note: $\alpha^2 + \beta^2 + \gamma^2 = 1$

PDG2018: $\alpha_{\Lambda} = 0.642 \pm 0.013, \alpha_{\bar{\Lambda}} = -0.71 \pm 0.08$