

JUNO实验探测 ${}^8\text{B}$ 太阳中微子的物理潜力

赵洁


中国科学院高能物理研究所

On behalf of the JUNO collaboration

2021-8-17 @ 第十三届全国粒子物理学学术会议

探测低能⁸B太阳中微子的物理潜力 @JUNO


■ Matter-vacuum Transition Phase ($E > 2$ MeV)




JUNO detector

20 kt LS, $3\%/\sqrt{E}$ energy resolution

- ✓ Afford excellent self-shielding
- ✓ Comparable statistics to Super-K
- ✓ Expected 2 MeV detection threshold




■ Solar & reactor measurement in Δm_{21}^2 using one single detector



2.2σ (2018) $\rightarrow 1.4 \sigma$ (2020)


SK results @ Neutrino2020



探测低能⁸B太阳中微子的物理潜力 @JUNO



■ Matter-vacuum Transition Phase




■ Two main observables:

- ✓ ES channel ($\nu + e \rightarrow \nu + e$): $E_{\text{vis}} > 2 \text{ MeV}$
- ✓ Day-night asymmetry

■ Challenge: Cosmogenic & Radioactivity


- ✓ LS radioactivity: 10^{-17} g/g (solar phase)
- ✓ Optimized fiducial volume
- ✓ Better muon veto approach

■ Solar & reactor measurement in Δm_{21}^2 using one single detector



2.2 σ (2018) \rightarrow 1.4 σ (2020)

SK results @ Neutrino2020



预期谱计算

■ Neutrino flux and spectrum

- ✓ Flux: $(5.25 \pm 0.20) \times 10^6 \text{ /cm}^2/\text{s}$ from SNO NC measurement (*Phys.Rev.C* 88 (2013) 025501)
- ✓ Spectra: <http://www.sns.ias.edu/~jnb>




■ Detection

In this talk

✓ Main channel: $\nu\text{-e ES}$ $\nu_\alpha + e^- \rightarrow \nu_\alpha + e^-$

✓ Other channel: $\nu_e + ^{13}\text{C CC}$ $\nu_e + ^{13}\text{C} \rightarrow e^- + ^{13}\text{N}$

$\nu + ^{13}\text{C NC}$ $\nu + ^{13}\text{C} \rightarrow \nu + ^{13}\text{C} + \gamma$




内禀天然放射性本底


■ ~2 MeV threshold achievable

- ✓ ^{210}Bi , ^{40}K are not easy to be removed.
- ✓ Alpha energy after quenching is $< 2 \text{ MeV}$
- ✓ ^{214}Bi , ^{212}Bi and ^{208}Tl decays can be removed by cascade decays.


- Well discrimination on α/β with PMT shape information



$^{238}\text{U}: 10^{-17} \text{ g/g}$
 $^{232}\text{Th}: 10^{-17} \text{ g/g}$
 $^{40}\text{K}: 10^{-18} \text{ g/g}$
 $^{210}\text{Pb}: 10^{-24} \text{ g/g}$
 $^{14}\text{C}: 10^{-17} \text{ g/g}$
 $^{39}\text{Ar}/^{85}\text{Kr}: 1\text{uBq/m}^3$
 $^{210}\text{Po}: 2600 \text{ cpd/kt}$




After reduction



外部天然放射性本底

External radioactivity

Main materials	Measurement (U/Th)	Information
Acrylic	< 1 ppt	Measured by NAA (LRT2019) and ICP-MS (arXiv2011.06817)
Stainless steel	< 1 ppb	Same supplier with Daya Bay
PMT glass	~200 ppb	Ref: NIMA 898 (2018) 67–71
Water	Radon < 0.1 Bq/m ³	Ref: RDTM (2018) 2:48
Rock	10~30 ppm	4 m water and 5 mm HDPE shielding, negligible



External neutron captures

- ✓ High energy gamma: neutron captures on iron, PMT glass and acrylic
- ✓ Residual: < 0.001 cpd with R < 16.5 m

Final optimized fiducial volume cut


- ✓ With about 5 m self-shielding, ^{208}Tl decays from external materials are negligible.

	R < 16.5 m	R < 15 m	R < 13 m
Energy (MeV)	> 5	(3, 5)	(2, 3)
Target mass (kt)	16.2	12.2	7.9

宇宙同位素本底

Muon rate

Overburden	Muon flux	$\langle E_\mu \rangle$	R_μ in LS	R_μ in water
~700 m	0.004 Hz/m ²	207 GeV	3.6 Hz	10 Hz





Optimized veto strategy

✓ Muon veto:

- Whole detector veto for non-track μ
- Cylinder veto around μ track
 - The nearer distance, the longer veto time
- Dead time: ~44%

✓ Three-Fold Coincidence cut (TFC):


- TFC: Muon, spallation neutron capture, isotope decay
- TFC fractions: > 90%
- Spherical veto around spallation neutron
- Dead time ~4%



宇宙同位素本底

Spectrum

Yield in LS is scaled from measurement by Borexino and KamLAND



After reduction

Systematics

Systematic uncertainty	^{12}B	^8Li	^6He	^{10}C	^{11}Be
livetime (τ)	29.1ms	1.21 s	1.16s	27.8 s	19.9s
KamLAND [47]	—	3.3%	—	—	10.8%
JUNO	1%	1~3%	1~3%	5~10%	5~10%

- ✓ The actual isotope yields, distance distributions, and TFC fractions will be measured *in-situ* in future.

本底和信号小结


■ ${}^8\text{B}$ signal efficiency

- ✓ Muon veto efficiency: 52%
- ✓ (3, 5) MeV: 52% (muon veto) * 80% (${}^{212}\text{Bi}$ - ${}^{208}\text{Tl}$ cut)

■ Other systematics

- ✓ FV cut: 1%, refer to Borexino
(*Phys. Rev. D*, 101(6):062001, 2020).
- ✓ Detector energy scale: 0.3%, refer to Daya Bay
(*Nucl. Instrum. Meth. A*, 940:230-242, 2019.)

The discontinuities at 3 MeV and 5 MeV come from the different FV cut.



$$\Delta m_{21}^{2*} = 4.8 \times 10^{-5} \text{ eV}^2, \text{ and } \Delta m_{21}^{2\dagger} = 7.5 \times 10^{-5} \text{ eV}^2$$

cpd/kt	FV	${}^8\text{B}$ signal eff.	${}^{12}\text{B}$	${}^8\text{Li}$	${}^{10}\text{C}$	${}^6\text{He}$	${}^{11}\text{Be}$	${}^{238}\text{U}$	${}^{232}\text{Th}$	$\bar{\nu}$ -e ES	Total bkg.	Signal rate at Δm_{21}^{2*}	Signal rate at $\Delta m_{21}^{2\dagger}$
(2, 3) MeV	7.9 kt	~51%	0.005	0.006	0.141	0.084	0.002	0.050	0.050	0.049	0.39	0.32	0.30
(3, 5) MeV	12.2 kt	~41%	0.013	0.018	0.014	0.008	0.005	0	0.012	0.016	0.09	0.42	0.39
(5, 16) MeV	16.2 kt	~52%	0.065	0.085	0	0	0.023	0	0	0.002	0.17	0.61	0.59
Syst. error	1%	<1%	3%	10%	3%	10%	1%	1%	2%				



Background after reduction, and muon veto efficiency corrected .

谱型畸变(Matter-vacuum Transition Phase)

Systematics

	Notation	Value	Reference
Detection efficiency	σ_d	1%	Borexino [23]
Detector energy scale	σ_e	0.3%	Daya Bay [35], Double Chooz [65]
The ${}^8\text{B}$ ν flux	σ_f	3.8%	SNO [25]
The ${}^8\text{B}$ ν spectrum shape	σ_s	1	Bahcall [26]
The j^{th} background	σ_b^j	Table 4	This study

Energy-independent hypothesis: flat P_{ee} for $E > 2 \text{ MeV}$



✓ This hypothesis is rejected at 2.7σ with large Δm_{21}^2 , and the statistic-only sensitivity can reach 5σ .



日夜效应

■ Day-night-asymmetry $DNA = \frac{R_D - R_N}{(R_N + R_D)/2}$

- ✓ R_N, R_D are background-subtracted signal rates during the Day ($\cos\theta_z < 0$) and Night ($\cos\theta_z > 0$)

■ Sensitivity (10 years data)

- ✓ Most systematics are cancelled, statistics dominate






Day/Night asymmetry: *Super-K results @ Neutrino2020*

$$A_{DN}^{Fit} = (-3.6 \pm 1.6(stat) \pm 0.6(syst)) \% \rightarrow A_{DN}^{Fit} = (-2.1 \pm 1.1) \%$$

- ✓ JUNO will reach SK 20y uncertainty (1.1%) in less than ten years due to the better S/B ratio.

振荡参数测量

■ χ^2 fitting with both energy and sun-detector angle



- ✓ Statistics/flux dominate the uncertainty of $\Delta m_{21}^2 / \sin^2 \theta_{12}$
- ✓ the discrimination on Δm_{21}^2 from $4.8 \times 10^{-5} \text{ eV}^2$ to $7.5 \times 10^{-5} \text{ eV}^2$ can reach 2σ .

■ Discussion on worse conditions

- ✓ ${}^{210}\text{Po}$ reach Phase I of Borexino (10^4 cpd/kt), ${}^{208}\text{Tl}$ could not be reduced. Dashed line in the right panel
- ✓ ${}^{238}\text{U}/{}^{232}\text{Th} \sim 10^{-15} \text{ g/g}$, $E_{\text{vis}} > 5 \text{ MeV}$. Dotted line in the right panel

总结

■ 详细研究了本底压低策略

- ✓ 内禀天然放射性本底: Bi-Po, Bi-Tl 级联衰变标记
- ✓ 外部天然放射性本底: 不同能量区域实行优化的靶体积
- ✓ 宇生同位素本底: muon 径迹 & 中子三重符合

■ JUNO 探测⁸B 太阳中微子的物理潜力 (10 years data)

- ✓ 信号 60,000, 本底 30,000, $E_{\text{th}} \sim 2 \text{ MeV}$
- ✓ 排除平的 P_{ee} vs. E : $\geq 2\sigma$ for the large Δm_{21}^2 ($7.5 \times 10^{-5} \text{ eV}^2$)
- ✓ 日夜不对称性测量精度: 0.9%, better than Super-K's latest result 1.1%
- ✓ 用同一探测器, 不同中微子源 (太阳, 反应堆), 测量振荡参数 Δm_{21}^2
 - the discrimination on Δm_{21}^2 from $4.8 \times 10^{-5} \text{ eV}^2$ to $7.5 \times 10^{-5} \text{ eV}^2$ can reach 2σ .
- ✓ 合作组文章已发表: *CPC Vol. 45, No.2 (2021)*

谢谢聆听!



Back up

Internal background



■ Strategy for Background reduction

- ✓ With correlation cuts (time, position, energy) most of ^{214}Bi , ^{212}Bi and ^{208}Tl decays can be removed

	^{214}Bi - ^{214}Po - ^{210}Pb	^{212}Bi - ^{212}Po - ^{208}Pb	^{212}Bi - ^{208}Tl - ^{208}Pb
Prompt signal	Beta (2, 3.5) MeV	Beta (2, 3.5) MeV	Alpha (0.5, 0.7) MeV
Delayed signal	Alpha (0.85, 1.05) MeV	Alpha (0.85, 1.05) MeV	Beta (3, 5) MeV
Time & position correlation	2 ms & 2 m	5 μs & 2 m	22 min & 1 m

Visible energy

Isotopes in the ^{238}U and ^{232}Th decay chains with decay energies larger than 2 MeV


Isotope	Decay mode	Decay energy	τ	Daughter	Daughter's τ	Removal eff.	Removed signal
^{214}Bi	β^-	3.27 MeV	28.7 min	^{214}Po	237 μs	>99.5%	<1%
^{212}Bi	β^- : 64%	2.25 MeV	87.4 min	^{212}Po	431 ns	93%	~0
^{212}Bi	α : 36%	6.21 MeV	87.4 min	^{208}Tl	4.4 min	N/A	N/A
^{208}Tl	β^-	5.00 MeV	4.4 min	^{208}Pb	Stable	99%	20%
$^{234}\text{Pa}^m$	β^-	2.27 MeV	1.7 min	^{234}U	245500 years	N/A	N/A
^{228}Ac	β^-	2.13 MeV	8.9 h	^{228}Th	1.9 years	N/A	N/A

External radioactivity and neutron captures



External radioactivity

- ✓ Simulation is done with JUNO offline software
- ✓ Contribution from rock is negligible with at least 4 m water and 5 mm HDPE liner shielding
- ✓ Effectively removed by the fiducial volume (FV) cut



External neutron captures

- ✓ High energy gamma: neutron captures on iron, PMT glass and acrylic

Unit: cpd	R < 17.7 m	R < 17 m	R < 16 m	R < 15 m	R < 14 m
Acrylic	1.3e-5	1.53e-7	2.7e-9	5.57e-11	0
Node	3.52e-5	8.9e-7	5.1e-8	4.1e-9	4.3e-10
Bar	9.8e-5	2.6e-6	1.8e-7	1.2e-8	2.7e-9
PMT glass	2.5e-2	1.8e-4	7.2e-6	0	0
Total	0.025	1.9e-4	7.5e-6	1.6e-8	3.1e-9

- ✓ PMT glass dominates the contribution, but can be effectively remove with R < 16.5 m FV cut

Final optimized fiducial volume cut

	R < 16.5 m	R < 15 m	R < 13 m
Energy (MeV)	> 5	(3, 5)	(2, 3)
Target mass (kt)	16.2	12.2	7.9

Long-lived cosmogenic isotopes




Muon rate

- ✓ With ~ 700 m overburden, muon flux is 0.004 Hz/m 2 with an average energy of 207 GeV.
- ✓ About 3.6/10 Hz muons passing through the LS/water

Yield

- ✓ Three-Fold Coincidence cut (TFC) among muon, spallation neutron capture and isotope decay
Used in this study



Isotope	Decay mode	Decay energy (MeV)	τ	Yield in LS (/day)		TFC fraction
				Geant4 simulation	Scaled	
¹² B	β^-	13.4	29.1 ms	1059	2282	90%
⁹ Li	β^- : 50%	13.6	257.2 ms	68	117	96%
⁹ C	β^+	16.5	182.5 ms	21	160	>99%
⁸ Li	$\beta^- + \alpha$	16.0	1.21 s	725	649	94%
⁶ He	β^-	3.5	1.16 s	526	2185	95%
⁸ B	$\beta^+ + \alpha$	~18	1.11 s	35	447	>99%
¹⁰ C	β^+	3.6	27.8 s	816	878	>99%
¹¹ Be	β^-	11.5	19.9 s	9	59	96%
¹¹ C	β^+	1.98	29.4 min	11811	46065	98%


Scaled: scale from measurement by Borexino and KamLAND

Long-lived cosmogenic isotopes



■ Optimized veto strategy

	For reconstructed track	Strategy	Dead time
Whole detector veto	w/ track	Veto 2 ms	44%
	w/o track	Veto 1 s	
Cylindrical volume veto (d : distance between candidate to muon track)	w/ track	Veto $d < 1$ m for 5 s Veto $1 \text{ m} < d < 3$ m for 4 s Veto $3 \text{ m} < d < 4$ m for 2 s Veto $4 \text{ m} < d < 5$ m for 0.2 s	
TFC veto		Veto 2 m spherical volume around neutron for 160 s	4%



Differential cross section of ν_e and $\nu_{\mu\tau}$




Figure 4: Differential cross section of $\nu_e - e$ (blue) and $\nu_{\mu,\tau} - e$ (black) elastic scattering for a 10 MeV neutrino. The stronger energy dependence of the $\nu_{\mu,\tau} - e$ cross section, as illustrated in red, produces another smooth upturn in the visible electron spectrum compared to the case of no $\nu_{\mu,\tau}$ appearance.