


$B_c^+ \rightarrow \tau^+ \nu_\tau$ Analysis

Taifan


Basic statistics

The following studies are done assuming 1×10^{11} Z bosons are produced. $\rightarrow 7.5 \times 10^5 B_c^+ \rightarrow \tau^+ \nu_\tau$ & $1 \times 10^6 B^+ \rightarrow \tau^+ \nu_\tau$ (according to the Soeren's slides. Not considering uncertainties)


Strategy




Parameter distribution



btag>0.6




Recoil side en –
signal side en >
10 GeV




Only one lepton
in the signal
hemisphere

Parameter distribution



That lepton is
electron




And it's the most energetic
particle in the signal hemisphere

Cut chain


	$B_c^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e$	$B^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e$	1/6100 ↓ dd	1/4900 ↓ uu	1/6100 ↓ ss	1/1100 ↓ cc	1/1500 ↓ bb
All	625177/134681	797230/195570	2530406	2415827	2531430	10414223	10532756
b-tag > 0.6	437048/94370	536144/133336	12495	11559	14920	590417	7885422
Energy asymmetry > 10 GeV	361063/83338	433750/119520	2048	1857	2525	108464	1892666
One lepton in sig hemi	127468/44500	153697/61805	610	549	784	38263	623432
Which is electron	32044/42386	30916/58652	206	181	245	16107	287334
And it's the most energetic one	2569/32458	2173/42475	8	5	4	2449	93945

BDT

Variable	Importance
Nominal B_c energy	0.201
The second largest momentum in sig hemi	0.151
Maximum neutral cluster energy inside 30 deg cone	0.151
Energy asymmetry	0.148
Electron energy	0.123
Second largest IP in sig hemi	0.120
Number of tracks in sig hemi	0.106



root 5.34.07
 ↑
 I chose this &
 cut at 0.99



root 5.34.18
 Set the weight to corresponding luminosity

Cut chain


	$B_c^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e$	$B^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e$	1/6100 ↓	1/4900 ↓	1/6100 ↓	1/1100 ↓	1/1500 ↓
	$B_c^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e$	$B^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e$	dd	uu	ss	cc	bb
All	625177/134681	797230/195570	2530406	2415827	2531430	10414223	10532756
b-tag > 0.6	437048/94370	536144/133336	12495	11559	14920	590417	7885422
Energy asymmetry > 10 GeV	361063/83338	433750/119520	2048	1857	2525	108464	1892666
One lepton in sig hemi	127468/44500	153697/61805	610	549	784	38263	623432
Which is electron	32044/42386	30916/58652	206	181	245	16107	287334
And it's the most energetic one	2569/32458	2173/42475	8	5	4	2449	93945
BDT > 0.99 (training data)	226/7226	65/5150	0	0	0	0	9 (5884)
BDT > 0.99 (test data)	223/7142	87/5178	0	0	0	1	8 (7441)

50/50 split between test and train samples


BDT

Using all of the previous variables + electron IP to do BDT again (ignore bb)

Variable	Importance
Electron IP	0.164
Electron energy	0.138
Nominal Bc energy	0.137
Energy asymmetry	0.134
Maximum neutral cluster energy inside 30 deg cone	0.133
The second largest momentum in sig hemi	0.127
Second largest IP in sig hemi	0.086
Number of tracks in sig hemi	0.082



Fit IP with exponential



Alternatively, we can fit the IP with exponential $C_1 \cdot e^{-E_1 x} + C_2 \cdot e^{-E_2 x}$ (again, ignoring bb).

EXT PARAMETER					
NO.	NAME	VALUE	ERROR	STEP SIZE	FIRST DERIVATIVE
1	C1	5.72575e+02	1.52615e+01	1.44289e-02	-7.51254e-06
2	E1	2.41352e-01	6.62978e-03	6.26832e-06	1.43783e-02
FCN=67.8532 FROM MIGRAD		STATUS=CONVERGED		80 CALLS	81 TOTAL
		EDM=5.09469e-08		STRATEGY= 1	ERROR MATRIX ACCURATE
EXT PARAMETER					
NO.	NAME	VALUE	ERROR	STEP SIZE	FIRST DERIVATIVE
1	C2	2.24564e+03	6.53438e+01	1.14074e-01	1.07633e-05
2	E2	7.25760e-01	1.12363e-02	1.96105e-05	-6.54741e-02
FCN=21.3704 FROM MIGRAD		STATUS=CONVERGED		327 CALLS	328 TOTAL
		EDM=7.43106e-08		STRATEGY= 1	ERROR MATRIX ACCURATE
EXT PARAMETER					
NO.	NAME	VALUE	ERROR	STEP SIZE	FIRST DERIVATIVE
1	C1	7.27399e+02	2.01951e+02	2.26757e-02	-5.88938e-06
2	E1	2.49481e-01	3.81610e-02	6.94000e-06	-2.18549e-03
3	C2	2.83089e+03	1.83058e+02	1.31003e-01	-4.50623e-06
4	E2	9.38499e-01	1.24337e-01	2.44118e-05	1.31922e-02

Number of events predicted by fit results


	Truth	Individual fit	Combined fit
$B_c^+ \rightarrow \tau^+ \nu_\tau, \tau^+ \rightarrow e \nu \nu$	6438	6525	5055
$B^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e \nu \nu$	6528	6375	8047

Cut chain ($\tau \rightarrow \mu\nu\nu$)

	$B_c^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow \mu$	$B^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow \mu$	1/6100 dd	1/4900 uu	1/6600 ss	1/1100 cc	1/1500 bb
All	628199/131659	802123/190677	2530406	2415827	2331430	10014223	10332758
b-tag > 0.6	439038/92380	539792/129688	12495	11559	13808	567891	7736169
Energy asymmetry > 10 GeV	363241/81160	437202/116068	2048	1857	2342	104333	1856916
One lepton in sig hemi	130225/41743	158052/57450	610	549	729	36843	611737
Which is muon	57721/39817	70506/55428	404	368	498	21313	329706
And it's the most energetic one	5820/31200	5173/40799	12	11	13	2779	98136

BDT ($\tau \rightarrow \mu\nu\nu$)

Variable	Importance
Nominal Bc energy	0.210
Maximum neutral cluster energy inside 30 deg cone	0.157
Energy asymmetry	0.151
The second largest momentum in sig hemi	0.141
Second largest IP in sig hemi	0.128
Electron energy	0.123
Number of tracks in sig hemi	0.089



root 5.34.07
↑
cut at 0.99

Set the weight to corresponding luminosity

Cut chain ($\tau \rightarrow \mu\nu\nu$)


	$B_c^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow \mu$	$B^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow \mu$	1/6100 ↓ dd	1/4900 ↓ uu	1/6600 ↓ ss	1/1100 ↓ cc	1/1500 ↓ bb
All	628199/131659	802123/190677	2530406	2415827	2331430	10014223	10332758
b-tag > 0.6	439038/92380	539792/129688	12495	11559	13808	567891	7736169
Energy asymmetry > 10 GeV	363241/81160	437202/116068	2048	1857	2342	104333	1856916
One lepton in sig hemi	130225/41743	158052/57450	610	549	729	36843	611737
Which is muon	57721/39817	70506/55428	404	368	498	21313	329706
And it's the most energetic one	5820/31200	5173/40799	12	11	13	2779	98136
BDT > 0.99 (training data)	756/7394	486/5170	0	0	0	2	9 (5598)
BDT > 0.99 (test data)	763/7313	455/5266	0	0	0	0	6 (5427)

50/50 split between test and train samples


BDT ($\tau \rightarrow \mu\nu\nu$)

Using all of the previous variables + electron IP to do BDT again (ignore bb)

Variable	Importance
Electron IP	0.159
Electron energy	0.143
Energy asymmetry	0.132
Maximum neutral cluster energy inside 30 deg cone	0.132
Nominal B meson energy	0.129
The second largest momentum in sig hemi	0.129
Number of tracks in sig hemi	0.091
Second largest IP in sig hemi	0.085



Fit IP with exponential ($\tau \rightarrow \mu\nu\nu$)



Alternatively, we can fit the IP with exponential $C_1 \cdot e^{-E_1 x} + C_2 \cdot e^{-E_2 x}$ (again, ignoring bb).

NO.	NAME	VALUE	ERROR	SIZE	DERIVATIVE
1	C1	5.51094e+02	1.45324e+01	1.95868e-02	-1.28790e-05
2	E1	2.35308e-01	6.48465e-03	8.73984e-06	1.71527e-02
FCN=64.654 FROM MIGRAD STATUS=CONVERGED 78 CALLS 79 TOTAL					
EDM=1.80213e-09 STRATEGY= 1 ERROR MATRIX ACCURATE					
EXT PARAMETER STEP FIRST					
NO.	NAME	VALUE	ERROR	SIZE	DERIVATIVE
1	C2	2.24316e+03	6.59072e+01	1.12545e-01	1.57326e-06
2	E2	7.35570e-01	1.14151e-02	1.94939e-05	-4.68803e-03
FCN=29.2032 FROM MIGRAD STATUS=CONVERGED 336 CALLS 337 TOTAL					
EDM=2.65854e-07 STRATEGY= 1 ERROR MATRIX ACCURATE					
EXT PARAMETER STEP FIRST					
NO.	NAME	VALUE	ERROR	SIZE	DERIVATIVE
1	C1	7.79201e+02	2.46309e+02	2.78175e-02	-1.86941e-05
2	E1	2.59603e-01	4.23892e-02	8.04291e-06	1.27713e-02
3	C2	2.66687e+03	1.67475e+02	1.44562e-01	-2.55748e-06
4	E2	9.03294e-01	1.34425e-01	2.80029e-05	2.87191e-02


Number of events predicted by fit results

	Truth	Individual fit	Combined fit
$B_c^+ \rightarrow \tau^+ \nu_\tau, \tau^+ \rightarrow \mu \nu \nu$	6284	6220	5121
$B^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow \mu \nu \nu$	6437	6410	8715


Training with xgboost ($\tau \rightarrow e\nu\nu$)

Variable	Importance
Nominal Bc energy	126
Maximum neutral cluster energy inside 30 deg cone	59
Energy asymmetry	58
The second largest momentum in sig hemi	43
Second largest IP in sig hemi	58
Impact parameter	39

Electron energy and #track in sig hemi are dropped due to low importance



Logarithmic y axis




Linear y axis


BDT response (training data)

Set the weight to the corresponding luminosity


Training with xgboost (train & test data overlay)




Signal



$B^+ \rightarrow \tau\nu, \tau \rightarrow e$




$B_c^+ \rightarrow \tau\nu$
(excl. $\tau \rightarrow e$)




$B^+ \rightarrow \tau\nu$
(excl. $\tau \rightarrow e$)

Training with xgboost (train & test data overlay)



cc





bb

Cut chain(xgboost, $\tau \rightarrow e\nu\nu$)

	$B_c^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e$	$B^+ \rightarrow \tau^+ \nu_\tau / \tau^+ \rightarrow e$	1/6100 dd	1/4900 uu	1/6100 ss	1/1100 cc	1/1500 bb
All	625177/134681	797230/195570	2530406	2415827	2531430	10414223	10532756
b-tag > 0.6	437048/94370	536144/133336	12495	11559	14920	590417	7885422
Energy asymmetry > 10 GeV	361063/83338	433750/119520	2048	1857	2525	108464	1892666
One lepton in sig hemi	127468/44500	153697/61805	610	549	784	38263	623432
Which is electron	32044/42386	30916/58652	206	181	245	16107	287334
And it's the most energetic one	2569/32458	2173/42475	8	5	4	2449	93945
BDT > 0.45 (training data)	200/8353	50/2806	0	0	0	0	0
BDT > 0.45 (test data)	247/7651	57/3232	0	0	0	0	0

50/50 split between test and train samples

Kinematic fit with pseudo track ($\tau \rightarrow e\nu\nu$)






χ^2

Distance (reco vtx – decay point)

Kinematic fit of $K_s^0 \rightarrow \pi^+\pi^-$ decay point
(just to check if it works)

Kinematic fit with pseudo track ($\tau \rightarrow e\nu\nu$)

Use the all of the particles on the recoil side to build a pseudo track that passes through the interaction point. And fit the vertex with the electron. Set $\text{cov}(\text{pseudo track}) = \text{cov}(\text{electron track})$.



$$\text{Relative error(IP)} = \text{abs}\left(\frac{\text{IP}.Mag() - (\tau \text{ end}).Mag()}{(\tau \text{ end}).Mag()}\right)$$

Pre-cut on the additional bkg samples ($\tau \rightarrow e\nu\nu$)

Reject obvious backgrounds at generator level.

bb:

1. Only one electron in the signal hemisphere
2. The electron is the most energetic particle in the signal hemisphere
3. Nominal B meson energy > 10 GeV

Results:

1. 4.6×10^8 events $\rightarrow 3.7 \times 10^7$ events (92% reduction)
2. bb survive rate after cut chain 0.9% \rightarrow 0.7%