

# Correlated Dirac Eigenvalues and Axial Anomaly in Chiral Symmetric QCD

*Sunday, 31 October 2021 16:50 (30 minutes)*

In this talk I will present the novel relations between the quark mass derivatives  $[\partial^n \rho(\lambda, m_l) / \partial m_l^n]$  of the Dirac eigenvalue spectrum and the  $(n + 1)$ -point correlations among the eigenvalues. Using these relations we present lattice QCD results for  $\partial^n \rho(\lambda, m_l) / \partial m_l^n$  ( $n = 1, 2, 3$ ) for  $m_l$  corresponding to pion masses  $m_\pi = 160 - 55$  MeV, and at a temperature of about 1.6 times the chiral phase transition temperature. Calculations were carried out using (2+1) flavors of highly improved staggered quarks with the physical value of strange quark mass, three lattice spacings  $a = 0.12, 0.08, 0.06$  fm. We find that  $\rho(\lambda \rightarrow 0, m_l)$  develops a peaked structure. This peaked structure arises due to non-Poisson correlations within the infrared part of the Dirac eigenvalue spectrum, becomes sharper as  $a \rightarrow 0$ , and its amplitude is proportional to  $m_l^2$ . We demonstrate that this  $\rho(\lambda \rightarrow 0, m_l)$  is responsible for the manifestations of axial anomaly in two-point correlation functions of light scalar and pseudoscalar mesons. After continuum and chiral extrapolations we find that axial anomaly remains manifested in two-point correlation functions of scalar and pseudoscalar mesons in the chiral limit. This talk is based on PRL 126 (2021) 082001.

**Presenter:** Dr 张, 瑜

**Session Classification:** session4