

Recent results from **BESIII** experiment

刘北江

中国科学院高能物理研究所

(On behalf of the BESIII collaboration)


中国格点QCD第一届年会, 华南师范大学

2021.10.30-2021.11.02


World's largest τ – charm data sets in e^+e^- annihilation

Beijing Electron Positron Collider (BEPCII)

beam energy: 1.0 – 2.3(2.45) GeV



- 1989-2004 (BEPC):
 $L_{\text{peak}}=1.0 \times 10^{31} / \text{cm}^2 \text{s}$
- 2004: started BEPCII upgrade, BESIII construction
- 2008: test run
- 2009-now (BEPCII):
 $L_{\text{peak}}=1.0 \times 10^{33} / \text{cm}^2 (4/5/2016)$
 2020: energy upgrade to 2.45 GeV & top-up mode




>> 1×10^5 had. Events each

Selected topics


- Light hadrons: glueballs & more
- XYZ particles: Y(4260), X(3872), Z_{cs}(3985)
- Charm decays: CKM, decay constants, form factors, LFU
- Hadronic corrections to muon g-2 : HPV & HLbL
- Baryons: form factors & polarization

Charmonium decays provide an ideal lab for light hadron physics



What's the role of gluonic excitation and how does it connect to the confinement?

- Clean high statistics data samples
- Well defined initial and final states
 - Kinematic constraints
 - $I(J^{PC})$ filter
- “Gluon-rich” process



$$\Gamma(J/\psi \rightarrow \gamma G) \sim O(\alpha\alpha_s^2), \Gamma(J/\psi \rightarrow \gamma H) \sim O(\alpha\alpha_s^3),$$

$$\Gamma(J/\psi \rightarrow \gamma M) \sim O(\alpha\alpha_s^4), \Gamma(J/\psi \rightarrow \gamma F) \sim O(\alpha\alpha_s^4)$$

Scalar glueball candidate

$$\Gamma(J/\psi \rightarrow \gamma G_{0+}) = \frac{4}{27} \alpha \frac{|p|}{M_{J/\psi}^2} |E_1(0)|^2 = 0.35(8) \text{ keV}$$


$$\Gamma/\Gamma_{tot} = 0.33(7)/93.2 = 3.8(9) \times 10^{-3}$$

CLQCD, *Phys. Rev. Lett.* 110, 021601 (2013)



Experimental results

- $B(J/\psi \rightarrow \gamma f_0(1710) \rightarrow \gamma K\bar{K}) = (8.5_{-0.9}^{+1.2}) \times 10^{-4}$
 - $B(J/\psi \rightarrow \gamma f_0(1710) \rightarrow \gamma \pi\pi) = (4.0 \pm 1.0) \times 10^{-4}$
 - $B(J/\psi \rightarrow \gamma f_0(1710) \rightarrow \gamma \omega\omega) = (3.1 \pm 1.0) \times 10^{-4}$
 - $B(J/\psi \rightarrow \gamma f_0(1710) \rightarrow \gamma \eta\eta) = (2.35_{-0.11}^{+0.13} {}_{-0.74}^{+1.24}) \times 10^{-4}$
- ⇒ $B(J/\psi \rightarrow \gamma f_0(1710)) > 1.7 \times 10^{-3}$



$f_0(1710)$ largely overlapped with scalar glueball

Recent interpretations

Scalar isoscalar mesons and the scalar glueball from radiative J/ψ decays

Andrey V. Sarantsev, Igor Denisenko, Ulrike Thoma, Eberhard Klempt

A coupled-channel analysis of BESIII data on radiative J/ψ decays into $\pi\pi$, $K\bar{K}$, $\eta\eta$ and $\omega\phi$ has been performed. The partial-wave amplitude is constrained by a large number of further data. The analysis finds ten isoscalar scalar mesons. Their masses, widths and decay modes are determined. The scalar mesons are interpreted as mainly SU(3)-singlet and mainly octet states. Octet isoscalar scalar states are observed with significant yields only in the 1500-2100 MeV mass region. Singlet scalar mesons are produced over a wide mass range but their yield peaks in the same mass region. The peak is interpreted as scalar glueball. Its mass and width are determined to $M = 1865 \pm 25^{+10}_{-30}$ (rm MeV) and $\Gamma = 370 \pm 50^{+30}_{-20}$ (rm MeV), its yield in radiative J/ψ decays to $(5.8 \pm 1.0) \cdot 10^{-3}$.

Comments: 11 pages, 4 figures

Subjects: **High Energy Physics - Phenomenology (hep-ph)**

DOI: [10.1016/j.physletb.2021.136227](https://doi.org/10.1016/j.physletb.2021.136227)

Cite as: [arXiv:2103.09680](https://arxiv.org/abs/2103.09680) [hep-ph]

Scalar and tensor resonances in J/ψ radiative decays

JPAC Collaboration: A. Rodas, A. Pilloni, M. Albaladejo, C. Fernandez-Ramirez, V. Mathieu, A. P. Szczepaniak

We perform a systematic analysis of the $J/\psi \rightarrow \gamma\pi^0\pi^0$ and $\rightarrow \gamma K_S^0 K_S^0$ partial waves measured by BESIII. We use a large set of amplitude parametrizations to reduce the model bias. We determine the physical properties of seven scalar and tensor resonances in the 1-2.5 GeV mass range. These include the well known $f_0(1500)$ and $f_0(1710)$, that are considered to be the primary glueball candidates. The hierarchy of resonance couplings determined from this analysis favors the latter as the one with the largest glueball component.

Comments: 17 pages, 11 figures + 28 pages of Supplemental Material

Subjects: **High Energy Physics - Phenomenology (hep-ph)**; High Energy Physics - Experiment (hep-ex); Nuclear Theory (nucl-th)


Cite as: [arXiv:2110.00027](https://arxiv.org/abs/2110.00027) [hep-ph]

To-do:


Implement coupled channel analysis in BESIII analysis

The X(2120) and X(2370)

- Observed in $J/\psi \rightarrow \gamma \eta' \pi^+ \pi^-$ at BESIII
[PRL106, 072002 (2011)][PRL117, 042002(2016)]
- Combined analysis of $J/\psi \rightarrow \gamma K^+ K^- \eta'$ and $\gamma K_S K_S \eta'$
- Search for X(2370) in $J/\psi \rightarrow \gamma \eta \eta \eta'$




EPJC80,746(2020)



1.3B

PRD103, 012009(2021)








$$M_{X(2370)} = 2341.6 \pm 6.5(\text{stat.}) \pm 5.7(\text{syst.}) \text{ MeV}/c^2,$$

$$\Gamma_{X(2370)} = 117 \pm 10(\text{stat.}) \pm 8(\text{syst.}) \text{ MeV},$$

- Observation of $X(2370) \rightarrow K \bar{K} \eta'$, 8.3σ
- No evidence of $X(2120) \rightarrow K \bar{K} \eta'$
- No evidence of $X(2370) \rightarrow \eta \eta \eta'$

$\chi(p\bar{p})/\chi(18??)$ from J/ψ radiative decays







PRL117, 042002 (2016)

PRD97, 051101(R)(2018)

$J/\psi \rightarrow \gamma \eta' \pi^+ \pi^-$

$J/\psi \rightarrow \gamma \gamma \phi$





Any relations? What is the role of the $p\bar{p}$ threshold ?

Selected topics

- Light hadrons: glueballs & more
- XYZ particles: Y(4260), X(3872), Z_{cs}(3985)
- Charm decays: CKM, decay constants, form factors, LFU
- Hadronic corrections to muon g-2 : HPV & HLbL
- Baryons: form factors & polarization



Charmonium and exotics at BESIII



direct production of vectors: ψ, Y
 radiative and hadronic transitions to others

$$e^+e^- \rightarrow \pi^+\pi^- J/\psi$$







Compare running at **Belle** and **BaBar**, with one month at **BESIII**!




BESIII: $\sigma^B = 62.9 \pm 1.9 \pm 3.7$ pb

PRL 110, 252001 (2013)


Y(4260) \rightarrow Y(4220) and new Y's



PRD 104, 052012 (2021)






Evidence for $Y(4220) \rightarrow \eta_c \pi^+ \pi^- \pi^0$



$e^+e^- \rightarrow \gamma\chi_{cJ}$ at $\sqrt{s}=3.8-4.6$ GeV

- No signals for $e^+e^- \rightarrow \gamma\chi_{c0}$
- Observations of $e^+e^- \rightarrow \gamma\chi_{c1,2}$

arXiv:2107.03604




- $\gamma\chi_{c1}$: Well describe with conventional charmonium states
- $\gamma\chi_{c2}$: Along with conventional ones, **an additional Y state** is needed

$$M = 4371.7 \pm 7.5 \pm 1.8 \text{ MeV}/c^2, \quad \Gamma = 51.1 \pm 17.6 \pm 1.9 \text{ MeV}$$

- ✓ statistical significance of 5.8σ
- ✓ consistent with the $Y(4360)/Y(4390)$

More X(3872) decay information

- Observation of $X(3872) \rightarrow \pi^0 \chi_{c1}$



BESIII, PRL122, 202001 (2019)

- Observation of $X(3872) \rightarrow \omega J/\psi$


BESIII, PRL 122, 232002 (2019)

- Observation of $X(3872) \rightarrow D^0 \bar{D}^{*0}$




BESIII, PRL 124, 242001 (2020)

- Transition of $X(3872) \rightarrow \gamma J/\psi, \gamma \psi(2S)$


BESIII, PRL 124, 242001 (2020)




$$R = \frac{\text{BF}(X(3872) \rightarrow \gamma \psi(2S))}{\text{BF}(X(3872) \rightarrow \gamma J/\psi)} < 0.59 \text{ at 90\% C.L. , agrees with Belle} (< 2.1), \text{ while challenges Babar} (3.4 \pm 1.1) \text{ and LHCb results} (2.46 \pm 0.70)$$




The Zc Family at BESIII




BESIII $e^+e^- \rightarrow \pi^+ \pi^- J/\psi$




BESIII $e^+e^- \rightarrow \pi^+ (D\bar{D}^*)^-$
 $Z_c(3900)^+$




BESIII $e^+e^- \rightarrow \pi^0 \pi^0 J/\psi$




BESIII $e^+e^- \rightarrow \pi^0 (D\bar{D}^*)^0$
 $Z_c(3900)^0$




BESIII $e^+e^- \rightarrow \pi^+ \pi^- h_c$




BESIII $e^+e^- \rightarrow \pi^+ (D^* \bar{D}^*)^-$
 $Z_c(4020)^+$




BESIII $e^+e^- \rightarrow \pi^0 \pi^0 h_c$



BESIII $e^+e^- \rightarrow \pi^0 (D^* \bar{D}^*)^0$
 $Z_c(4020)^0$



Which is the nature of these states?
If exists, there should be SU(3) counter-part
Zcs state with strangeness





Observation of the $Z_{cs}(3985)^\pm$

$$e^+e^- \rightarrow K^+(D_s^-D^{*0} + D_s^{*-}D^0)$$

PRL126, 102001 (2021)

- Simultaneous fit to the five energy points




| State | Signif. | JP | Mass (MeV) | Width (MeV) |
|----------------|-------------|------|--------------------------------|------------------------------|
| $Z_{cs}(3985)$ | 5.3σ | ?? | $3982.5^{+1.8}_{-2.6} \pm 2.1$ | $12.8^{+5.3}_{-4.4} \pm 3.0$ |
| $Z_{cs}(4000)$ | 15σ | $1+$ | $4003 \pm 6_{-14}^{+4}$ | $131 \pm 15 \pm 26$ |
| $Z_{cs}(4220)$ | 5.9σ | $1+$ | $4216 \pm 24_{-30}^{+43}$ | $233 \pm 52_{-73}^{+97}$ |

High statistics analysis of $e^+e^- \rightarrow K^+K^-J/\psi$ is desirable

Selected topics

- Light hadrons: glueballs & more
- XYZ particles: Y(4260), X(3872), Z_{cs}(3985)
- Charm decays: CKM, decay constants, form factors, LFU
- Hadronic corrections to muon g-2: HPV & HLbL
- Baryons: form factors & polarization



- **Single tag (ST):**
fully reconstruct one D^-

$$\Delta E = E_{D^-} - E_{\text{beam}}$$

$$M_{\text{BC}} = \sqrt{E_{\text{beam}}^2 - |\vec{p}_{D^-}|^2}$$


- **Double tag (DT):**
in the recoil ST $D_{(s)}^-$,
analyze the signal $D_{(s)}^+$

$$MM^2 = E_{\text{miss}}^2 - |\vec{p}_{\text{miss}}|^2$$

$$E_{\text{miss}} = E_{\text{cm}} - \sqrt{|\vec{p}_{D_{(s)}^-}|^2 + M_{D_{(s)}}^2} - E_X$$

$$\vec{p}_{\text{miss}} = -\vec{p}_{D_{(s)}^-} - \vec{p}_X$$

$$U_{\text{miss}} = E_{\text{miss}} - |\vec{p}_{\text{miss}}|$$



- **Single tag (ST):**
fully reconstruct one D_s^-

$$M_{\text{rec}} = \sqrt{\left(E_{\text{cm}} - \sqrt{|\vec{p}_{D_s^-}|^2 + m_{D_s^-}^2}\right)^2 - |\vec{p}_{D_s^-}|^2}$$

ST yield: $N_{\text{ST}}^i = 2 \times N_{D\bar{D}} \times B_{\text{ST}}^i \times \varepsilon_{\text{ST}}^i$


DT yield: $N_{\text{DT}}^i = 2 \times N_{D\bar{D}} \times B_{\text{ST}}^i \times B_{\text{sig}} \times \varepsilon_{\text{ST vs. sig}}^i$

Average eff.: $\bar{\varepsilon}_{\text{sig}} = \frac{\sum_{i=1}^N (N_{\text{ST}}^i \times \varepsilon_{\text{ST vs. sig}}^i / \varepsilon_{\text{ST}}^i)}{\sum_{i=1}^N N_{\text{ST}}^i}$

Absolute Br.

$$B_{\text{sig}} = \frac{N_{\text{DT}}^{\text{tot}}}{N_{\text{ST}}^{\text{tot}} \times \bar{\varepsilon}_{\text{sig}}}$$

Pure leptonic decay



$$\Gamma(D_{(s)}^+ \rightarrow l^+ \nu) = \frac{G_F^2 f_{D_{(s)}^+}^2}{8\pi} |V_{cd(s)}|^2 m_l^2 m_{D_{(s)}^+} \left(1 - \frac{m_l^2}{m_{D_{(s)}^+}^2}\right)^2$$

- Decay constant $f_{D_{(s)}^+}$

Exp. decay rate + $|V_{cs(d)}|^{CKM\text{fitter}}$ \rightarrow calibrate LQCD @charm & extrapolate to Beauty

- CKM matrix element $|V_{cs(d)}|$


Exp. decay rate + LQCD \rightarrow CKM matrix elements

$$D_s^+ \rightarrow \mu^+ \nu_\mu \text{ and } D_s^+ \rightarrow \tau^+ \nu_\tau \text{ via } \tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$$

• An unbinned **simultaneous** maximum likelihood fit to **two-dimensional** distributions


Only show @ 4.178 GeV

For all data samples



$$N_{D_s^+ \rightarrow \mu^+ \nu_\mu}^{\text{signal}} = 2198 \pm 55$$

$$N_{D_s^+ \rightarrow \tau^+ \nu_\tau}^{\text{signal}} = 946^{+46}_{-45}$$



The most precise result to date

$$B(D_s^+ \rightarrow \mu^+ \nu_\mu) = (5.35 \pm 0.13_{\text{stat.}} \pm 0.16_{\text{syst.}}) \times 10^{-3}$$


$$B(D_s^+ \rightarrow \tau^+ \nu_\tau) = (5.21 \pm 0.25_{\text{stat.}} \pm 0.17_{\text{syst.}}) \times 10^{-2}$$

$$D_s^+ \rightarrow \tau^+ \nu_\tau \text{ via } \tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$$

✓ $E_{\text{extra}}^{\text{tot}}$: the total energy of the good EMC showers, excluding those associated with the ST D_s^- candidates and those within 5° of the initial direction of the positron.

✓ DT yield $N_{\text{DT}} = N_{\text{DT}}^{\text{tot}} - N_{\text{DT}}^{\text{non-}D_s^-} - N_{\text{DT}}^{K_L^0 e^+ \nu_e} - N_{\text{DT}}^{\chi e^+ \nu_e}$

(in signal $E_{\text{extra}}^{\text{tot}} < 0.4 \text{ GeV}$)



Phys. Rev. Lett. 127 (2021) 171801

The most precise result to date

BESIII results

| Mode | $\mathcal{B}(D_s^+ \rightarrow \tau^+ \nu_\tau)$ | $\mathcal{B}(D_s^+ \rightarrow \mu^+ \nu_\mu)$ |
|---|--|--|
| $\tau^+ \rightarrow \pi^+ \pi^0 \bar{\nu}_\tau$ | $(5.29 \pm 0.25 \pm 0.20)\%$ | - |
| $\tau^+ \rightarrow \pi^+ \bar{\nu}_\tau$ | $(5.21 \pm 0.25 \pm 0.17)\%$ | $(0.535 \pm 0.013 \pm 0.016)\%$ |
| $\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$ | $(5.27 \pm 0.10 \pm 0.12)\%$ | - |
| Average | $(5.26 \pm 0.09 \pm 0.09)\%$ | $(0.535 \pm 0.013 \pm 0.016)\%$ |

Combining our results with world averages


$$\mathcal{B}_{D_s^+ \rightarrow \tau^+ \nu_\tau} / \mathcal{B}_{D_s^+ \rightarrow \mu^+ \nu_\mu} = 9.72 \pm 0.37$$

SM prediction 9.75 ± 0.01

No LFU violation is found with the current precision


$$f_{D_S^+} |V_{cs}| = (244.4 \pm 2.3 \pm 2.9) \text{ MeV}$$

Input $|V_{cs}| = 0.97320 \pm 0.00011$ from CKM global fit




Most precise measurement

Input $f_{D_S^+} = 249.9 \pm 0.5$ from LQCD calculations (FLAVG19)



Semi-leptonic decay



$$\frac{d\Gamma}{dq^2} = X \frac{G_F^2 p^3}{24\pi^3} |f_+(q^2)|^2 |V_{cd(s)}|^2 \quad (X = 1 \text{ for } K^-, \pi^-, \bar{K}^0, \eta^{(\prime)}; X = \frac{1}{2} \text{ for } \pi^0)$$

- Analyze exp. partial decay rates \rightarrow q^2 dependence of $f_+^{K(\pi)}(q^2)$, extract $f_+^{K(\pi)}(0)$ with $|V_{cs(d)}|^{\text{CKMfitter}}$ as input ---- calibrate QCD
- Exp. + LQCD calculation of $f_+^{K(\pi)}(0)$ and $f_+^{\pi}(0) \rightarrow V_{cs(d)}$ ---- constrain CKM

First observation of $D^+ \rightarrow \eta \mu \nu_\mu$

$2.93 \text{fb}^{-1} @ E_{cm} = 3.773 \text{GeV}$
 $e^+ e^- \rightarrow \Psi(3770) \rightarrow D D \bar{D}$


BESIII: PRL 124, 231801 (2020)

$B[D^+ \rightarrow \eta \mu^+ \nu] = (0.104 \pm 0.010 \pm 0.005)\%$

$$R_{D\eta} = \frac{\Gamma[D^+ \rightarrow \eta \mu^+ \nu]}{\Gamma[D^+ \rightarrow \eta e^+ \nu]} = 0.91 \pm 0.13$$



(SM prediction: 0.93-0.96)

$$f_+^{D \rightarrow \eta}(0) |V_{cd}| = 0.087(08)(02)$$



No. of single tags: $(1522.5 \pm 2.1) \times 10^3$

No. of double tags: 234 ± 22







Selected topics


- Light hadrons: glueballs & more
- XYZ particles: Y(4260), X(3872), Z_{cs}(3985)
- Charm decays: CKM, decay constants, form factors, LFU
- Hadronic corrections to muon g-2: HPV & HLbL
- Baryons: form factors & polarization

Standard Model contributions to muon g-2

$$a_\mu = a_\mu(\text{QED}) + a_\mu(\text{Weak}) + a_\mu(\text{Hadronic})$$


| | | |
|---|--------------------------------------|-----------|
| <p>QED</p>  <p>+ ...</p> | $116\,584\,718.9(1) \times 10^{-11}$ | 0.001 ppm |
| <p>Weak</p>  <p>+ ...</p> | $153.6(1.0) \times 10^{-11}$ | 0.01 ppm |
| <p>Hadronic...</p> | | |
| <p>...Vacuum Polarization (HVP)</p>  <p>+ ...</p> | $6845(40) \times 10^{-11}$ [0.6%] | 0.37 ppm |
| <p>...Light-by-Light (HLbL)</p>  <p>+ ...</p> | $92(18) \times 10^{-11}$ [20%] | 0.15 ppm |

From Xu Feng





Uncertainty: dominated by **strong interactions**

BESIII contributions to HVP




$$a_{\mu}^{HVP} = \frac{1}{4\pi^3} \int_{4m_{\pi}^2}^{\infty} ds K(s) \sigma_{had}(s)$$

$\sigma_{had}(s) = \sigma_{tot}(e^+e^- \rightarrow \text{Hadrons})$
relevant mass range < 2...3 GeV



$e^+e^- \rightarrow \pi^+\pi^-\gamma_{ISR}$



| | |
|--------------------------------|--------------------------|
| CMD-2 03,06 | 372.4 ± 3.0 |
| SND 04 | 371.7 ± 5.0 |
| CLEO 18 | 376.9 ± 6.3 |
| BaBar 09 | 376.7 ± 2.7 |
| KLOE18 avg. of KLOE08/10/12 | 366.9 ± 2.1 |
| BESIII 16 | 368.2 ± 2.5 ± 3.3 |
| BESIII (Updated) | 368.2 ± 1.5 ± 3.3 |
| BESIII (Future) | 368.2 ± 0.7 ± 2.2 |


$a_{\mu}^{\pi\pi, LO}(600-900 \text{ MeV}) [10^{-10}]$

Great achievement with coming 20 fb⁻¹ @3773 MeV

From Achim

BESIII contribution to Hadronic light-by-light scattering: transition form factors


$$\gamma\gamma^* \rightarrow \pi^0, \eta, \eta', \pi^0\pi^0, \pi^+\pi^-, K\bar{K}$$



HLbL


Data-driven approach!

Exp. Input: **Transition Form Factors TFF** $F(Q^2)$
momentum transfer Q^2 below $\sim 2 \text{ GeV}^2$





Meson

Estimate of (g-2) Theory Initiative:
 $(9.2 \pm 1.8) \cdot 10^{-10}$ was $(10.5 \pm 2.6) \cdot 10^{-10}$



PPNP107 (2019) 20

Status of hadronic light-by-light contribution





Large data sample at 3.773 GeV: 20 fb⁻¹ (coming soon)
 Large data sample in high energy point: 4.0 – 5.6 GeV

Selected topics

- Light hadrons: glueballs & more
- XYZ particles: $Y(4260)$, $X(3872)$, $Z_{cs}(3985)$
- Charm decays: CKM, decay constants, form factors, LFU
- Hadronic corrections to muon $g-2$: HPV & HLbL
- Baryons: form factors & polarization

Oscillation Structure in neutron Form Factor


arXiv:2103.12486, accepted in Nature Physics




$$G_{osc}(q^2) = |G_n| - G_{D^*},$$

$$G_{D^*} = G_D \cdot \frac{1}{1 + \frac{q^2}{m_a^2}},$$

$$G_D = \frac{A_n}{\left(1 - \frac{q^2}{0.71(\text{GeV}^2)}\right)^2}$$



- a similar periodic structure of $|G_{eff}|$ as proton
- Simultaneous fit to $|G_{eff}|$ of neutron and proton yields a shared frequency $5.55 \pm 0.28 \text{ GeV}^{-1}$
- a large phase difference $\Delta b^{osc} = |b_{2p}^{osc} - b_{2n}^{osc}| = (125 \pm 12)^\circ$



$$F(p) = b_0^{osc} e^{-b_1^{osc} p} \cos(b_2^{osc} p + b_3^{osc})$$

Planned future data set

Table 7.1: List of data samples collected by BESIII/BEPCII up to 2019, and the proposed samples for the remainder of the physics program. The most right column shows the number of required data taking days in current (T_C) or upgraded (T_U) machine. The machine upgrades include top-up implementation and beam current increase.

| Energy | Physics motivations | Current data | Expected final data | T_C / T_U |
|---------------------|---|--|--|---------------|
| 1.8 - 2.0 GeV | R values Nucleon cross-sections | N/A | 0.1 fb ⁻¹ (fine scan) | 60/50 days |
| 2.0 - 3.1 GeV | R values Cross-sections | Fine scan (20 energy points) | Complete scan (additional points) | 250/180 days |
| ✓ J/ψ peak | Light hadron & Glueball J/ψ decays | 3.2 fb ⁻¹ (10 billion) | 3.2 fb ⁻¹ (10 billion) | N/A |
| ✓ $\psi(3686)$ peak | Light hadron & Glueball Charmonium decays | 0.67 fb ⁻¹ (0.45 billion) | 4.5 fb ⁻¹ (3.0 billion) | 150/90 days |
| $\psi(3770)$ peak | D^0/D^\pm decays | 2.9 fb ⁻¹ | 20.0 fb ⁻¹ | 610/360 days |
| 3.8 - 4.6 GeV | R values XYZ /Open charm | Fine scan (105 energy points) | No requirement | N/A |
| 4.180 GeV | D_s decay XYZ /Open charm | 3.2 fb ⁻¹ | 6 fb ⁻¹ | 140/50 days |
| 4.0 - 4.6 GeV | XYZ /Open charm Higher charmonia cross-sections | 16.0 fb ⁻¹ at different \sqrt{s} | 30 fb ⁻¹ at different \sqrt{s} | 770/310 days |
| 4.6 - 4.9 GeV | Charmed baryon/ XYZ cross-sections | 0.56 fb ⁻¹ at 4.6 GeV | 15 fb ⁻¹ at different \sqrt{s} | 1490/600 days |
| 4.74 GeV | $\Sigma_c^+ \Lambda_c^-$ cross-section | N/A | 1.0 fb ⁻¹ | 100/40 days |
| 4.91 GeV | $\Sigma_c \Sigma_c$ cross-section | N/A | 1.0 fb ⁻¹ | 120/50 days |
| 4.95 GeV | Ξ_c decays | N/A | 1.0 fb ⁻¹ | 130/50 days |

to be complete
in 2022-23



~55 fb⁻¹

Proposal of the upgrade BEPCII

✓ An upgrade of BEPCII (**BEPCII-U**) has been approved in July 2021:
the optimized energy is 2.35 GeV with luminosity 3 times higher than current BEPCII and extend the maximum energy to 5.6 GeV

- Add another cavity per beam to improve the RF power
- Change optics slightly, increase number of bunches
- Challenges: high beam intensities, backgrounds and aging effect in the detector
- Small risk: can continue running with better performance than BEPCII
- Timescale: 2.5 years construction + 0.5 year installation
- Installation: July – December 2024 and the upgraded machine ready in Jan. 2025

| | BEPCII | BEPCII-U |
|--|--------|----------|
| Lum [$10^{32}\text{cm}^{-2}\text{s}^{-1}$] | 3.5 | 11 |
| β_y^* [cm] | 1.5 | 1.35 |
| Bunch Current [mA] | 7.1 | 7.5 |
| Bunch Num | 56 | 120 |
| SR Power [kW] | 110 | 250 |
| $\xi_{y,\text{lum}}$ | 0.029 | 0.033 |
| Emittance [nmrad] | 147 | 152 |
| Coupling [%] | 0.53 | 0.35 |
| Bucket Height | 0.0069 | 0.011 |
| $\sigma_{z,0}$ [cm] | 1.54 | 1.07 |
| σ_z [cm] | 1.69 | 1.22 |
| RF Voltage [MV] | 1.6 | 3.3 |



Summary

- Data with unprecedented statistical accuracy from BESIII provides great opportunities to hadron physics and flavor physics. Will continue to run for ~10 years
- BESIII is in good status, inner detector upgrade in progress
- High-lumi. fine scan between 3.8 GeV and 5.6 GeV is planned
→ BEPCII-U: 3x upgrade on luminosity
- To explore the high statistics data sets, synergies between experiment and theory are essential

Thank you for your attention