

# CR scattering against pre-existing MHD turbulent modes

*Wednesday, 27 October 2021 16:40 (20 minutes)*

We present the phenomenological implications of the micro-physics of CR diffusion as resulting from particle scattering onto the three modes in which \textit{Magneto-Hydro-Dynamics} (MHD) cascades are decomposed. We calculate the diffusion coefficients from first principles based on reasonable choices of the physical quantities characterizing the different environments of our Galaxy, namely the \textit{Halo} and the \textit{Warm Ionized Medium}, and implement for the first time these coefficients in the \texttt{DRAGON2} numerical code. Remarkably, we obtain the correct propagated slope and normalization for all the charged species taken into account, without any \textit{ad-hoc} tuning of the transport coefficients. We show that fast magnetosonic modes dominate CR confinement up to  $\sim 100$  TeV; Alfvénic modes are strongly subdominant due to the anisotropy of the cascade (in agreement with previous findings) up to rigidities in the sub-PeV domain, where their contribution may show up as a spectral feature, potentially observable in the upcoming years. We also find that such framework cannot be responsible for CR confinement below  $\sim 200$  GeV, possibly leaving room for an additional confinement mechanism, and that the Kolmogorov-like scaling of the  $B/C$  ratio cannot be reproduced. Therefore this scaling might not be the imprint of the pre-existing turbulence spectrum

## Please choose the session this abstract belongs to

Cosmic rays

**Primary author:** FORNIERI, Ottavio (The Gran Sasso Science Institute)

**Presenter:** FORNIERI, Ottavio (The Gran Sasso Science Institute)

**Session Classification:** Session 3