



# Time-dependent treatment of cosmic-ray spectral steepening due to turbulence driving

Martin Pohl



# Acceleration process




Shock acceleration:

Significant energy gain  
by multiple shock crossing

Test-particle spectrum  $E^{-2}$

Requires continuous driving of turbulence





# Question



**Turbulence driving implies energy loss**

**Is there an impact on CR spectra?**



# Time dependence



**Nonresonant modes (aka Bell)**

**Early analysis in steady-state limit:**

**Spectral steepening can be significant**

**Is there enough time to reach the steady state?**

**Is there additional energy transfer to heat, etc?**



# 1st method



**Peak growth rate**

$$\gamma_{\max} \simeq \Omega_p \frac{v_{\text{sh}} N_{\text{cr}}}{2 v_A N_p} = \omega_{\text{p,p}} \frac{v_{\text{sh}} N_{\text{cr}}}{2 c N_p}$$

**Energy density transfer**

$$\dot{U} \simeq \int dk \gamma(k) \frac{B_k^2}{4\pi\epsilon} \lesssim \gamma_{\max} \frac{(\delta B)^2}{4\pi\epsilon}$$

**Insert and take ratio**

$$\tau_{\text{loss}} \simeq \frac{U_{\text{cr}}}{\dot{U}} \gtrsim \frac{2\epsilon\Gamma_{\text{cr}}}{\omega_{\text{p,p}}} \frac{U_{\text{bulk}}}{U_{\delta B}} \frac{c^3}{v_{\text{sh}}^3}$$



# 1st method



Comparison with acceleration time  
requires diffusion coefficient

$$\kappa = \eta r_L c / 3$$

Yields spectral modification

$$\Delta s \lesssim \frac{2(s-1)\eta M_A}{3\epsilon} \frac{U_{\delta B}}{U_{\text{bulk}}}$$

Here  $M_A$  is written with the full field amplitude



# 2nd method



Integrate energy loss rate over the entire precursor

$$\dot{E}_{\text{tot}} \lesssim \frac{\omega_{\text{pp,sh}} N_{\text{cr,sh}}}{8\pi \epsilon c N_{\text{p,sh}} \sqrt{v_{\text{sh}}}} \int_{r_{\text{sh}}}^{\infty} dr \frac{v(r)^{3/2} (\delta B(r))^2}{\exp\left(\int_{r_{\text{sh}}}^r dr' \frac{v(r')}{\kappa(r')}\right)}$$

Allows treatment of spatial variations in precursor

They don't matter  $\rightarrow$  Steepening


$$\lesssim \frac{2(s-1)^2 \eta M_A}{3\epsilon} \frac{U_{\delta B}}{U_{\text{bulk}}}$$

# Generalization

Replace Alfvénic Mach number for  $\delta B \gg B_0$

$$\Delta s \lesssim \frac{2(s-1)^2 \eta}{3\epsilon M_A}$$

$$= \frac{2(s-1)^2 \eta}{3\epsilon} \sqrt{\frac{U_{\delta B}}{U_{\text{bulk}}}}$$







# Generalization




Bell's instability requires  $kr_L \gg 1$  or  $\frac{v_{sh}}{4c} \frac{U_{cr}}{U_{\delta B}} \gg 1$

which gives

$$\Delta s \lesssim \frac{(s-1)\eta}{3\sqrt{2}\epsilon} \sqrt{\frac{v_{sh}}{c} \frac{U_{cr}}{U_{bulk}}}$$

These are upper limits!





# Summary



**Spectral steepening on account of turbulence driving happens**

**Its level is limited even for fast shocks,  $\Delta s < 0.1$ .**

**Nonthermal X rays indicate very weak steepening for TeV CRs**

**Important are the time and space available.**

**Earlier steady-state estimate gives less than one growth time**

