

ATLAS measurements of CP violation and rare decays processes with beauty mesons



Marcella Bona
(QMUL)
on behalf of the
ATLAS collaboration



School of Chemical and Physical Sciences
Queen Mary University of London



Queen Mary
University of London

Royal Society Leverhulme Trust
Senior Research Fellow


THE
ROYAL
SOCIETY

LEVERHULME
TRUST _____

TeV Particle Astrophysics 2021 (TeVPA 2021)
Chengdu, China and online
October 26th, 2021

B physics in ATLAS

- ATLAS has collected 25 fb^{-1} of data in Run 1, and 139 fb^{-1} in Run 2
- Has access to B, B_s , B_c , Λ_b , etc.
- Focus mostly on final states with muons
 - Typical trigger: di-muons with p_T thresholds at 4, 6 and 11 GeV
 - In 2018, a di-electron high-level trigger implemented and being analysed now
- **Rare and semi-rare decays: state of the art at ATLAS**
 - B to $K^*\mu\mu$ angular analysis in Run 1 [*JHEP 10 (2018) 047*]
 - $B_{(s)}$ to $\mu\mu$ in 2015-2016 Run 2 [*JHEP 04 (2019) 098*]
 - LHC combination $B_{(s)}$ to $\mu\mu$ for Summer 2020, partial Run 2 [*ATLAS-CONF-2020-049*]
- **CP Violation in B_s system: most recent result at ATLAS**
 - CP violating phase φ_s in $B_s^0 \rightarrow J/\psi\phi$ angular analysis in 2015-2017 Run 2 [*Eur. Phys. J. C 81 (2021) 342*]




CP violation in the SM and NP:

- $B_{(s)}$ systems are giving us a rather precise picture
- However there is some space for NP
- Could appear as new contributions in $\Delta F=2$ loop processes

$$A_q = C_{B_q} e^{2i\phi_{B_q}} A_q^{SM} e^{2i\phi_q^{SM}}$$



$$A_q = \left(1 + \frac{A_q^{NP}}{A_q^{SM}} e^{2i(\phi_q^{NP} - \phi_q^{SM})} \right) A_q^{SM} e^{2i\phi_q^{SM}}$$



[0707.0636 hep-ph]

The ratio of NP/SM amplitudes need to be:

- < 30% @95% prob. in B_d mixing
- < 18% @95% prob. in B_s mixing



Flavour Changing Neutral Currents: b to sll

There are a lot of measurements that can test b to sll transitions:


Branching ratios,
Angular analyses
SM symmetry tests

$$B_s \rightarrow \ell^+ \ell^-, B \rightarrow K \ell^+ \ell^-,$$

$$B \rightarrow K^* \ell^+ \ell^-, B_s \rightarrow \phi \ell^+ \ell^-,$$

$$\Lambda_b \rightarrow p K^- \ell^+ \ell^-, \dots$$

Suppressed: with branching ratios from 10^{-6} down
hence new physics effects can enhance their rates
Clean: varying levels of cleanliness



Weak effective theory:


four-fermion interaction with effective couplings:

$$\text{Wilson coefficients } C_i = C_i^{\text{SM}} + C_i^{\text{NP}}$$

Main SM contributions:

Vector (C_9) and Axial-vector (C_{10})
leptonic currents

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_i C_i O_i$$



Angular analysis on $B \rightarrow K^* \mu\mu$

Run1 result:

JHEP 10 (2018) 047, arXiv:1805.04000



HL-LHC prospects:

ATL-PHYS-PUB-2019-003

Angular analysis on $B \rightarrow K^* \mu \mu$

JHEP 10 (2018) 047, arXiv:1805.04000

- angular distribution of the 4 particles in the final state sensitive to new physics for the interference of NP and SM diagrams
- allows measuring a large set of angular parameters sensitive to Wilson coefficients $C_7^{(\prime)}$, $C_9^{(\prime)}$, $C_{10}^{(\prime)}$, $C_{S,P}^{(\prime)}$



- decay described by three angles (θ_L , θ_K , ϕ) and the di-muon mass squared q^2 \rightarrow the angular distribution is analysed in finite bins of q^2 as a function of θ_L , θ_K and ϕ .
- LHCb reports a 3.4σ deviation from the SM.

JHEP 02 (2016) 104
arXiv:1512.04442

Angular analysis on $B \rightarrow K^* \mu \mu$

JHEP 10 (2018) 047, arXiv:1805.04000

- B^0 flavour eigenstate can be identified through the $K^* \rightarrow K^- \pi^+$ decay
- angular distribution given by:

$$\frac{1}{d\Gamma/dq^2} \frac{d^4\Gamma}{d\cos\theta_\ell d\cos\theta_K d\phi dq^2} = \frac{9}{32\pi} \left[\frac{3(1-F_L)}{4} \sin^2\theta_K + F_L \cos^2\theta_K + \frac{1-F_L}{4} \sin^2\theta_K \cos 2\theta_\ell \right. \\ \left. - F_L \cos^2\theta_K \cos 2\theta_\ell + S_3 \sin^2\theta_K \sin^2\theta_\ell \cos 2\phi + S_4 \sin 2\theta_K \sin 2\theta_\ell \cos \phi \right. \\ \left. + S_5 \sin 2\theta_K \sin \theta_\ell \cos \phi + S_6 \sin^2\theta_K \cos \theta_\ell \right. \\ \left. + S_7 \sin 2\theta_K \sin \theta_\ell \sin \phi + S_8 \sin 2\theta_K \sin 2\theta_\ell \sin \phi \right. \\ \left. + S_9 \sin^2\theta_K \sin^2\theta_\ell \sin 2\phi \right].$$

- the S parameters are translated into the $P^{(\prime)}$ parameters via

$$P_1 = \frac{2S_3}{1-F_L} \quad P'_{i=4,5,6,8} = \frac{S_{j=4,5,7,8}}{\sqrt{F_L(1-F_L)}}$$


- the $P^{(\prime)}$ parameters are expected to have a reduced dependence on the hadronic form factors.
- ATLAS needs to fold the angular distribution via trigonometric relations to reduce the number of free parameters
- Measurement in 6 (overlapping) bins of q^2 in the range $[0.04, 6] \text{ GeV}^2$ → 4 sets of fits for three parameters (F_L , S_3 and S_j with $j=4,5,7,8$)

$K^* \mu\mu$ angular analysis

JHEP 10 (2018) 047, arXiv:1805.04000

- Data collected in 2012 at 8 TeV with 20.3 fb⁻¹ Run 1 data
- Selection of triggers with muon p_T thresholds starting at 4 GeV
- 787 events selected with $q^2 < 6 \text{ GeV}^2$
- Extended unbinned maximum likelihood fits in each of the fit variants in each q^2 bin: first fit of the invariant mass distributions and then fit of angular distributions to extract F_L and $S(P)$ parameters $\bar{a}_{\text{FB}}^{\mu\mu}$
- Results compatible with predictions
- Deviations of about 2.5σ (2.7σ) from DHMV in P'_4 (P'_5) in $[4,6] \text{ GeV}^2$

q^2 [GeV ²]	P'_4	P'_5
[0.04, 2.0]	$0.31 \pm 0.40 \pm 0.20$	$0.67 \pm 0.26 \pm 0.16$
[2.0, 4.0]	$-0.76 \pm 0.31 \pm 0.21$	$-0.33 \pm 0.31 \pm 0.13$
[4.0, 6.0]	$0.64 \pm 0.33 \pm 0.18$	$0.26 \pm 0.35 \pm 0.18$
[0.04, 4.0]	$-0.30 \pm 0.24 \pm 0.17$	$0.32 \pm 0.21 \pm 0.11$
[1.1, 6.0]	$0.05 \pm 0.22 \pm 0.14$	$0.01 \pm 0.21 \pm 0.08$
[0.04, 6.0]	$0.05 \pm 0.20 \pm 0.14$	$0.27 \pm 0.19 \pm 0.06$



OPE and LHCb data fit: CFFMPSV: Ciuchini et al.; JHEP 06 (2016) 116.
QCD factorisation: DMVH: Decotes-Genon et al.; JHEP 12 (2014) 125.
 JC: Jäger-Camalich; Phys. Rev. D93 (2016) 014028.

rare B decays $B_{(s)} \rightarrow \mu^+ \mu^-$

Run1 result:

EPJ C76 (2016) 513, arXiv:1604.04263

Run2 result on 2015-2016 data:





JHEP 04 (2019) 098, arXiv:1812.03017

LHC combination:

ATLAS-CONF-2020-049

HL-LHC prospects:


ATL-PHYS-PUB-2018-005



Motivations and Predictions

- Decays of B^0 and B_s^0 into two leptons have to proceed through Flavour Changing Neutral Currents (FCNC)
 - forbidden at tree level in the SM
- In addition, they are CKM and helicity suppressed.
- Depends on a single term in the effective Hamiltonian: C_{10}
- Within the SM, they can be calculated with small theoretical uncertainties of order 6-8%

meson type	Lepton type		
	e	μ	τ
B^0	$(2.48 \pm 0.21)10^{-15}$	$(1.06 \pm 0.09)10^{-10}$	$(2.22 \pm 0.19)10^{-8}$
B_s^0	$(8.54 \pm 0.55)10^{-14}$	$(3.65 \pm 0.23)10^{-9}$	$(7.73 \pm 0.49)10^{-7}$





PRL 112 (2014) 101801

- Latest predictions: $\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (3.66 \pm 0.14) \times 10^{-9}$
- $\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) = (1.03 \pm 0.05) \times 10^{-10}$

JHEP10(2019)232

- Perfect ground for indirect new physics searches:


- virtual new particles can contribute to the loop





rare B decays $B_{(s)} \rightarrow \mu^+ \mu^-$

Run: 302137
Event: 1093131714
2016-06-16 01:48:30 CEST



ATLAS analysis on 2015-2016 Run 2 data

- 36.2 fb⁻¹ dataset of 2015-2016 data taking:
 - Effectively 26.3 fb⁻¹ for B → μμ
- Trigger: higher thresholds [4-6 GeV] than in Run1,
 - L_{xy} > 0 request at trigger level


JHEP 04 (2019) 098
arXiv:1812.03017

$$\mathcal{B}(B_{(s)}^0 \rightarrow \mu^+ \mu^-) = \frac{N_{d(s)}}{\epsilon_{\mu^+ \mu^-}} \times \frac{\epsilon_{J/\psi K^+}}{N_{J/\psi K^+}} \times \frac{f_u}{f_{d(s)}} \times [\mathcal{B}(B^+ \rightarrow J/\psi K^+) \times \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)]$$

- correction for the different hadronisation probabilities for B_s⁰ and B⁰ vs B[±]
- include the B[±] and J/ψ branching fractions
- correction for the efficiencies of the two channels



Normalisation B yield extraction

- unbinned maximum likelihood fit of the invariant mass $m_{J/\psi K} \rightarrow m_{\mu\mu K}$



Backgrounds and control samples

- combinatorial background: μ 's from other b quarks
 - BDT classifier with 15 variables
- partially reconstructed B decays:
 - Same Vertex (SV): $B \rightarrow \mu\mu X$ decays
 - Same Side (SS): $b \rightarrow c\mu\nu \rightarrow s(d)\mu\mu\nu$
 - B_c decays: like $B_c \rightarrow J/\psi \mu\nu$
- semileptonic B and B_S decays: μ and charged hadron
- peaking background from hadronic $B_{(S)}$ decays:
 - B decays to two hadrons h (K/ π): $B_{(S)}^0 \rightarrow hh'$



Tight muon-ID against hadron misID

- negligible misidentification of protons ($< 0.01\%$)
- misidentification is 0.08% (0.10%) for K (π).

peaking-background events: 2.7 ± 1.3

Efficiency ratio $\epsilon_{\mu\mu}/\epsilon_{J/\psi K}$

- from MC and systematic from data-MC discrepancies
- For B_S^0 : 2.7% correction for lifetime difference of the B_S^0 mass eigenstates


Source	Contribution (%)
Statistical	0.8
BDT Input Variables	3.2
Kaon Tracking Efficiency	1.5
Muon trigger and reconstruction	1.0
Kinematic Reweighting (DDW)	0.8
Pile-up Reweighting	0.6

JHEP 04 (2019) 098, arXiv:1812.03017

Signal yield extraction

- unbinned maximum likelihood fit to the dimuon mass simultaneously in 4 BDT bins
 - 18% signal efficiency each bin
 - signals, B to hh: 3 double Gaussians
 - continuum: first order polynomial
 - partially reconstructed B: exponential
 - semi-leptonic: exponential

JHEP 04 (2019) 098, arXiv:1812.03017




Run 2 results and combinations with Run 1

- yields unconstrained:
 - $N_S = 80 \pm 22$ and $N_d = -12 \pm 20$
 - expected from the SM:
 - $N_S = 91 \pm$ and $N_d = 10$

Neyman Contours for Run 2:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (3.21^{+0.96+0.49}_{-0.91-0.30}) \times 10^{-9} = (3.2^{+1.1}_{-1.0}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 4.3 \times 10^{-10} \text{ @ 95\% CL}$$



Run 1 + Run 2 (2015+2016):
Compatible with SM at 2.4σ

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.8^{+0.8}_{-0.7}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 2.1 \times 10^{-10}$$

LHC combination from Summer 2020



ATLAS-CONF-2020-049

- Combination from binned two-dimensional profile likelihoods
- Independent systematics, except for ratio of fragmentation fractions f_d/f_s ,
 - f_d/f_s profiled separately and its uncertainty included in one likelihood.

Latest LHCb result not included

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.69^{+0.37}_{-0.35}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 1.9 \times 10^{-10} \text{ at 95\% CL.}$$



CP violation parameters from time-dependent angular analysis on $B_s \rightarrow J/\psi\phi$

Run1 result:


JHEP 08 (2016) 147, arXiv:1601.03297

Run2 result with 2015-2017 data:

Eur. Phys. J. C 81 (2021) 342, arXiv:2001.07115

HL-LHC prospects:

ATL-PHYS-PUB-2018-041




Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

- Golden mode: penguin diagrams can contribute to the decay either with the same weak phase (λ^2) or they are CKM suppressed (λ^4)

- Parameters of the B_s system:

- Mixing \rightarrow Decay width difference $\Delta\Gamma_s = \Gamma_L - \Gamma_S$

$$\Delta\Gamma_s = 0.087 \pm 0.021 \text{ ps}^{-1} \text{ in the SM [arXiv:1102.4274]}$$



- CPV phase $\varphi_s \rightarrow$ weak phase between mixing and $b \rightarrow ccs$ decay

$$\varphi_s = -2\beta_s \text{ with } \beta_s = \arg[-(V_{ts}V_{tb}^*)/(V_{cs}V_{cb}^*)]$$

$$\text{SM: } -2\beta_s = -0.0363 \pm 0.0016 \text{ [arXiv:1106.4041]}, 0.0370 \pm 0.0010 \text{ [UTfit18]}$$

- Pseudoscalar B0s to the vector–vector $J/\psi(\mu^+\mu^-)\phi(K^+K^-)$ final state
 \rightarrow admixture of CP-odd and CP-even states ($L = 0, 1$ or 2).


- $L = 0$ or $2 \rightarrow$ CP-even states, while $L = 1 \rightarrow$ CP-odd state.
- Same final state can also be K^+K^- pairs in S-wave \rightarrow CP-odd.
- CP states are separated statistically using an angular analysis

Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

- Differential decay rate:

$$\frac{d^4\Gamma}{dt d\Omega} = \sum_{k=1}^{10} O^{(k)}(t) g^{(k)}(\theta_T, \psi_T, \phi_T),$$

with $O^{(k)}(t)$ time-dependent functions corresponding to the contributions of amplitudes (A_0 , A_{\parallel} , A_{\perp} , and A_S) (and interferences) and $g^{(k)}(\theta_T, \psi_T, \phi_T)$ are angular functions.



- Flavour tagging is used to distinguish between the initial B_s^0 and \bar{B}_s^0 states.

ATLAS Run-2 result



- 80.5 fb^{-1} of 13 TeV data (Run 2, 2015-2017)
- J/ψ trigger with muon p_T of 4 or 6 GeV
- Measurement of the proper decay time $t = L_{xy} m_B / p_T^B$
- Combination with Run-1 result

Eur. Phys. J. C 81 (2021) 342,
arXiv:2001.07115

ATLAS $B_s \rightarrow J/\psi\phi$ analysis: flavour tagging

- Flavour tagging to identify the flavour of the b quark:
 - opposite-side tagging (OST) using p_T -weighted charge of tracks in cone around muons / electrons / b jets
 - Calibrated on self-tagged $B^\pm \rightarrow J/\psi K^\pm$ events
 - Tag probabilities included in the B_s fit
 - Dilution $D(Q_x)$ and tagging power T_x defined as:

$$Q_x = \frac{\sum_i^{N \text{ tracks}} q_i \cdot (p_{Ti})^\kappa}{\sum_i^{N \text{ tracks}} (p_{Ti})^\kappa}$$



$$D(Q_x) = 2P(B|Q_x) - 1$$


$$T_x = \sum_i \epsilon_{xi} \cdot (2P(B|Q_{xi}) - 1)^2$$

Tag method	ϵ_x [%]	D_x [%]	T_x [%]
Tight muon	4.50 ± 0.01	43.8 ± 0.2	0.862 ± 0.009
Electron	1.57 ± 0.01	41.8 ± 0.2	0.274 ± 0.004
Low- p_T muon	3.12 ± 0.01	29.9 ± 0.2	0.278 ± 0.006
Jet	12.04 ± 0.02	16.6 ± 0.1	0.334 ± 0.006
Total	21.23 ± 0.03	28.7 ± 0.1	1.75 ± 0.01

ATLAS $B_s \rightarrow J/\psi\phi$ analysis: ML fit

Eur. Phys. J. C 81 (2021) 342, arXiv:2001.07115

- Unbinned maximum-likelihood fit
- B_s properties: mass m_B (and its error), proper decay time t , proper decay time error σ_t , tagging probability $P(B|Q_x)$
- Transversity angles: $\Omega(\theta_T, \psi_T, \phi_T)$
- Physical parameters: $\Delta\Gamma_s, \varphi_s, \Gamma_s, |A_0(0)|^2, |A_{||}(0)|^2, \delta_{||}, \delta_{\perp}, |A_s(0)|^2$ and δ_s
- Systematics:
 - Lifetime model: varying p_T bins and signal fraction
 - Backgrounds: $B_d / \Lambda_b /$ angular models varied / p_T bins varied
 - Tagging: variation of the parameterisation / recalibration from MC samples / pile-up effects



ATLAS $B_s \rightarrow J/\psi\phi$ analysis: Run-2 results

ATLAS Run-2 result on 80.5 fb^{-1} of 2015-2017 data



Run 2 only (80.5 fb^{-1}):

Parameter	Value	Statistical uncertainty	Systematic uncertainty
ϕ_s [rad]	-0.081	0.041	0.022
$\Delta\Gamma_s$ [ps^{-1}]	0.0607	0.0047	0.0043
Γ_s [ps^{-1}]	0.6687	0.0015	0.0022
$ A_{\parallel}(0) ^2$	0.2213	0.0019	0.0023
$ A_0(0) ^2$	0.5131	0.0013	0.0038
$ A_S(0) ^2$	0.0321	0.0033	0.0046
$\delta_{\perp} - \delta_S$ [rad]	-0.25	0.05	0.04
Solution (a)			
δ_{\perp} [rad]	3.12	0.11	0.06
δ_{\parallel} [rad]	3.35	0.05	0.09
Solution (b)			
δ_{\perp} [rad]	2.91	0.11	0.06
δ_{\parallel} [rad]	2.94	0.05	0.09

Two solutions in $\delta_{\parallel} - \delta_{\perp}$ plane,
negligible impact on other parameters

Eur. Phys. J. C 81 (2021) 342, arXiv:2001.07115

Run 1 (19.2 fb^{-1}) & Run 2 (80.5 fb^{-1}):




ATLAS $B_s \rightarrow J/\psi\phi$ analysis: Run1+2 combination

Eur. Phys. J. C 81 (2021) 342,
arXiv:2001.07115

ATLAS Run 1 & Run 2 combined
(19.2 fb⁻¹ + 80.5 fb⁻¹)

Parameter	Value	Solution (a)	
		Statistical uncertainty	Systematic uncertainty
ϕ_s [rad]	-0.087	0.036	0.021
$\Delta\Gamma_s$ [ps ⁻¹]	0.0657	0.0043	0.0037
Γ_s [ps ⁻¹]	0.6703	0.0014	0.0018
$ A_{\parallel}(0) ^2$	0.2220	0.0017	0.0021
$ A_0(0) ^2$	0.5152	0.0012	0.0034
$ A_S ^2$	0.0343	0.0031	0.0045
δ_{\perp} [rad]	3.22	0.10	0.05
δ_{\parallel} [rad]	3.36	0.05	0.09
$\delta_{\perp} - \delta_S$ [rad]	-0.24	0.05	0.04

Comparison with CMS & LHCb:



$$\phi_s = -0.087 \pm 0.036 \text{ (stat)} \pm 0.021 \text{ (syst) rad}$$



$$\Delta\Gamma_s = 0.0657 \pm 0.0043 \text{ (stat)} \pm 0.0037 \text{ (syst) ps}^{-1}$$

- ϕ_s result consistent with results from CMS, LHCb and SM
- Competitive single measurement of $\Delta\Gamma_s$, Γ_s and helicity parameters
- Still to add 60 fb⁻¹ of 2018 data

$B_s \rightarrow J/\psi\phi$ results: HFLAV average

HFLAV average for PDG21:
 $\phi_s = -0.050 \pm 0.019$ rad

Because of tensions, errors on Γ_s
 and $\Delta\Gamma_s$ scaled by 2.5 and 1.77



Summary and Conclusions

- **Results on FCNC b to s transitions:**
 - B to $K^*\mu\mu$ angular analysis and $B_{(s)}$ to $\mu\mu$
- **Recent results on CP Violation in B_s system:**
 - CP violating phase in $B_s^0 \rightarrow J/\psi\phi$ angular analysis
- **A number of Run 2 analyses ongoing**
 - Updates and new analyses
- ATLAS is competitive in B physics
 - Thanks to accumulated statistical samples
 - Thanks to some detector performance (tracking)
 - Perfect example the angular analysis of the golden mode $B_s \rightarrow J/\psi\phi$
 - **Working on the updates of all the above to full Run-2 statistics and preparing for Run 3**







back-up slides

Analysis strategy for $B \rightarrow K^* \mu \mu$

- Data collected in 2012 at 8 TeV with 20.3 fb⁻¹ Run 1 data
- Measured in 6 (overlapping) bins of q^2 in the range [0.04, 6] GeV²
- 4 sets of fits for three parameters (F_L , S_3 and S_j with $j=4,5,7,8$)
- Selection of triggers with muon p_T thresholds starting at 4 GeV
- K^* tagged by the kaon sign:
 - dilution from mistag probability included in $(1-2\langle w \rangle)$:
 - $\langle w \rangle \sim 10.9(1)\%$ with small dependence on q^2
- 787 events selected with $q^2 < 6$ GeV²
- Extended unbinned maximum likelihood fits in each of the fit variants in each q^2 bin:
 - two step fit procedure: first fit the invariant mass distribution
 - then add to the fit the angular distributions to extract the F_L and $S(P)$ parameters
- Signal shape studies from control samples $K^* J/\psi$ and $K^* \psi(2S)$

Fit projections




- fit $m(K^*\mu\mu)$, $\cos\theta_L$, $\cos\theta_K$ and ϕ to isolate signal and extract parameters of interest.




- Data shown for $[0.04, 2.0] \text{ GeV}^2$
- projections for the S_5 fit.
- Approx 106-128 signal events in $2 \text{ GeV}^2 q^2$ bin.
- Similar results for the other q^2 bins and other fit variants.

Angular analysis results

- Results are compatible with theoretical calculations & fits:




$q^2 \text{ [GeV}^2\text{]}$	P_1	P'_4	P'_5
[0.04, 2.0]	$-0.05 \pm 0.30 \pm 0.08$	$0.31 \pm 0.40 \pm 0.20$	$0.67 \pm 0.26 \pm 0.16$
[2.0, 4.0]	$-0.78 \pm 0.51 \pm 0.34$	$-0.76 \pm 0.31 \pm 0.21$	$-0.33 \pm 0.31 \pm 0.13$
[4.0, 6.0]	$0.14 \pm 0.43 \pm 0.26$	$0.64 \pm 0.33 \pm 0.18$	$0.26 \pm 0.35 \pm 0.18$
[0.04, 4.0]	$-0.22 \pm 0.26 \pm 0.16$	$-0.30 \pm 0.24 \pm 0.17$	$0.32 \pm 0.21 \pm 0.11$
[1.1, 6.0]	$-0.17 \pm 0.31 \pm 0.13$	$0.05 \pm 0.22 \pm 0.14$	$0.01 \pm 0.21 \pm 0.08$
[0.04, 6.0]	$-0.15 \pm 0.23 \pm 0.10$	$0.05 \pm 0.20 \pm 0.14$	$0.27 \pm 0.19 \pm 0.06$



OPE and LHCb data fit: CFFMPSV: Ciuchini et al.; JHEP 06 (2016) 116.
 QCD factorisation: DMVH: Decotes-Genon et al.; JHEP 12 (2014) 125.
 JC: Jäger-Camalich; Phys. Rev. D93 (2016) 014028.

Angular analysis results

- ATLAS gets deviations of about 2.5σ (2.7σ) from DHMV in P'_4 (P'_5) in $[4,6] \text{ GeV}^2$




CFFMPSV: Ciuchini et al.; JHEP 06 (2016) 116.
 DHMV: Decotes-Genon et al.; JHEP 12 (2014) 125.
 JC: Jäger-Camalich; Phys. Rev. D93 (2016) 014028.

Projections for $K^* \mu\mu$ angular analysis at HL-LHC

- Extrapolation from signal/background yields in Run 1 and toy-MC simulations
- Accounting for improved performance of the ATLAS Upgraded tracking system
- Three trigger scenarios: high-yield, intermediate and low-statistics for signal.
- The precision on, for example, the P'_5 parameter expected to improve by factors of $\sim 9\times$, $\sim 8\times$, $\sim 5\times$ (for the three trigger scenarios) relative to Run 1



LHC phase	q^2 [GeV 2]	$\delta_{F_L}^{\text{tot}}$	$\delta_{P'_1}^{\text{tot}}$	$\delta_{P'_4}^{\text{tot}}$	$\delta_{P'_5}^{\text{tot}}$	$\delta_{P'_6}^{\text{tot}}$	$\delta_{P'_8}^{\text{tot}}$
Run 1	[0.04, 2.0]	0.11	0.31	0.45	0.31	0.21	0.51
	[2.0, 4.0]	0.12	0.61	0.37	0.34	0.34	0.57
	[4.0, 6.0]	0.18	0.50	0.38	0.39	0.30	0.43
HL-LHC $\mu 6\mu 6$	[0.04, 2.0]	0.010	0.027	0.037	0.037	0.019	0.046
	[2.0, 4.0]	0.008	0.093	0.040	0.038	0.040	0.070
	[4.0, 6.0]	0.016	0.083	0.032	0.047	0.033	0.041
HL-LHC $\mu 10\mu 6$	[0.04, 2.0]	0.011	0.037	0.046	0.040	0.023	0.055
	[2.0, 4.0]	0.011	0.103	0.047	0.042	0.044	0.075
	[4.0, 6.0]	0.018	0.100	0.040	0.053	0.038	0.052
HL-LHC $\mu 10\mu 10$	[0.04, 2.0]	0.018	0.065	0.076	0.059	0.041	0.093
	[2.0, 4.0]	0.017	0.15	0.074	0.068	0.059	0.100
	[4.0, 6.0]	0.026	0.17	0.074	0.082	0.063	0.090



ATL-PHYS-PUB-2019-003

Signal yield extraction




- signal yields extracted with a unbinned maximum likelihood fit to the dimuon mass
- fit performed simultaneously in four BDT bins
 - 18% signal efficiency



- signals, B to hh: 3 double Gaussians
- continuum: first order polynomial
- partially reconstructed B: exponential
- semi-leptonic: exponential

Signal yield extraction


- yields unconstrained: $N_S = 80 \pm 22$ and $N_d = -12 \pm 20$
- *expected from the SM: $N_S = 91 \pm$ and $N_d = 10$*



- consistent with Standard Model predictions
- likelihood maximum:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = \left(3.21_{-0.83-0.31}^{+0.90+0.48} \right) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) = \left(-1.3_{-1.9-0.8}^{+2.2+0.7} \right) \times 10^{-10}$$





Combination of Run 1 and Run 2 results

Neyman Contours yield for Run 2:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (3.21_{-0.91-0.30}^{+0.96+0.49}) \times 10^{-9} = (3.2_{-1.0}^{+1.1}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 4.3 \times 10^{-10} \text{ @ 95\% CL}$$




Run 1 + Run 2 (2015+2016) combination:
Compatible with SM at 2.4σ

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.8_{-0.7}^{+0.8}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 2.1 \times 10^{-10}$$

Neutral B mesons into two muons

We have been chasing this decay for a long time:



LHC combination from Summer 2020

- Combination from binned two-dimensional profile likelihoods
- Independent systematics, except for ratio of fragmentation fractions f_d/f_s ,
 - f_d/f_s profiled separately and its uncertainty included in one likelihood.

Latest LHCb result not included



$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.69^{+0.37}_{-0.35}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 1.9 \times 10^{-10} \text{ at 95\% CL.}$$

Prospect on $B_{(s)} \rightarrow \mu^+ \mu^-$ at ATLAS

- HL-LHC \rightarrow 3 trigger scenarios: with thresholds ($p_{T}^{\mu 1}, p_{T}^{\mu 2}$):
 - Conservative: (10 GeV, 10 GeV) \rightarrow $\times 15$ Run 1
 - Intermediate: (6 GeV, 10 GeV) \rightarrow $\times 60$ Run 1
 - High-yield: (6 GeV, 6 GeV) \rightarrow $\times 75$ Run 1

	$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)$		$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-)$	
	stat [10^{-10}]	stat + syst [10^{-10}]	stat [10^{-10}]	stat + syst [10^{-10}]
Run 2	7.0	8.3	1.42	1.43
HL-LHC: Conservative	3.2	5.5	0.53	0.54
HL-LHC: Intermediate	1.9	4.7	0.30	0.31
HL-LHC: High-yield	1.8	4.6	0.27	0.28



ATL-PHYS-PUB-2018-005

LHC combination from Summer 2020

ATLAS-CONF-2020-049



- Combination from binned two-dimensional profile likelihoods
- Independent systematics, except for ratio of fragmentation fractions f_d/f_s ,
 - f_d/f_s profiled separately and its uncertainty included in one likelihood.

Latest LHCb result not included

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.69^{+0.37}_{-0.35}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 1.9 \times 10^{-10} \text{ at 95\% CL.}$$

2.1 σ from SM



Around and outside B physics...

● Searches for Lepton Flavour Violating decay tau to 3 muons

- *Eur. Phys. J. C* (2016) 76:232, *arXiv:1601.03567*
 - Tau production from W decays in Run 1
 - Analysis on Run 2 ongoing: more abundant production mechanism

● Test of Lepton Flavour Universality as $BR(W \rightarrow \tau\nu_\mu)/BR(W \rightarrow \mu\nu_\mu)$

- *Nat. Phys.* (2021), *arXiv:2007.14040*
- Whole Run 2: di-leptonic tt events with either one electron and one muon (e-μ channel) or two muons (μ-μ channel).



$$R(\tau/\mu) = 0.992 \pm 0.013 [\pm 0.007 \text{ (stat)} \pm 0.011 \text{ (syst)}]$$

● For the two light generations:

$BR(W \rightarrow \mu\nu_\mu)/BR(W \rightarrow e\nu_e)$

- *Eur. Phys. J. C* 77 (2017) 367, *arXiv:1612.03016*
- From inclusive W^\pm and Z/γ^* production cross section measurements in Run 1
- Analysis on Run 2 ongoing on top decays:

$$R_{\text{top}} = \frac{B(t \rightarrow b\mu^+\nu_\mu)}{B(t \rightarrow be^+\nu_e)}$$




Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

ATLAS Run-1 result

- 4.9 fb⁻¹ of 7 TeV data (Run 1, 2011)
and 14.3 fb⁻¹ of 8 TeV data (Run 1, 2012)

$$\begin{aligned}\phi_s &= -0.090 \pm 0.078 \text{ (stat.)} \pm 0.041 \text{ (syst.) rad} \\ \Delta\Gamma_s &= 0.085 \pm 0.011 \text{ (stat.)} \pm 0.007 \text{ (syst.) ps}^{-1} \\ \Gamma_s &= 0.675 \pm 0.003 \text{ (stat.)} \pm 0.003 \text{ (syst.) ps}^{-1}\end{aligned}$$



ATLAS Run-2 result

- 80.5 fb⁻¹ of 13 TeV data (Run 2, 2015-2017)
- J/ψ trigger with muon p_T of 4 or 6 GeV
- Measurement of the proper decay time $t = L_{xy} m_B / p_T^B$
- Flavour tagging to identify the flavour of the b quark
- Combination with Run-1 result


Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

- Systematics:
 - *Lifetime model*: varying p_T bins and signal fraction
 - *Backgrounds*: $B_d / \Lambda_b /$ angular models varied / p_T bins varied
 - *Tagging*: variation of the parameterisation / recalibration from MC samples / pile-up effects

	ϕ_s [10^{-3} rad]	$\Delta\Gamma_s$ [10^{-3} ps $^{-1}$]	Γ_s [10^{-3} ps $^{-1}$]	$ A_{\parallel}(0) ^2$ [10^{-3}]	$ A_0(0) ^2$ [10^{-3}]	$ A_S(0) ^2$ [10^{-3}]	δ_{\perp} [10^{-3} rad]	δ_{\parallel} [10^{-3} rad]	$\delta_{\perp} - \delta_S$ [10^{-3} rad]
Tagging	19	0.4	0.3	0.2	0.2	1.1	17	19	2.3
ID alignment	0.8	0.2	0.5	< 0.1	< 0.1	< 0.1	11	7.2	< 0.1
Acceptance	0.5	0.3	< 0.1	1.0	0.9	2.9	37	64	8.6
Time efficiency	0.2	0.2	0.5	< 0.1	< 0.1	0.1	3.0	5.7	0.5
Best candidate selection	0.4	1.6	1.3	0.1	1.0	0.5	2.3	7.0	7.4
Background angles model:									
Choice of fit function	2.5	< 0.1	0.3	1.1	< 0.1	0.6	12	0.9	1.1
Choice of p_T bins	1.3	0.5	< 0.1	0.4	0.5	1.2	1.5	7.2	1.0
Choice of mass window	9.3	3.3	0.2	0.4	0.8	0.9	17	8.6	6.0
Choice of sidebands intervals	0.4	0.1	0.1	0.3	0.3	1.3	4.4	7.4	2.3
Dedicated backgrounds:									
B_d^0	2.6	1.1	< 0.1	0.2	3.1	1.5	10	23	2.1
Λ_b	1.6	0.3	0.2	0.5	1.2	1.8	14	30	0.8
Alternate Δm_s	1.0	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1	15	4.0	< 0.1
Fit model:									
Time res. sig frac	1.4	1.1	0.5	0.5	0.6	0.8	12	30	0.4
Time res. p_T bins	0.7	0.5	0.8	0.1	0.1	0.1	2.2	14	0.7
S-wave phase	0.3	< 0.1	< 0.1	< 0.1	< 0.1	0.2	8.0	15	37
Fit bias	5.7	1.3	1.2	1.3	0.4	1.1	3.3	19	0.3
Total	22	4.3	2.2	2.3	3.8	4.6	55	88	39

Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

$$\frac{d^4\Gamma}{dt d\Omega} = \sum_{k=1}^{10} \mathcal{O}^{(k)}(t) g^{(k)}(\theta_T, \psi_T, \phi_T),$$



k	$\mathcal{O}^{(k)}(t)$	$g^{(k)}(\theta_T, \psi_T, \phi_T)$
1	$\frac{1}{2} A_0(0) ^2 \left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$2 \cos^2 \psi_T (1 - \sin^2 \theta_T \cos^2 \phi_T)$
2	$\frac{1}{2} A_{ }(0) ^2 \left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\sin^2 \psi_T (1 - \sin^2 \theta_T \sin^2 \phi_T)$
3	$\frac{1}{2} A_{\perp}(0) ^2 \left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\sin^2 \psi_T \sin^2 \theta_T$
4	$\frac{1}{2} A_0(0) A_{ }(0) \cos \delta_{ }$ $\left[(1 + \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 - \cos \phi_s) e^{-\Gamma_H^{(s)} t} \pm 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{1}{\sqrt{2}} \sin 2\psi_T \sin^2 \theta_T \sin 2\phi_T$
5	$ A_{ }(0) A_{\perp}(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \cos(\delta_{\perp} - \delta_{ }) \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\sin(\delta_{\perp} - \delta_{ }) \cos(\Delta m_s t) - \cos(\delta_{\perp} - \delta_{ }) \cos \phi_s \sin(\Delta m_s t)) \right]$	$-\sin^2 \psi_T \sin 2\theta_T \sin \phi_T$
6	$ A_0(0) A_{\perp}(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \cos \delta_{\perp} \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\sin \delta_{\perp} \cos(\Delta m_s t) - \cos \delta_{\perp} \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{1}{\sqrt{2}} \sin 2\psi_T \sin 2\theta_T \cos \phi_T$
7	$\frac{1}{2} A_S(0) ^2 \left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{2}{3} (1 - \sin^2 \theta_T \cos^2 \phi_T)$
8	$\alpha A_S(0) A_{ }(0) \left[\frac{1}{2}(e^{-\Gamma_L^{(s)} t} - e^{-\Gamma_H^{(s)} t}) \sin(\delta_{ } - \delta_S) \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\cos(\delta_{ } - \delta_S) \cos(\Delta m_s t) - \sin(\delta_{ } - \delta_S) \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{1}{3} \sqrt{6} \sin \psi_T \sin^2 \theta_T \sin 2\phi_T$
9	$\frac{1}{2}\alpha A_S(0) A_{\perp}(0) \sin(\delta_{\perp} - \delta_S)$ $\left[(1 - \cos \phi_s) e^{-\Gamma_L^{(s)} t} + (1 + \cos \phi_s) e^{-\Gamma_H^{(s)} t} \mp 2e^{-\Gamma_s t} \sin(\Delta m_s t) \sin \phi_s \right]$	$\frac{1}{3} \sqrt{6} \sin \psi_T \sin 2\theta_T \cos \phi_T$
10	$\alpha A_0(0) A_S(0) \left[\frac{1}{2}(e^{-\Gamma_H^{(s)} t} - e^{-\Gamma_L^{(s)} t}) \sin \delta_S \sin \phi_s \right.$ $\left. \pm e^{-\Gamma_s t} (\cos \delta_S \cos(\Delta m_s t) + \sin \delta_S \cos \phi_s \sin(\Delta m_s t)) \right]$	$\frac{4}{3} \sqrt{3} \cos \psi_T (1 - \sin^2 \theta_T \cos^2 \phi_T)$





TD angular analysis of $B_s \rightarrow J/\psi\phi$

ATLAS Run-1 result:

- 14.3 fb⁻¹ of ATLAS data from 2012 at 8 TeV
 - Results:
 - $\phi_s = -0.090 \pm 0.078$ (stat) ± 0.041 (syst) rad
 - $\Delta\Gamma_s = 0.085 \pm 0.011$ (stat) ± 0.007 (syst) ps⁻¹
- [JHEP 08 (2016) 147]*



- Agrees with SM
- Consistent with other experiments
- Consistent with previous analysis, using 2011 data at 7 TeV
 - [Phys. Rev. D 90, 052007 (2014)]*
- A Best Linear Unbiased Estimate (BLUE) combination used to combine 7 and 8 TeV measurements



$B_s \rightarrow J/\psi\phi$ results: HFLAV average


HFLAV average for PDG21:
 $\phi_s = -0.050 \pm 0.019$ rad

Because of tensions, errors on Γ_s and $\Delta\Gamma_s$ scaled by 2.5 and 1.77



ATLAS $B_s \rightarrow J/\psi\phi$ results: HL-LHC projections

- Updated tracking (ITk): proper decay time resolution improved by 21% w.r.t. Run 2
- Three trigger scenarios for thresholds
- Improvements w.r.t. Run 1:
 - ϕ_s stat: better by $\sim 9x$ to $20x$
 - uncertainty on ϕ_s at least as the theory error
 - $\Delta\Gamma_s$ stat: better by $\sim 4x$ to $10x$





ATL-PHYS-PUB-2018-041

ϕ_s^{ccs} [rad]


Time-dependent angular analysis of $B_s \rightarrow J/\psi\phi$

ATLAS combined Run-1 result:



Motivations and predictions



- Decays of B^0 and B_s^0 into two leptons have to proceed through Flavour Changing Neutral Currents (FCNC)
 - forbidden at tree level in the SM
- In addition, they are CKM and helicity suppressed.
- Within the SM, they can be calculated with small theoretical uncertainties of order 6-8%



meson type	Lepton type		
	e	μ	τ
B^0	$(2.48 \pm 0.21)10^{-15}$	$(1.06 \pm 0.09)10^{-10}$	$(2.22 \pm 0.19)10^{-8}$
B_s^0	$(8.54 \pm 0.55)10^{-14}$	$(3.65 \pm 0.23)10^{-9}$	$(7.73 \pm 0.49)10^{-7}$

Bobeth et al., PRL 112 (2104) 101801 [includes NLO EM and NNLO QCD corrections]

- Perfect ground for indirect new physics searches:
 - virtual new particles can contribute to the loop
 - both enhancement and suppression effects are possible



ATLAS analysis on 2015-2016 Run 2 data

JHEP 04 (2019) 098, arXiv:1812.03017

- 36.2/fb dataset of 2015-2016 data taking:
 - effectively 26.3/fb for $B \rightarrow \mu\mu$
 - 15.1/fb for $B \rightarrow J/\psi\Phi$ and $B \rightarrow J/\psi K$
- Trigger: higher thresholds [4-6 GeV] than in Run1,
 - $L_{xy} > 0$ request at trigger level

$$\mathcal{B}(B_{(s)}^0 \rightarrow \mu^+ \mu^-) = \frac{N_{d(s)}}{\epsilon_{\mu^+ \mu^-}} \times \frac{\epsilon_{J/\psi K^+}}{N_{J/\psi K^+}} \times \frac{f_u}{f_{d(s)}} \times [\mathcal{B}(B^+ \rightarrow J/\psi K^+) \times \mathcal{B}(J/\psi \rightarrow \mu^+ \mu^-)]$$



- correction for the different hadronisation probabilities for B_s^0 and B^0 vs B^\pm
- include the B^\pm and J/ψ branching fractions
- correction for the efficiencies of the two channels
- normalisation yield and efficiency ratio define the factor:

$$\mathcal{D}_{\text{norm}} = N_{J/\psi K^+} \left(\frac{\epsilon_{\mu^+ \mu^-}}{\epsilon_{J/\psi K^+}} \right)$$

Background contributions

In order of relative magnitude:


- combinatorial background:
 - two real muons from different b quarks
- partially reconstructed B decays:
 - two real muons
 - Same Vertex (SV): $B \rightarrow mmX$ decays
 - Same Side (SS): semileptonic decay cascades ($b \rightarrow cmn \rightarrow s(d)mmnn$)
 - B_c decays: like $B_c \rightarrow J/\psi mn$
 - all these accumulate at low values of the dimuon invariant mass
- semileptonic B and B_s decays:
 - one real muon and a charged hadron.
- peaking background from charmless hadronic $B_{(s)}$ decays:
 - B decays into two hadrons h (kaons and pions): $B^0_{(s)} \rightarrow hh'$
 - smaller component, but overlays with the signal in dimuon invariant mass



Tight muon-ID against hadron misidentification


- mis-identification reduced by 0.39^2 using standard 'tight' ATLAS selections
- studied on simulated samples
- validated on control regions
- negligible misidentification of protons ($< 0.01\%$)
- misidentification is 0.08% (0.10%) for K(p).

peaking-background events: 2.7 ± 1.3



BDT against combinatorial bkg

- MVA classifier to discriminate from signal
- trained and tested on mass sidebands
 - divided in 3 subsets
 - 3 independent BDTs
 - compatible performance
- 15 variables related to properties of B candidates, muons from the B decay, other tracks from the same collision and to pile-up vertices.



Normalisation B yield extraction



- unbinned maximum likelihood fit of the invariant mass $m_{J/\psi K} \rightarrow m_{mmK}$
- cross-checked with raw relative yield of J/ψ over J/ψK ratio
 $r_{p/K} = (3.71 \pm 0.09)\%$

$$D_{\text{norm}} = N_{J/\psi K^+} \left(\frac{\epsilon_{\mu^+ \mu^-}}{\epsilon_{J/\psi K^+}} \right)$$

Efficiency ratio $\epsilon_{\mu\mu}/\epsilon_{J/\psi K}$





- efficiency ratio from MC
- systematic from data-MC discrepancies
- For B_s^0 : 2.7% correction for lifetime difference of the B_s^0 mass eigenstates

Source	Contribution (%)
Statistical	0.8
BDT Input Variables	3.2
Kaon Tracking Efficiency	1.5
Muon trigger and reconstruction	1.0
Kinematic Reweighting (DDW)	0.8
Pile-up Reweighting	0.6




Signal yield extraction

- signal yields extracted with a unbinned maximum likelihood fit to the dimuon mass
- fit performed simultaneously in four BDT bins
- 18% signal efficiency








- signals, B to hh: 3 double Gaussians
- continuum: first order polynomial
- partially reconstructed B: exponential
- semi-leptonic: exponential



Signal yield extraction

- yields unconstrained:
 - $N_S = 80 \pm 22$ and $N_D = -12 \pm 20$
- expected from the SM:
 - $N_S = 91 \pm$ and $N_D = 10$



- consistent with Standard Model predictions
- likelihood maximum:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = \left(3.21_{-0.83-0.31}^{+0.90+0.48} \right) \times 10^{-9}$$



$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) = \left(-1.3_{-1.9-0.8}^{+2.2+0.7} \right) \times 10^{-10}$$

Combination of Run 1 and Run 2 results

Neyman Contours yield for Run 2:

$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (3.21^{+0.96+0.49}_{-0.91-0.30}) \times 10^{-9} = (3.2^{+1.1}_{-1.0}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 4.3 \times 10^{-10} \text{ @ 95\% CL}$$



Run 1 + Run 2 (2015+2016) combination:
 Compatible with SM at 2.4σ



$$\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-) = (2.8^{+0.8}_{-0.7}) \times 10^{-9}$$

$$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-) < 2.1 \times 10^{-10}$$

Angular analysis on $B \rightarrow K^* \mu \mu$

JHEP 10 (2018) 047, arXiv:1805.04000

- FCNC b to s transition with a BR $\sim 1.1 \cdot 10^{-6}$
- Angular distribution of the 4 particles in the final state sensitive to new physics for the interference of NP and SM diagrams



- Decay described by three angles (q_L, q_K, ϕ) and the di-muon mass squared $q^2 \rightarrow$ angular distribution in bins of q^2 as function of q_L, q_K and ϕ .

$$\frac{1}{d\Gamma/dq^2} \frac{d^4\Gamma}{d \cos \theta_\ell d \cos \theta_K d\phi dq^2} = \frac{9}{32\pi} \left[\frac{3(1-F_L)}{4} \sin^2 \theta_K + F_L \cos^2 \theta_K + \frac{1-F_L}{4} \sin^2 \theta_K \cos 2\theta_\ell - F_L \cos^2 \theta_K \cos 2\theta_\ell + S_3 \sin^2 \theta_K \sin^2 \theta_\ell \cos 2\phi + S_4 \sin 2\theta_K \sin 2\theta_\ell \cos \phi + S_5 \sin 2\theta_K \sin \theta_\ell \cos \phi + S_6 \sin^2 \theta_K \cos \theta_\ell + S_7 \sin 2\theta_K \sin \theta_\ell \sin \phi + S_8 \sin 2\theta_K \sin 2\theta_\ell \sin \phi + S_9 \sin^2 \theta_K \sin^2 \theta_\ell \sin 2\phi \right].$$



- The S parameters are translated into the $P^{(\prime)}$ parameters via

$$P_1 = \frac{2S_3}{1-F_L} \quad P'_{i=4,5,6,8} = \frac{S_{j=4,5,7,8}}{\sqrt{F_L(1-F_L)}}$$

with reduced dependence on the hadronic form factors.

Angular analysis on $B \rightarrow K^* \mu \mu$

- another way to look at FCNC: $b \rightarrow s$ transition with a BR $\sim 1.1 \cdot 10^{-6}$
- angular distribution of the 4 particles in the final state sensitive to new physics for the interference of NP and SM diagrams
- allows measuring a large set of angular parameters sensitive to Wilson coefficients $C^{(i)}_7, C^{(i)}_9, C^{(i)}_{10}, C^{(i)}_{S,P}$



- decay described by three angles (θ_L, θ_K, ϕ) and the di-muon mass squared $q^2 \rightarrow$ the angular distribution is analysed in finite bins of q^2 as a function of θ_L, θ_K and ϕ .
- LHCb reports a 3.4σ deviation from the SM.

JHEP 02 (2016) 104
arXiv:1512.04442

Angular analysis on $B \rightarrow K^* \mu \mu$

- B^0 flavour eigenstate can be identified through the $K^* \rightarrow K^- \pi^+$ decay
- angular distribution given by:

$$\frac{1}{d\Gamma/dq^2} \frac{d^4\Gamma}{d\cos\theta_\ell d\cos\theta_K d\phi dq^2} = \frac{9}{32\pi} \left[\frac{3(1-F_L)}{4} \sin^2\theta_K + F_L \cos^2\theta_K + \frac{1-F_L}{4} \sin^2\theta_K \cos 2\theta_\ell \right. \\ \left. - F_L \cos^2\theta_K \cos 2\theta_\ell + S_3 \sin^2\theta_K \sin^2\theta_\ell \cos 2\phi + S_4 \sin 2\theta_K \sin 2\theta_\ell \cos \phi \right. \\ \left. + S_5 \sin 2\theta_K \sin \theta_\ell \cos \phi + S_6 \sin^2\theta_K \cos \theta_\ell \right. \\ \left. + S_7 \sin 2\theta_K \sin \theta_\ell \sin \phi + S_8 \sin 2\theta_K \sin 2\theta_\ell \sin \phi \right. \\ \left. + S_9 \sin^2\theta_K \sin^2\theta_\ell \sin 2\phi \right].$$

- the S parameters are translated into the $P^{(\prime)}$ parameters via

$$P_1 = \frac{2S_3}{1-F_L} \quad P'_{i=4,5,6,8} = \frac{S_{j=4,5,7,8}}{\sqrt{F_L(1-F_L)}}$$





- the $P^{(\prime)}$ parameters are expected to have a reduced dependence on the hadronic form factors.
- ATLAS and CMS need to fold the angular distribution via trigonometric relations to reduce the number of free parameters

Analysis strategy for $B \rightarrow K^* \mu\mu$

- Data collected in 2012 at 8 TeV with 20.3 fb⁻¹ Run 1 data
- Measured in 6 (overlapping) bins of q^2 in the range [0.04, 6] GeV²
- 4 sets of fits for three parameters (F_L , S_3 and S_j with $j=4,5,7,8$)
- Selection of triggers with muon p_T thresholds starting at 4 GeV
- K^* tagged by the kaon sign:
 - dilution from mistag probability included in $(1-2\langle w \rangle)$:
 - $\langle w \rangle \sim 10.9(1)\%$ with small dependence on q^2
- 787 events selected with $q^2 < 6$ GeV²
- Extended unbinned maximum likelihood fits in each of the fit variants in each q^2 bin:
 - two step fit procedure: first fit the invariant mass distribution
 - then add to the fit the angular distributions to extract the F_L and $S(P)$ parameters
- Signal shape studies from control samples $K^* J/\psi$ and $K^* \psi(2S)$

Fit projections




- fit $m(K^*_{\mu\mu})$, $\cos\theta_L$, $\cos\theta_K$ and ϕ to isolate signal and extract parameters of interest.




- Data shown for $[0.04, 2.0] \text{ GeV}^2$
- projections for the S_5 fit.
- Approx 106-128 signal events in 2 GeV^2 q^2 bin.
- Similar results for the other q^2 bins and other fit variants.

Angular analysis results

- Results are compatible with theoretical calculations & fits:




q^2 [GeV ²]	P_1	P'_4	P'_5
[0.04, 2.0]	$-0.05 \pm 0.30 \pm 0.08$	$0.31 \pm 0.40 \pm 0.20$	$0.67 \pm 0.26 \pm 0.16$
[2.0, 4.0]	$-0.78 \pm 0.51 \pm 0.34$	$-0.76 \pm 0.31 \pm 0.21$	$-0.33 \pm 0.31 \pm 0.13$
[4.0, 6.0]	$0.14 \pm 0.43 \pm 0.26$	$0.64 \pm 0.33 \pm 0.18$	$0.26 \pm 0.35 \pm 0.18$
[0.04, 4.0]	$-0.22 \pm 0.26 \pm 0.16$	$-0.30 \pm 0.24 \pm 0.17$	$0.32 \pm 0.21 \pm 0.11$
[1.1, 6.0]	$-0.17 \pm 0.31 \pm 0.13$	$0.05 \pm 0.22 \pm 0.14$	$0.01 \pm 0.21 \pm 0.08$
[0.04, 6.0]	$-0.15 \pm 0.23 \pm 0.10$	$0.05 \pm 0.20 \pm 0.14$	$0.27 \pm 0.19 \pm 0.06$



OPE and LHCb data fit: CFFMPSV: Ciuchini et al.; JHEP 06 (2016) 116.
 QCD factorisation: DMVH: Decotes-Genon et al.; JHEP 12 (2014) 125.
 JC: Jäger-Camalich; Phys. Rev. D93 (2016) 014028.

Angular analysis results

- ATLAS gets deviations of about 2.5σ (2.7σ) from DHMV in P'_4 (P'_5) in $[4,6] \text{ GeV}^2$



CFFMPSV: Ciuchini et al.; JHEP 06 (2016) 116.
 DHMV: Decotes-Genon et al.; JHEP 12 (2014) 125.
 JC: Jäger-Camalich; Phys. Rev. D93 (2016) 014028.



TD angular analysis of $B_s \rightarrow J/\psi\phi$

ATL-PHYS-PUB-2018-041

Period	L_{int} [fb^{-1}]	N_{sig}	f_{sig}	Tag Power [%]	$\sigma(\tau)$ [ps]	$\delta_{\phi_s}^{\text{stat}}$ [rad] measured (extrapolated)	$\delta_{\Delta\Gamma_s}^{\text{stat}}$ [ps^{-1}] measured (extrapolated)
2012	14.3	73693	0.20	1.49	0.091	0.082	0.013
2011	4.9	22690	0.17	1.45	0.100	0.25 (0.22)	0.021 (0.023)
						$\delta_{\phi_s}^{\text{stat}}$ [rad] extrapolated	
HL-LHC	3000						
Trigger $\mu 6\mu 6$		$9.72 \cdot 10^6$	0.17	1.49	0.048	0.004	0.0011
Trigger $\mu 10\mu 6$		$5.93 \cdot 10^6$	0.17	1.49	0.044	0.005	0.0014
Trigger $\mu 10\mu 10$		$1.75 \cdot 10^6$	0.15	1.49	0.038	0.009	0.003

Around and outside B physics...


Test of the universality of τ and μ in W decays from $t\bar{t}$ events

Muons from W and muons from tau distinguished using the lifetime of the τ , through the muon transverse impact parameter, and differences in the muon transverse momentum spectra.

Tag and probe approach: tag leptons to select the events, probe muon from prompt decay, $W \rightarrow \mu\nu_\mu$, or via intermediate τ , $W \rightarrow \tau\nu_\tau \rightarrow \mu\nu_\mu\nu_\tau\nu_\tau$.

Di-leptonic $t\bar{t}$ events with either one electron and one muon ($e\text{-}\mu$ channel), or two muons ($\mu\text{-}\mu$ channel).

[arXiv:2007.14040](https://arxiv.org/abs/2007.14040) [hep-ex]



$$R(\tau/\mu) = 0.992 \pm 0.013 [\pm 0.007 \text{ (stat)} \pm 0.011 \text{ (syst)}]$$

CP violation in the SM and NP:

- $B_{(s)}$ systems are giving us a rather precise picture
- However there is some space for NP
- Could appear as new contributions in $\Delta\Gamma^{\circ}$ loop processes

$$A_q = C_{B_q} e^{2i\phi_{B_q}} A_q^{SM} e^{2i\phi_{B_q}^{\circ}}$$

$$A_q = \left(1 + \frac{A_q^{NP}}{A_q^{SM}} e^{2i(\phi_q^{NP} - \phi_q^{SM})} \right) A_q^{SM} e^{2i\phi_q^{SM}}$$

The ratio of NP/SM amplitudes need to be:

