



# Searching for dark matter decay with gamma-ray and neutrino telescopes



**DIYASELIS M. DELGADO LÓPEZ**  
**EMAIL: [ddelgado@g.harvard.edu](mailto:ddelgado@g.harvard.edu)**

**TEVPA (CHENGDU, CHINA) - PARALLEL TALK**

**OCT 27, 2021**



---

# Standard Model is great and all but...

# Standard Model is great and all but...

Need SM extension


Looking for a theory that explains observed neutrino masses and the nature of Dark Matter

Weakly-interacting massive particles (WIMPs) are a simple solution. A new particle connected to the Standard Model by a new force.

Overwhelming astrophysical and cosmological evidence for the existence of Dark Matter (DM).

Hints from bullet cluster and dwarf galaxies pointing to corpuscular nature of DM

# Indirect detection




No need for specialized detectors: **Neutrino detectors, Gamma-ray telescopes, CR-experiments**  
Search for products of dark matter decay processes: **Focus on large reservoirs of dark matter**

Figure from Juan Aguilar




# Motivation: Dark Matter Annihilation to Neutrinos



Argüelles, et al., 10.1103




# Dark Matter Decay to neutrinos





# Dark Matter Decay to neutrinos




**FLUX FROM DARK MATTER IN OUR GALAXY  
(GALACTIC CONTRIBUTION)**

$$\frac{d\Phi}{dE} = \frac{1}{4\pi} \frac{dN}{dE} D(\Omega, x) \frac{\Gamma_{\chi}}{m_{\chi}}$$



# Dark Matter Decay to neutrinos



**FLUX FROM DARK MATTER IN OUR GALAXY  
(GALACTIC CONTRIBUTION)**

$$\frac{d\Phi}{dE} = \frac{1}{4\pi} \frac{dN}{dE} D(\Omega, x) \frac{\Gamma_{\chi}}{m_{\chi}}$$


**NEUTRINO  
PRODUCTION SPECTRUM  
FOR DECAY OF DM TO  
NEUTRINOS \***

$$\frac{dN_{\nu}}{dE} = \delta\left(\frac{m_{\chi}}{2} - E_{\nu}\right)$$

**\* TRUE FOR NEUTRINOS BUT FOR GAMMAS BECOMES MORE  
COMPLICATED WITH ELECTROWEAK CORRECTIONS**



# Dark Matter Decay to neutrinos



**FLUX FROM DARK MATTER IN OUR GALAXY  
(GALACTIC CONTRIBUTION)**

$$\frac{d\Phi}{dE} = \frac{1}{4\pi} \frac{dN}{dE} D(\Omega, x) \frac{\Gamma_\chi}{m_\chi}$$

**NEUTRINO  
PRODUCTION SPECTRUM  
FOR DECAY OF DM TO  
NEUTRINOS \***


$$\frac{dN_\nu}{dE} = \delta\left(\frac{m_\chi}{2} - E_\nu\right)$$

**D FACTOR: 3D INTEGRAL  
OVER THE SKY SOLID ANGLE  
AND LINE OF SIGHT**

$$D = \int d\Omega \int_{l.o.s.} \rho_\chi(x) dx$$



# Dark Matter Decay to neutrinos



**FLUX FROM DARK MATTER IN OUR GALAXY  
(GALACTIC CONTRIBUTION)**

$$\frac{d\Phi}{dE} = \frac{1}{4\pi} \frac{dN}{dE} D(\Omega, x) \frac{\Gamma_\chi}{m_\chi}$$

**NEUTRINO  
PRODUCTION SPECTRUM  
FOR DECAY OF DM TO  
NEUTRINOS \***

$$\frac{dN_\nu}{dE} = \delta\left(\frac{m_\chi}{2} - E_\nu\right)$$

**D FACTOR: 3D INTEGRAL  
OVER THE SKY SOLID ANGLE  
AND LINE OF SIGHT**

$$D = \int d\Omega \int_{l.o.s.} \rho_\chi(x) dx$$


**DARK MATTER DENSITY:  
NFW PROFILE**

$$\rho_\chi = \frac{2^{3-\gamma} \rho_s}{\left(\frac{r}{r_s}\right)^\gamma \left(1 + \frac{r}{r_s}\right)^{3-\gamma}}$$

J. Navarro, C. Frenck, S. White, [10.1086/177173](https://arxiv.org/abs/10.1086/177173)



# Dark Matter Decay to neutrinos




**FLUX FROM DARK MATTER IN OUR GALAXY  
(GALACTIC CONTRIBUTION)**

$$\frac{d\Phi}{dE} = \frac{1}{4\pi} \frac{dN}{dE} D(\Omega, x) \frac{\Gamma_{\chi}}{n_{\chi}}$$

**WE WANT TO DETERMINE  
THE LIMIT ON THE LIFETIME**

$$\Gamma_{\chi} = \frac{1}{\tau_{\chi}}$$

# Dark Matter Decay to neutrinos



**Extragalactic contribution also plays an important role!**  
**Depends on the dark matter distribution.**

FLUX

INTEGRAL SOLID ANGLE  
 LINE OF SIGHT

$$D = \int d\Omega \int_{l.o.s.} \rho_\chi(x) dx$$

**DARK MATTER DENSITY:**


**NFW PROFILE**

$$\rho_\chi = \frac{2^{3-\gamma} \rho_s}{\left(\frac{r}{r_s}\right)^\gamma \left(1 + \frac{r}{r_s}\right)^{3-\gamma}}$$


Navarro, et al., 9508025



# Dark Matter Decay to neutrinos



# Dark Matter Search using neutrinos





	Energy Range	Experimental Analysis	Directionality	Detected Flavor
MeV	2.5 – 15 MeV	Borexino (Bellini <i>et al.</i> , 2011)	×	$\bar{\nu}_e$ (IBD)
	8.3 – 18.3 MeV	KamLAND (Gando <i>et al.</i> , 2012)	✓	$\bar{\nu}_e$ (IBD)
	10 – 40 MeV	JUNO (An <i>et al.</i> , 2016)	✓	$\bar{\nu}_e$ (IBD)
	GeV	15 – 10 <sup>3</sup> MeV	SK (Olivares-Del Campo <i>et al.</i> , 2018a)	×
DARWIN (McKeen and Raj, 2018)			×	All Flavors (Coherent)
TeV	0.1 – 30 GeV	DUNE (Abi <i>et al.</i> , 2020b)	×	$\nu_e, \bar{\nu}_e, \nu_\tau, \bar{\nu}_\tau$ (CC)
		HK (Olivares-Del Campo <i>et al.</i> , 2018b)		
	1 – 10 <sup>4</sup> GeV	SK (Abe <i>et al.</i> , 2020; Frankiewicz, 2015)	✓	All Flavors
	20 – 10 <sup>4</sup> GeV	IceCube (Aartsen <i>et al.</i> , 2016a)	✓	All Flavors
	50 – 10 <sup>5</sup> GeV	ANTARES (Adrian-Martinez <i>et al.</i> , 2015)	✓	$\nu_\mu, \bar{\nu}_\mu$ (CC)
	0.2 – 100 TeV	CTA (Queiroz <i>et al.</i> , 2016)	✓	All Flavors (Bremsstrahlung)
PeV	10 – 10 <sup>4</sup> GeV	IC-Upgrade (Baur, 2019)	✓	All Flavors
	> 10 PeV	IC Gen-2 (Aartsen <i>et al.</i> , 2014b)	✓	All Flavors
	10 – 10 <sup>4</sup> TeV	KM3Net (Adrian-Martinez <i>et al.</i> , 2016)	✓	All Flavors
	1 – 100 PeV	TAMBO (Wissel <i>et al.</i> , 2019)	✓	$\nu_\tau, \bar{\nu}_\tau$ (CC)
	EeV	> 100 PeV	GRAND (Alvarez-Muniz <i>et al.</i> , 2018)	✓

Argüelles, et al., 10.1103





# Neutrino Experiments



- Cherenkov detector at the South Pole.
- 1 gigaton of ice target with 5160 PMTs
- IceCube has a measured diffuse astrophysical neutrino flux in the TeV-PeV range.

# Neutrino Experiments



Liquid scintillator.  
Solar neutrinos (MeV)



Liquid scintillator (Reactor).  
Extraterrestrial neutrino  
fluxes (MeV)



Liquid Argon TPC.  
Atmospheric neutrino  
fluxes (GeV)



Water Cherenkov.  
Atmospheric neutrinos  
(GeV-TeV)




**ICECUBE**  
SOUTH POLE NEUTRINO OBSERVATORY

- Cherenkov detector at the South Pole.
- 1 gigaton of ice target with 5160 PMTs
- IceCube has a measured diffuse astrophysical neutrino flux in the TeV-PeV range.




Water Cherenkov.  
Atmospheric  
neutrinos (GeV-TeV)



Sea Water Cherenkov  
Extraterrestrial  
neutrino fluxes (PeV)



Water Cherenkov.  
Astrophysical Tau  
Neutrino (PeV)



PIERRE  
AUGER  
OBSERVATORY

Water Cherenkov.  
Ultra High Energy  
Cosmic Rays (EeV)




Radio Array. Tau  
Neutrinos (EeV)




# Lifetime Limits

C. Argüelles, **D. Delgado**, A. Vincent, A. Friedlander,  
H. White, A. Kheirandish, I. Safa




# Dark Matter Decay to neutrinos






# Dark Matter Search detecting gammas



# Gamma-Ray Experiments



Water Cherenkov.  
Gamma Rays and  
Cosmic Rays  
(GeV - TeV)



Hybrid Air  
Shower. Gamma  
Rays (GeV - PeV)



Air Cherenkov. High Energy  
Gamma Rays (TeV)



MeV      GeV      TeV      PeV      EeV



# Recipe for Lifetime Limits

## EXPECTED BACKGROUND (GAMMA-RAYS)


- Gamma-Ray Sensitivities from the experiments
- Effective Areas and Observation Time
- Expected Number of Events

## EXPECTED SIGNAL (DARK MATTER)

- Neutrino Production Spectrum for Decay of DM to neutrinos
- Dark Matter Expected Flux (Galactic & Extragalactic)
- Expected Number of Events


$$N_{evt} = \int dE A_{eff} \frac{d\Phi}{dE} T_{obs}$$

# Gamma-Ray Experimental Sensitivities







# Expected gamma flux (Galactic)





All Sky,  $\tau_\chi = 10^{26}$  s



All Sky,  $\tau_\chi = 10^{26}$  s



# Expected gamma flux (Extragalactic)



---

# Summary

- The nature of Dark Matter and origin of neutrino mass remain a mystery.
- Dark Matter neutrino connections offer solutions to both problems.
- We can measure both neutrinos and gamma-rays as final products of Dark matter decay to neutrinos → Correlated signal.
- Major experimental advances in neutrino and gamma-ray detection allows us to explore a wide mass range (MeV - ZeV).
- New constraints for gamma rays lifetime limits will be reported on an upcoming paper. Stay tuned!



---

# Backup

# Electro weak corrections: why gamma-rays?

- The standard  $1 \rightarrow 2$  decay process is  $\chi \rightarrow \bar{\nu}\nu$ .
- Higher orders involve the bremsstrahlung of an electroweak gauge boson.
- The branching ratio  $R = \sigma(\chi \rightarrow \bar{\nu}\nu W) / \sigma(\chi \rightarrow \bar{\nu}\nu)$  only depends generally only on the details of the underlying  $1 \rightarrow 2$  process for  $Q^2 \sim m_\chi^2$ .
- We have three cases:
  1. Fermi regime  $m_\chi \lesssim m_W$
  2. Perturbative electroweak regime  $m_\chi \lesssim m_W \lesssim 10^6 \text{ GeV}$
  3. Non-perturbative regime where large logarithms over-compensate the small electroweak coupling  $\alpha_2$

Kachelrieß, et al.,0707.0209

# Converting Gamma-Ray Diffuse Flux Limits to Limits on the Dark Matter Differential Spectrum

- The reported gamma-ray flux limit,  $\left. \frac{d\phi}{dE} \right|_{lim} \equiv f_0 E^{-\alpha}$ , for which the actual limit at the bin center  $E = \bar{E}$  is:

$$\phi_{lim}(\bar{E}) = 4\pi \int_{a_-}^{a^+} f_0 E^{-\alpha} dE \quad \text{with } a_{\pm} \equiv \bar{E} 10^{\pm\Delta/2}$$

$\Delta$  is the bin width.



# Converting Gamma-Ray Diffuse Flux Limits to Limits on the Dark Matter Differential Spectrum

- The reported gamma-ray flux limit,  $\left. \frac{d\phi}{dE} \right|_{lim} \equiv f_0 E^{-\alpha}$ , for which the actual limit at the bin center  $E = \bar{E}$  is:

$$\phi_{lim}(\bar{E}) = 4\pi \int_{a_-}^{a^+} f_0 E^{-\alpha} dE \quad \text{with } a_{\pm} \equiv \bar{E} 10^{\pm\Delta/2}$$

$\Delta$  is the bin width.

- Dark matter flux is given by:

$$\phi = \int dE \frac{1}{4\pi} \frac{1}{m_{\chi} \tau_{\chi}} \frac{dN_{\gamma}}{dE} D(\Omega, x)$$

---

# Calculating lifetime limits

- We determine the number of events from the following equation:

$$N_{evt} = \int dE A_{eff} \Phi T_{obs}$$

# Calculating lifetime limits

- We determine the number of events from the following equation:

$$N_{evt} = \int dE A_{eff} \Phi T_{obs}$$


- Equate using both Gamma-ray flux and Expected Dark Matter Flux:


$$N_{evt} = \int dE A_{eff} \frac{1}{4\pi} \frac{1}{m_\chi \tau_\chi} \frac{dN_\gamma}{dE} D(\Omega, x) T_{obs} = \int_{a_-}^{a^+} dE A_{eff} \underline{f_0} E^{-\alpha} T_{obs}$$

- We are interested in the limit of  $\tau_\chi$  that makes the equation true.



# Expected neutrino flux





IceCube Collaboration: 10.1140/epjc/s10052-018-6273-3