



Improvement of single π^- reconstruction with CGEM+ODC

Liangliang Wang (IHEP)



III Assumptions in CGEM simulation

- 100% cluster efficiency
- Spatial resolution 130 μm in both X and V direction

Recent activities to improve soft π^- reconstruction

- New Least-Square global track fitting used (circle and helix fitting)
 - ✓ circle fitting rejects hits with large χ^2
 - ✓ helix fitting rejects outermost hits if χ^2 is large => favor track segment near IP
=> track parameters at IP
 - ✓ latest tag: [DotsConnection-00-00-04](#)

- Tuning HoughTransAlg for π^- with $p_T=50$ MeV/c
 - ✓ Circle search/reconstruction criteria loosen => keep efficiency high
 - ✓ V-hits association procedure modified
 - ✓ latest tag: [HoughTransAlg-00-00-16](#)



III Good track and tracking efficiency

- Good track: $|dr| < 1.0\text{cm}$, $|dz| < 10\text{cm}$, $|\cos\theta| < 0.93$, correct charge
- Tracking efficiency for single track events:

$$\varepsilon = N_{\text{good}}/N_{\text{gen}}$$



where N_{good} is the number of events with one or more good tracks reconstructed,
 N_{gen} is the number of events generated/simulated

Tracking efficiency for single π^-



Improved significantly!

Kalman filter success rate for good π^- track



$$\text{Success rate} = N_{\text{success}} / N_{\text{good}}$$

Under investigation