

Precision QCD Calculations for Heavy Quark Decays

Yu-Ming Wang

Nankai University

The 6th China LHC Physics Workshop
November 8, 2020

Why precision calculations?

- Understanding the general properties of power expansion in EFTs (HQET, SCET, NRQCD).
- Interesting to understand the strong interaction dynamics of heavy quark decays.
 - ▶ Factorization properties of the subleading-power amplitudes.
 - ▶ Renormalization and asymptotic properties of the higher-twist B -meson DAs.
 - ▶ Interplay of different QCD techniques.
- Precision determinations of the CKM matrix elements $|V_{ub}|$ and $|V_{cb}|$.
Power corrections, QED corrections, BSM physics.
- Crucial to understand the CP violation in B -meson decays.
Strong phase of $\mathcal{A}(B \rightarrow M_1 M_2)$ @ m_b scale in the leading power.
- Indispensable for understanding the flavour puzzles (continuously updated).
 - ▶ P'_5 and $R_{K^{(*)}}$ anomalies in $B \rightarrow K^{(*)} \ell^+ \ell^-$.
 - ▶ $R_{D^{(*)}}$ anomalies in $B \rightarrow D^{(*)} \ell \bar{\nu}_\ell$.
 - ▶ Color suppressed hadronic B -meson decays.
 - ▶ Polarization fractions of penguin dominated $B_{(s)} \rightarrow VV$ decays.

Highlights of Heavy Quark Physics: Biased Summary

- B -meson light-cone distribution amplitudes:
 - ▶ Matching heavy-meson LCDA from QCD to HQET [Ishaq, Jia, Xiong, Yang, 2020].
 - ▶ Determining the B -meson LCDA with the lattice QCD simulation [Wang, YMW, Xu, Zhao, 2020].
- Higher-order QCD/QED corrections to hadronic B -meson decays:
 - ▶ NNLO QCD correction to the QCD penguin amplitude [Bell, Beneke, Huber, Li, 2020].
 - ▶ QED factorization with generalized distribution amplitudes [Beneke, Böer, Toelstede, Vos, 2020].
- Subleading power corrections to exclusive B -meson decays:
 - ▶ Systematic calculations of the helicity form factors for $B_q \rightarrow \gamma \ell \bar{\ell}$ [Beneke, Bobeth and YMW, 2020].
 - ▶ A complete NLL factorization at LP and the LO factorization at NLP for $B_q \rightarrow \gamma \gamma$ [Shen, YMW, Wei, 2020].
The so-called “soft form factor” defined in [Beneke, Rohrwild, 2011] is factorizable!
- BSM searches with the FCNC B -meson decays:
 - ▶ Novel determination of the photon helicity in $B \rightarrow K_1 \gamma$ [Wang, Yu, Zhao, 2020].
- Apologies for omitting many interesting topics due to the limited time.

Theory tools for precision flavor physics

New Physics: $\mathcal{L}_{NP}$

↓

EW scale (m_W): $\mathcal{L}_{SM} + \mathcal{L}_{D>4}$

↓

Heavy-quark scale (m_b): $\mathcal{L}_{eff} = -\frac{G_F}{\sqrt{2}} \sum_i C_i Q_i + \mathcal{L}_{eff,D>6}$

↓

QCD scale (Λ_{QCD})

- Aim: $\langle f | Q_i | \bar{B} \rangle = ?$
- QCD factorization [Diagrammatic approach].
- SCET factorization [Operator formalism].
- TMD factorization.
- (Light-cone) QCD sum rules.
- Lattice QCD.

- Key concepts: Factorization, Resummation, Evolution.

B-meson distribution amplitudes

- The light-ray HQET matrix element [Grozin, Neubert, 1997]:

$$\langle 0 | \bar{q}_\beta(z) [z, 0] h_{v\alpha}(0) | \bar{B}(v) \rangle = -\frac{i\tilde{f}_B m_B}{4} \left[\frac{1 + \not{v}}{2} \left\{ 2\tilde{\phi}_B^+(t, \mu) + \frac{\tilde{\phi}_B^-(t, \mu) - \tilde{\phi}_B^+(t, \mu)}{t} \not{z} \right\} \gamma_5 \right]_{\alpha\beta}.$$

- Evolution equation at one loop [Lange, Neubert, 2003]:

$$\frac{d\phi_B^+(\omega, \mu)}{d\ln \mu} = -\left[\Gamma_{\text{cusp}}(\alpha_s) \ln \frac{\mu}{\omega} + \gamma_+(\alpha_s) \right] \phi_B^+(\omega, \mu) - \omega \int_0^\infty d\eta \Gamma_+(\omega, \eta, \alpha_s) \phi_B^+(\eta, \mu).$$

This is an integro-differential equation!

- (Relatively) complicated solution [Lee, Neubert, 2005]:

$$\begin{aligned} \phi_B^+(\omega, \mu) &= e^{V-2\gamma_E g} \frac{\Gamma(2-g)}{\Gamma(g)} \int_0^\infty \frac{d\eta}{\eta} \phi_B^+(\eta, \mu_0) \left(\frac{\max(\omega, \eta)}{\mu_0} \right)^g \\ &\quad \times \frac{\min(\omega, \eta)}{\max(\omega, \eta)} {}_2F_1 \left(1-g, 2-g, 2, \frac{\min(\omega, \eta)}{\max(\omega, \eta)} \right), \\ g(\mu, \mu_0) &= \int_{\alpha_s(\mu_0)}^{\alpha_s(\mu)} d\alpha \frac{\Gamma_{\text{cusp}}(\alpha)}{\beta(\alpha)} \approx \frac{2C_F}{\beta_0} \ln \frac{\alpha_s(\mu_0)}{\alpha_s(\mu)}. \end{aligned}$$

Making the QCD resummation for enhanced logarithms complicated!

B -meson distribution amplitudes

- Fourier/Mellin transformation:

$$\varphi_B^+(\theta, \mu) = \int_0^\infty \frac{d\omega}{\omega} \left(\frac{\omega}{\mu}\right)^{-i\theta} \phi_B^+(\omega, \mu) \quad \Leftrightarrow \quad \phi_B^+(\omega, \mu) = \int_{-\infty}^\infty \frac{d\theta}{2\pi} \left(\frac{\omega}{\mu}\right)^{i\theta} \varphi_B^+(\theta, \mu).$$

- Solution to the RGE in Mellin space:

$$\varphi_B^+(\theta, \mu) = e^{V-2\gamma_E g} \left(\frac{\mu}{\mu_0}\right)^{i\theta} \frac{\Gamma(1-i\theta)}{\Gamma(1+i\theta)} \cdot \frac{\Gamma(1+i(\theta+ig))}{\Gamma(1-i(\theta+ig))} \varphi_B^+(\theta+ig, \mu_0).$$

Already very symmetric solution in Mellin space.

- Yet simpler solution to the integral-differential equation exists?

$\Leftrightarrow$ Eigenfunctions of the Lange-Neubert kernel [Bell, Feldmann, YMW and Yip, 2013].

$$\begin{aligned} \varphi_B^+(\theta, \mu) &:= \frac{\Gamma(1-i\theta)}{\Gamma(1+i\theta)} \bar{\rho}_B^+(\theta, \mu) \\ &= \frac{\Gamma(1-i\theta)}{\Gamma(1+i\theta)} \int_0^\infty \frac{d\omega'}{\omega'} \rho_B^+(\omega', \mu) \left(\frac{\omega'}{\mu}\right)^{-i\theta}. \end{aligned}$$

B -meson distribution amplitudes

- Linear differential equation [Bell, Feldmann, YMW and Yip, 2013]:

$$\frac{d\rho_B^+(\omega', \mu)}{d\ln \mu} = - \left[\Gamma_{\text{cusp}}(\alpha_s) \ln \frac{\mu}{\omega'} + \gamma_+(\alpha_s) \right] \rho_B^+(\omega', \mu).$$

Local evolution in the dual space!

- Integral transformation:

$$\begin{aligned}\phi_B^+(\omega, \mu) &= \int_0^\infty \frac{d\omega'}{\omega'} \sqrt{\frac{\omega}{\omega'}} J_1 \left(2 \sqrt{\frac{\omega}{\omega'}} \right) \rho_B^+(\omega', \mu), \\ \rho_B^+(\omega', \mu) &= \int_0^\infty \frac{d\omega}{\omega} \sqrt{\frac{\omega}{\omega'}} J_1 \left(2 \sqrt{\frac{\omega}{\omega'}} \right) \phi_B^+(\omega, \mu).\end{aligned}$$

Eigenfunction of the Lange-Neubert kernel at one-loop is the **Bessel function**!

- Solution to the RGE in dual space [Bell, Feldmann, YMW, Yip, 2013]:

$$\rho_B^+(\omega', \mu) = e^V \left(\frac{\mu_0}{\omega'} \right)^{-g} \rho_B^+(\omega', \mu_0).$$

Very compact expression in a full analytical form!

B -meson distribution amplitudes

- **Collinear conformal symmetry** for the Lange-Neubert kernel [Braun, Manashov, 2014]:

$$\left(\frac{d}{d \ln \mu} + \mathcal{H}_{\text{LN}} \right) O_+(z, \mu) = 0.$$

$\mathcal{H}_{\text{LN}}$ is almost determined by the commutation relations completely [Knoedlseder, Offen, 2011]

$$[S_+, \mathcal{H}_{\text{LN}}] = 0, \quad [S_0, \mathcal{H}_{\text{LN}}] = 1.$$

The beautiful solution in terms of S_+ :

$$\mathcal{H}_{\text{LN}} = \ln(i\mu S^+) - \psi(1) - \frac{5}{4}.$$

- Generators of the collinear conformal group:

$$S_+ = z^2 \partial_z + 2jz, \quad S_0 = z \partial_z + j, \quad S_- = -\partial_z.$$

Eigenfunctions of S_+ [Braun, Manashov, 2014]:

$$\begin{aligned} iS_+ Q_s(z) &= s Q_s(z), & Q_s(z) &= -\frac{1}{z^2} e^{is/z}. \\ \langle e^{-i\omega z} | Q_s(z) \rangle &= \frac{1}{\sqrt{\omega s}} J_1(2\sqrt{\omega s}). \end{aligned}$$

Wide applications of the conformal symmetry in high energy physics!

B-meson distribution amplitudes

- RG evolution of $\phi_B^+(\omega, \mu)$ at two loops [Braun, Ji, Manashov, 2019; Liu, Neubert, 2020]:

$$\frac{d\phi_B^+(\omega, \mu)}{d\ln\mu} = \left[\Gamma_{\text{cusp}}(\alpha_s) \ln \frac{\omega}{\mu} - \gamma_\eta(\alpha_s) \right] \phi_B^+(\omega, \mu) + \Gamma_{\text{cusp}}(\alpha_s) \int_0^\infty dx \Gamma(1, x) \phi_B^+(\omega/x, \mu) + \left(\frac{\alpha_s}{2\pi} \right)^2 C_F \int_0^1 \frac{dx}{1-x} h(x) \phi_B^+(\omega/x, \mu).$$

The last missing element for the NLL predictions of exclusive B-meson decay observables!

- The two-loop eigenfunctions depend on the strong coupling α_s [Braun, Ji, Manashov, 2019].
- Applying the Laplace transformation of the LCDA [Galda, Neubert, 2020]

$$\tilde{\phi}_B^+(\eta, \mu) = \int_0^\infty \frac{d\omega}{\omega} \phi_B^+(\omega, \mu) \left(\frac{\omega}{\bar{\omega}} \right)^{-\eta},$$

$\Rightarrow$ the general solution to the two-loop RGE of $\phi_B^+(\omega, \mu)$



$$\begin{aligned} \tilde{\phi}_B^+(\eta, \mu) &= \exp \left[S(\mu_0, \mu) + a_\gamma(\mu_0, \mu) + 2 \gamma_E a_\Gamma(\mu_0, \mu) \right] \left(\frac{\bar{\omega}}{\mu_0} \right)^{-a_\Gamma(\mu_0, \mu)} \\ &\times \frac{\Gamma(1 + \eta + a_\Gamma(\mu_0, \mu)) \Gamma(1 - \eta)}{\Gamma(1 - \eta - a_\Gamma(\mu_0, \mu)) \Gamma(1 + \eta)} \exp \left[\int_{\alpha_s(\mu_0)}^{\alpha_s(\mu)} \frac{d\alpha}{\beta(\alpha)} \mathcal{G}(\eta + a_\Gamma(\mu_\alpha, \mu), \alpha) \right] \\ &\times \tilde{\phi}_B^+(\eta + a_\Gamma(\mu_0, \mu), \mu_0). \end{aligned}$$

B -meson distribution amplitudes

- Better determination of the B -meson LCDA essential to precision predictions.
 - ▶ **Dominant uncertainties** for QCD calculations of exclusive B -meson decays.
 - ▶ The only **QCD-motivated construction** from the sum rule approach [Braun, Ivanov, Korchemsky, 2004].
 - ▶ Drawback I: can only determine $\tilde{\phi}_B^+(t, \mu_0)$ for t around $(1-3) \text{ GeV}^{-1}$.
 - ▶ Drawback II: sensitive to the parton-hadron duality ansatz.
- Model independent determination of the B -meson LCDA in HQET.
 - ▶ Key observation: **Time-independent quasi-quantities** calculable on the Euclidean lattice.
 - ▶ Construct the relations between quasi-distribution functions and light-cone matrix elements.
 $\Leftrightarrow$ Establish **perturbative QCD factorization theorems** for the quasi-quantities.
 - ▶ Ji's proposal under the name of the large-momentum effective theory (LaMET) [Ji, 2013].
 - ▶ Wide applications in the determinations of the nucleon PDF and the pion LCDA [Cichy, Constantinou, 2019; Ji, Liu, Liu, Zhang, Zhao, 2020].
 - ▶ Alternative proposals with the aid of the coordinate-space factorization approach, pseudo-distribution functions, etc.
 - ▶ **Determining the B -meson LCDA from the lattice QCD simulation technique** [Wang, YMW, Xu, Zhao, 2020].

B-meson distribution amplitudes

- The characteristic feature of $\phi_B^+(\xi, \mu)$ [Wang, YMW, Xu, Zhao, 2020]:



- Two different models for the B -meson LCDA:

$$\phi_{B,I}^+(\omega, \mu_0) = \frac{\omega}{\omega_0^2} e^{-\omega/\omega_0},$$

$$\phi_{B,II}^+(\omega, \mu_0) = \left[\frac{1}{k^2 + 1} - \frac{2(\sigma_B^{(1)} - 1)}{\pi^2} \ln k \right] \times \frac{4}{\pi \omega_0} \frac{k}{k^2 + 1}, \quad k = \frac{\omega}{1.0 \text{ GeV}}.$$


- Small $\omega = \xi$ region excluded to validate the light-cone OPE.
- $\phi_B^+(\xi, \mu)$ at large ξ region can be computed perturbatively in QCD.

- Further theory developments from both the continuum and lattice QCD aspects in demand.

- Perturbative matching between different regularization schemes.
- Lattice simulations for the hadronic matrix elements defined with the moving Wilson lines.
- Uncertainties due to the finite lattice spacing and the truncation of Fourier transformation

Radiative leptonic B meson decays in QCD

- Schematic structure of the distinct mechanisms:



A: hard subgraph that includes both photon and W^* vertices

B: real photon emission at large distances

C: Feynman mechanism: soft quark spectator

$$\left(\frac{\Lambda}{m_b}\right)^{1/2} + \left(\frac{\Lambda}{m_b}\right)^{3/2} + \dots$$

$$\left(\frac{\Lambda}{m_b}\right)^{3/2} + \dots$$

$$\left(\frac{\Lambda}{m_b}\right)^{3/2} + \dots$$

- Operator definitions of different terms needed for an unambiguous classification.

Current status of $B \rightarrow \gamma \ell \bar{\nu}_\ell$


- **Factorization properties at leading power** [Korchensky, Pirjol, Yan, 2000; Descotes-Genon, Sachrajda, 2002; Lunghi, Pirjol, Wyler, 2003; Bosch, Hill, Lange, Neubert, 2003].
- Leading power contributions at NLL and **(partial)-subleading power corrections at tree level** [Beneke, Rohrwild, 2011].
- **Subleading power corrections from the dispersion technique:**
 - ▶ Soft two-particle correction **at tree level** [Braun, Khodjamirian, 2013].
 - ▶ Soft two-particle correction **at one loop** [YMW, 2016].
 - ▶ **Three-particle B -meson DA's contribution** at tree level [YMW, 2016; Beneke et al, 2018].
 - ▶ Subleading effective current and twist-5 and 6 corrections at tree level. [Beneke et al, 2018].
- **Subleading power corrections from the direct QCD approach:**
 - ▶ Hadronic photon corrections **at tree level** up to the twist-4 accuracy [Khodjamirian, Stoll, Wyler, 1995; Ali, Braun, 1995; Eilam, Halperin, Mendel, 1995].
 - ▶ Hadronic photon corrections of **twist-two at one loop** and of **higher-twist at tree level** [Ball, Kou, 2003; YMW, Shen, 2018].
- **Leading power contributions at NNLL and the updated NLP corrections:**
 - ▶ Two-loop RG evolution of $\phi_B^+(\omega, \mu)$ derived in [Braun, Ji, Manashov, 2019].
 - ▶ Two-loop jet function obtained in [Liu, Neubert, 2020].
 - ▶ Further improvement on [Shen, Wei, Zhao, Zhou, 2020] will appear soon [Cui, Jin, Shen, Wang, YMW, Wei, 2020].

Theory predictions for $B \rightarrow \gamma \ell \bar{\nu}_\ell$

- Integrated decay rate $\Delta BR(E_{\text{cut}})$:

$$\Delta BR(E_{\text{cut}}) = \tau_B \int_{E_{\text{cut}}}^{m_B/2} dE_\gamma \frac{d\Gamma}{dE_\gamma} (B \rightarrow \gamma \ell \bar{\nu}_\ell).$$


- $\lambda_B(\mu_0)$ dependence of $\Delta BR(E_{\text{cut}})$ [YMW, Shen, 2018]:



- Belle 2015 data:
 $\Delta BR(1 \text{ GeV}) < 3.5 \times 10^{-6}$.
- Belle 2018 data
[arXiv:1810.12976]:
 $\Delta BR(1 \text{ GeV}) =$
 $(1.4 \pm 1.0 \pm 0.4) \times 10^{-6}$.
- Expected statistical error for
 $\Delta BR(1 \text{ GeV})$ with 50 ab^{-1} of
Belle-II data: $^{+0.18}_{-0.17} \times 10^{-6}$.
- The photon-energy cut not sufficiently large.**
Power corrections numerically important for $E_\gamma < 1.5 \text{ GeV}$.

Double Radiative B_q -Meson Decays

- Leading-order contributions at $\mathcal{O}(\alpha_s^0)$:



Kinematics:

$$p_\mu = \frac{n \cdot p}{2} \bar{n}_\mu \equiv \frac{m_{B_q}}{2} \bar{n}_\mu, \quad q_\mu = \frac{\bar{n} \cdot q}{2} n_\mu \equiv \frac{m_{B_q}}{2} n_\mu.$$

Interplay of the soft and collinear QCD dynamics!

- Decay amplitude:

$$\mathcal{A}(\bar{B}_q \rightarrow \gamma\gamma) = -\frac{4G_F}{\sqrt{2}} \frac{\alpha_{\text{em}}}{4\pi} \epsilon_1^{*\alpha}(p) \epsilon_2^{*\beta}(q) \sum_{p=u,c} V_{pb} V_{pq}^* \sum_{i=1}^8 C_i T_{i,\alpha\beta}^{(p)}.$$

Hadronic tensors:

$$\begin{aligned} T_{7,\alpha\beta} &= 2\bar{m}_b(v) \int d^4x e^{iq \cdot x} \langle 0 | T \left\{ j_\beta^{\text{em}}(x), \bar{q}_L(0) \sigma_{\mu\alpha} p^\mu b_R(0) \right\} | \bar{B}_q(p+q) \rangle \\ &\quad + [p \leftrightarrow q, \alpha \leftrightarrow \beta], \\ T_{i,\alpha\beta}^{(p)} &= -(4\pi)^2 \int d^4x \int d^4y e^{ip \cdot x} e^{iq \cdot y} \langle 0 | T \left\{ j_\alpha^{\text{em}}(x), j_\beta^{\text{em}}(y), P_i^{(p)}(0) \right\} | \bar{B}_q(p+q) \rangle, \\ &\quad (i = 1, \dots, 6, 8). \end{aligned}$$


Main task: Construct the SCET factorization formulae beyond the leading power.

Current status of $B_q \rightarrow \gamma\gamma$

- QCD factorization at leading power in Λ/m_b and at NLO in α_s [Descotes-Genon, Sachrajda, 2003].
 - ▶ No collinear strong interaction dynamics at LP.
 - ▶ The two-loop $b \rightarrow q\gamma$ matrix elements of QCD penguin operators NOT included.
 $\Rightarrow$ A complete factorization-scale independence at NLO is absent!
- Subleading power contributions from the weak annihilation diagrams [Bosch, Buchalla, 2002].
 - ▶ Complex perturbative hard functions evaluated at one loop.
 - ▶ Diagrammatic factorization established at two loops.
- Previous model-dependent calculations introduce additional systematic uncertainties.
- The new (technical)-ingredients from [Shen, YMW, Wei, 2020]:
 - ▶ A complete NLL calculation for the LP contribution $\Rightarrow$ 2-loop evolution of ϕ_B^+ .
 - ▶ The NLP factorization for the energetic photon radiation off the light quark.
 - ▶ The NLP factorization for the light-quark mass effect.
 - ▶ The NLP factorization for the SCET current $J^{(A2)} \supset (\bar{\xi}_{hc} W_{hc}) \gamma_\alpha^\perp P_L \left(\frac{i \vec{D}_\perp}{2m_b} \right) h_v$.
 - ▶ The NLP factorization for the subleading twist B -meson LCDAs.
 - ▶ The resolved photon contribution with the dispersion technique.

The helicity amplitudes of $B_q \rightarrow \gamma\gamma$


- Breakdown of the various QCD mechanisms [Shen, YMW, Wei, 2020]:



- NLL effects stabilize the factorization-scale dependence.
- Factorizable NLP effects around $\mathcal{O}(30\%)$.
- Destructive effects from the NLP soft corrections.
- Strong phase from the 2-loop matrix element of P_2^c and the weak annihilation.

Phenomenological observables for $B_q \rightarrow \gamma\gamma$

- Time-integrated branching fraction [Shen, YMW, Wei, 2020]:



- The yielding theory predictions

$$\mathcal{BR}(B_d \rightarrow \gamma\gamma) = \left(1.44^{+1.35}_{-0.80}\right) \times 10^{-8}, \quad \mathcal{BR}(B_s \rightarrow \gamma\gamma) = \left(3.17^{+1.96}_{-1.74}\right) \times 10^{-7}.$$

with the dominant uncertainties from λ_{B_q} , $\hat{\sigma}_{B_q}^{(1)}$, $\hat{\sigma}_{B_q}^{(2)}$ and the QCD renormalization scale v .

- The ratio of the two branching ratios for $B_{d,s} \rightarrow \gamma\gamma$


$$\frac{\mathcal{BR}(B_s \rightarrow \gamma\gamma)}{\mathcal{BR}(B_d \rightarrow \gamma\gamma)} = 33.80 \left(\frac{\lambda_{B_d}}{\lambda_{B_s}}\right)^2 + \mathcal{O}\left(\frac{\Lambda}{m_b}, \alpha_s\right).$$

The λ_{B_q} -scaling violation effect due to the NLP contributions approximately (10–20) %.

Phenomenological observables for $B_q \rightarrow \gamma\gamma$

- Time-dependent CP asymmetries [Shen, YMW, Wei, 2020]:

$$A_{\text{CP}}^{\chi}(t) = \frac{\bar{\Gamma}^{\chi}(\bar{B}_q(t) \rightarrow \gamma\gamma) - \Gamma^{\chi}(B_q(t) \rightarrow \gamma\gamma)}{\bar{\Gamma}^{\chi}(\bar{B}_q(t) \rightarrow \gamma\gamma) + \Gamma^{\chi}(B_q(t) \rightarrow \gamma\gamma)} = -\frac{\mathcal{A}_{\text{CP}}^{\text{dir},\chi} \cos(\Delta m_q t) + \mathcal{A}_{\text{CP}}^{\text{mix},\chi} \sin(\Delta m_q t)}{\cosh(\Delta\Gamma_q t/2) + \mathcal{A}_{\Delta\Gamma}^{\chi} \sinh(\Delta\Gamma_q t/2)}.$$




- $|\mathcal{A}_{\text{CP}}^{\text{dir},\chi}|$ and $|\mathcal{A}_{\text{CP}}^{\text{mix},\chi}|$ around (10–40)% for $B_d \rightarrow \gamma\gamma$.
- Tiny CP asymmetries for $B_s \rightarrow \gamma\gamma$ as expected.
- The difference between $\mathcal{A}_{\text{CP}}^{\text{dir},||}$ and $\mathcal{A}_{\text{CP}}^{\text{dir},\perp}$ due to the NLP corrections.

Radiative leptonic $B_q \rightarrow \gamma \ell \bar{\ell}$ decays

- Systematic calculations for the helicity form factors [Beneke, Bobeth and YMW, 2020]:

- ▶ SCET factorization for the LP contribution at $\mathcal{O}(\alpha_s)$.
- ▶ The NLP corrections from the real or virtual photon radiation off the b -quark at $\mathcal{O}(\alpha_s^0)$.
- ▶ The NLP corrections from the hard-collinear light-quark propagator at $\mathcal{O}(\alpha_s^0)$.
- ▶ The NLP corrections from the weak-annihilation diagrams at $\mathcal{O}(\alpha_s^0)$.

- Three possible contractions of the four-quark operators:




- ▶ The left and middle diagrams are defined as the A and B -type insertions (also for P_7).
- ▶ Need the more general $B \rightarrow \gamma^{(*)}$ form factors [YMW, 2016].
- ▶ Factorization works for $q^2 \leq 6 \text{ GeV}^2$ (excluding the narrow light-meson resonance region).
- ▶ Strong duality violation of $B_s \rightarrow \gamma \ell \bar{\ell}$ due to the narrow width of $\phi(1020)$.

- Previous investigations with different techniques/methods [Guadagnoli, Reboud and Zwicky, 2017; Albrecht, Stamou, Ziegler, Zwicky, 2019; Kozachuk, Melikhov, Nikitin, 2018; etc].

Phenomenological observables for $B_q \rightarrow \gamma \ell \bar{\ell}$

- Theory predictions for $d\mathcal{B}/dq^2$ and $\mathcal{A}_{\text{FB}}(q^2)$ [Beneke, Bobeth and YMW, 2020].



- Adding successively in quadrature the uncertainties due to $\mu_{h,hc}$ [green], $(\lambda_{B_q}, \hat{\sigma}_{B_q}^{(1)})$ [red], r_{LP} [blue].
- The largest uncertainty due to the LCDA parameters: about $^{+70}_{-30}\%$ for $\mathcal{B}(B_s \rightarrow \gamma \ell \bar{\ell})$ in the q^2 -bin $[2.0, 6.0] \text{GeV}^2$.

Heavy-to-light B -meson decay form factors

- QCD/SCET factorization formulae [BBNS, BPRS, and many others].


$$F_i^{B \rightarrow M}(E) = C_i^{(A0)}(E) \xi_a(E) + \int_0^\infty \frac{d\omega}{\omega} \int_0^1 dv \underbrace{T_i(E; \ln \omega, v)}_{C_i^{(B1)} * J_i} \phi_B^+(\omega) \phi_M(v).$$

Need the hadronic matrix elements of both the LP and NLP SCET currents!

- Diagrammatic factorization for heavy-to-light form factors at one loop [Beneke, Feldmann, 2001].
- Perturbative calculations of the hard matching coefficients $C_i^{(A0)}(E)$:
 - ▶ One-loop SCET computations in [Bauer, Fleming, Pirjol, Stewart, 2001; Beneke, Kiyo, Yang, 2004].
 - ▶ Two-loop SCET computations in [Bonciani, Ferroglia, 2008; Asatryan, Greub, Pecjak, 2008; Beneke, Huber, Li, 2009; Bell, 2009; Bell, Beneke, Huber, Li, 2011].
- Perturbative calculations of the hard matching coefficients $C_i^{(B1)}$:
 - ▶ Infrared subtractions complicated by the appearance of evanescent operators and the D -dimensional Fierz transformation.
 - ▶ One-loop SCET computations in [Becher, Hill, 2004; Hill, Becher, Lee, Neubert, 2004; Beneke, Yang, 2006].

Semileptonic $B \rightarrow V$ form factors in QCD

- A long-standing puzzle [Bell, Beneke, Huber, Li, 2011]:




Discrepancies between the SCET predictions (blue solid) and the LCSR results with light vector meson DAs (black dashed-dotted) [Ball, Zwicky, 2005].

- Subleading power correction in heavy quark expansion, systematic uncertainties?

Semileptonic $B \rightarrow V$ form factors in QCD

- QCD factorization (yellow) versus SCET sum rules (red) [Gao, Lü, Shen, YMW, Wei, 2019]:



- The underlying mechanism responsible for the discrepancies:

$$\begin{aligned} \mathcal{R}_{1,\text{LCSR}} &= 1 + (-0.049) \Big|_{C_i^{(\text{A0})}} + (+0.054) \Big|_{C_i^{(\text{B1})}} + (-3.5 \times 10^{-5}) \Big|_{3\text{PHT}}, \\ \mathcal{R}_{1,\text{QCDF}} &= 1 + (-0.023) \Big|_{C_i^{(\text{A0})}} + (+0.086) [1 + \mathcal{O}(\alpha_s)] \Big|_{C_i^{(\text{B1})}}. \end{aligned}$$

- (a) No perturbative expansion for the A0-type SCET_I form factor in QCDF.
- (b) Almost a factor of two smaller prediction of the B1-type SCET_I form factor in LCSR.

QCD/SCET sum rules with heavy-hadron LCDA

- **LO QCD calculations** of $B \rightarrow M$ form factors [Khodjamirian, Offen, Mannel, 2006; Faller, Khodjamirian, Klein, Mannel, 2008; Gubernari, Kokulu, van Dyk, 2018].
- **NLO QCD calculations** of $B \rightarrow \pi$ form factors **at (two-particle) twist-3** [De Fazio, Feldmann, Hurth, 2006, 2008; YMW, Shen, 2015].
- **NLO QCD calculations** of $B \rightarrow \pi, K$ form factors **at (two-particle) twist-3** and LO QCD calculations of **higher-twist corrections** up to twist-6 [Lü, Shen, YMW, Wei, 2018].
- **NLO QCD calculations** of $B \rightarrow D$ form factors **at (two-particle) twist-3** and LO QCD calculations of **higher-twist corrections** up to twist-6 [YMW, Wei, Shen, Lü, 2017].
 - ▶ NLO twist-3 jet function complicated by two distinct hard-collinear variables.
 - ▶ Power-enhanced charm-quark mass effect.
 - ▶ Updated subleading power corrections at twist-6 in progress.
- **NLO QCD calculations** of $B \rightarrow V$ form factors **at (two-particle) twist-3** and LO QCD calculation of **higher-twist corrections** up to twist-6 [Gao, Lü, Shen, YMW, Wei, 2019].
 - ▶ **Rigorous perturbative matching** with the evanescent-operator approach.
 - ▶ **First SCET computation** of the light-quark mass effect.
 - ▶ Three-particle higher-twist corrections **compatible with the EOM constraints**.
- **NLO QCD calculations** of $\Lambda_b \rightarrow \Lambda$ form factors **at twist-four** accuracy [YMW, Shen, 2016].
 - ▶ Incomplete NLO calculation of the jet function [Feldmann, Yip, 2015].
 - ▶ **Confirmed by the lattice QCD calculations** [Detmold, Meinel, 2016].
 - ▶ SCET factorization at leading power in Λ/m_b [Wei Wang, 2011].

The charmless hadronic B -meson decays


- **QCD/SCET factorization** [BBNS, BPRS, Chay, Kim, and many others].

Heavy quark limit: $m_b \gg \Lambda_{\text{QCD}}$


Large-energy limit: $E_M \approx m_b/2 \gg \Lambda_{\text{QCD}}$

Scales: $m_b, \sqrt{m_b \Lambda_{\text{QCD}}}, \Lambda_{\text{QCD}}, (M_{\text{EW}}, \Lambda_{\text{NP}})$

$$\mu \approx m_b$$



$$\mu \approx 1 \text{ GeV}$$



- Reduces $\langle M_1 M_2 | \mathcal{O} | B \rangle$ to simpler $\langle M | \mathcal{O} | B \rangle$ (form factors), $\langle 0 | \mathcal{O} | B \rangle$, $\langle M | \mathcal{O} | 0 \rangle$ (decay constants and distribution amplitudes).
- Calculation from first principles, but limited accuracy by Λ_{QCD}/m_b corrections.

Status of the NNLO QCD calculations

$$\langle M_1 M_2 | C_i O_i | \bar{B} \rangle_{\mathcal{L}_{\text{eff}}} = \sum_{\text{terms}} C(\mu_h) \times \left\{ F_{B \rightarrow M_1} \times \underbrace{T^I(\mu_h, \mu_s)}_{1+\alpha_s+\dots} \star f_{M_2} \Phi_{M_2}(\mu_s) \right. \\ \left. + f_B \Phi_B(\mu_s) \star \left[\underbrace{T^{II}(\mu_h, \mu_I)}_{1+\dots} \star \underbrace{J^{II}(\mu_I, \mu_s)}_{\alpha_s+\dots} \right] \star f_{M_1} \Phi_{M_1}(\mu_s) \star f_{M_2} \Phi_{M_2}(\mu_s) \right\}$$

	T^I , tree	T^I , penguin	T^{II} , tree	T^{II} , penguin
LO: $\mathcal{O}(1)$				
NLO: $\mathcal{O}(\alpha_s)$ BBNS '99-'04				
NNLO: $\mathcal{O}(\alpha_s^2)$	 Bell '07,'09 Beneke, Huber, Li '09 Huber, Krankl, Li '16	 Kim, Yoon '11, Bell Beneke, Huber, Li '15 Bell, Beneke, Huber, Li '20	 Beneke, Jager '05 Kivel '06, Pilipp '07	 Beneke, Jager '06 Jain, Rothstein, Stewart '07

- **Strong cancellation** of the NNLO corrections to **the topological tree amplitudes** between the vertex term and the spectator-scattering mechanism.
- **Strong cancellation** of the NNLO corrections to **the QCD penguin amplitudes** between the current-current operators $\mathcal{Q}_{1,2}^p$ and the penguin operators $\mathcal{Q}_{3-6,8,g}$.

PQCD factorization for hard exclusive reactions

- Pioneer works on hard exclusive reactions:
 - ▶ **Hard scattering approach** formulated in [Lepage, Brodsky, 1979, 1980].
 - ▶ The hadronic wave function in QCD [Brodsky, Lepage, Huang, 1980].
 - ▶ **Sudakov effects** in hadron-hadron elastic (**Landshoff scattering**) [Botts, Sterman, 1989].
 - ▶ Sudakov resummation for the pion electromagnetic form factor [Li, Sterman, 1992].
Key observation: Perturbative QCD formalism to hard exclusive processes applicable for $\sqrt{Q^2} \sim 20 \Lambda_{\text{QCD}}$.
Saving us from the strong doubts raised in [Isgur, Llewellyn Smith, 1988; Radyushkin, 1984].
 - ▶ More references can be found in the **Bible by John Collins**.

Important pieces of work accomplished in China:

- Applicability of PQCD factorization for the pion electromagnetic form factor [Huang, Shen, 1990].
- **Sudakov suppression** for hard exclusive reactions [PhD thesis by Jun Cao].
- Many more interesting papers to be discussed.

微扰量子色动力学应用到遍举 过程中的几个问题

曹 俊

指导教师姓名 : 黄 涛 吴慧芳
职称 : 研究员 研究员
单位 : 中国科学院高能物理所

PQCD factorization for heavy hadron decays

- Birth of the modern PQCD approach for charmless hadronic B -meson decays:
 - ▶ Penguin enhancement for $B \rightarrow \pi K$ [Keum, Li, Sanda, 2001].
 - ▶ Tree-dominated processes $B \rightarrow \pi\pi$ [Lü, Ukai, Yang, 2001].
- Hundreds of papers on the tree-level PQCD calculations [Groups led by Li, Lü, Xiao, etc].
- PQCD calculations at NLO complicated by the appearance of multi-scales:
 - ▶ The pion-photon form factor at large momentum transfer [Nandi, Li, 2007].
 - ▶ The pion electromagnetic form factor at large Q^2 [Li, Shen, Wang, Zou, 2011].
 - ▶ $B \rightarrow \pi$ form factors at large recoil [Li, Shen, Wang, 2012].
 - ▶ Increasing complexities mainly due to the infrared subtractions!
- Intensive NLO PQCD calculations subsequently:
 - ▶ NLO twist-3 correction to the pion electromagnetic form factor [Cheng, Fan, Xiao, 2014].
 - ▶ NLO twist-3 correction to $B \rightarrow \pi$ form factors [Cheng, Fan, Yu, Lü, Xiao, 2014].
 - ▶ NLO correction to $B \rightarrow \rho$ form factors [Hua, Zhang, Xiao, 2018].
 - ▶ (Partial)-NLO correction to $B \rightarrow \pi\pi$ [Cheng, Xiao, Zhang, 2014].
 - ▶ (Partial)-NLO correction to $B_s \rightarrow PP$ [Yan, Liu, Xiao, 2019].
 - ▶ (Partial)-NLO correction to $B_c \rightarrow J/\psi$ form factors [Liu, Li, Xiao, 2020].
 - ▶ Review on the NLO PQCD computations [Cheng, Xiao, 2020].
 - ▶ Many more papers on the NLO PQCD calculations.

Theoretical wishlist

- **Systematic understanding of the (high-twist) B -meson distribution amplitudes.**
 - ▶ Renormalization properties **beyond the one-loop** approximation [conformal symmetry].
 - ▶ Perturbative constraints at large ω_i [OPE technique].
 - ▶ **Renormalon analysis** and the renormalization-scheme dependence.
 - ▶ Precision determinations of the inverse moment λ_B .
- **QCD factorization for the subleading power corrections.**
 - ▶ SCET analysis for the pion-photon form factor as the first step [operator structures, symmetry constraints, etc].
 - ▶ General treatment of the **rapidity divergences** in the (naïve)-factorization formulae.
 - ▶ Rigorous factorization proof taking into account the **Glauber gluons**.
 - ▶ Novel resummation techniques for enhanced logarithms.
- **Technical issues for future improvements.**
 - ▶ Factorization techniques for **the electromagnetic corrections**.
 - ▶ NNLO QCD computations for $B \rightarrow V\gamma$ and $B \rightarrow V\ell\ell$.
 - ▶ QCD factorization for the radiative and electroweak penguin decays of the Λ_b -baryon.
 - ▶ Improved understanding of the parton-hadron **duality violation**.
- **Very promising future for QCD aspects of heavy-quark physics!**