

QPT 2021

Guiyang, China

Contribution ID: 7

Type: not specified

## Measurement of Intermittency for Charged Particles in Au + Au Collisions at $\sqrt{s_{NN}} = 7.7 - 200$ GeV from STAR

One of the main goals of RHIC beam energy scan (BES) program is to search for signatures of a critical point in the QCD phase diagram through heavy-ion collisions. It is predicted that the local density fluctuations near the critical point exhibit power-law scaling, which can be probed with an intermittency analysis of the scaled factorial moments,  $F_q(M)$ , for charged particles. The power-law behavior of  $q^{th}$  order scaled factorial moments can be expressed as:  $F_q(M) \sim F_2(M)^{\beta_q}$ , where  $M$  is the number of equally sized cells in one dimension of momentum space, and  $\beta_q$  is the intermittency exponent. The scaling exponent,  $\nu$ , related to the critical component can be derived from the equation:  $\beta_q = (q - 1)^\nu$ . The energy dependence of  $\nu$  could be used to search for the signature of the QCD critical point. Similar measurements have been carried out by NA49 and NA61 experiments in heavy-ion collisions with different system sizes.

In this talk, we will present the scaled factorial moments ( $F_q(M)$ , up to sixth order) of charged particles in Au + Au collisions at  $\sqrt{s_{NN}} = 7.7 - 200$  GeV measured by STAR experiment in the first phase of RHIC BES. Then, we will show the energy and centrality dependence of the extracted  $\nu$  values. The physical implications of these results will be discussed.

**Primary author:** Mr WU, Jin (CCNU)

**Presenter:** Mr WU, Jin (CCNU)