

Contribution ID: 64

Type: not specified

## Higher-order diagonal cumulants of net-Lambda multiplicity distributions and off-diagonal cumulants of net-proton, net-kaon, and net-charge multiplicity distributions in the STAR experiment in Au+Au collisions at $\sqrt{s_{NN}}$ =27 GeV

Susceptibilities of conserved quantities, such as net-charge, net-baryon, and net-strangeness, are sensitive to the quantum chromodynamics (QCD) phase transition, and also the QCD critical point. The prime aim of the RHIC beam-energy scan program is to explore the QCD phase diagram and search for the location of QCD critical , which can be studied by higher order cumulants of net-proton ( $\Delta$ p), netcharge ( $\Delta$ Q), and net-kaon ( $\Delta$ k) multiplicity distributions. Additionally,the cumulants of net- $\Lambda$ ( $\Delta\Lambda$ ) multiplicity distributions could provide an access to explore flavor-dependent chemical freeze-out parameters in the QCD phase diagram. In the year 2018, the STAR experiment has collected high statistics minimum bias data in Au+Au collisions at = 27 GeV. We present the higher-order diagonal cumulants (1st to 4th order) and their ratios of  $\Delta\Lambda$  multiplicity distributions are discussed. Furthermore, the off-diagonal cumulants between  $\Delta$ p- $\Delta$ k at this collision energy along with other published BES results will be discussed.

**Primary author:** LI, changfeng (SDU) **Presenter:** LI, changfeng (SDU)