



Contribution ID: 16

Type: not specified

## Splitting scales in reclustered large-radius jets in high-energy nuclear collisions.

Jet has been proposed as an excellent probe to jet quenching and the properties of Quark-gluon plasma. Recently, large radius jets using a radius parameter of  $R = 1.0$ , by re-clustering anti- $k_t$   $R = 0.2$  jets having transverse momentum  $p_T > 35$  GeV/c, are measured by ATLAS in Pb+Pb collisions at 5.02 TeV. By re-clustering the large-radius jet constituents with  $k_t$  algorithm, its hardest parton splitting scale  $\sqrt{d_{12}}$  are measured. Those measurements provide additional constraints on the jet-medium interaction and jet transport coefficient.

In the talk, we carry out the first theoretical investigation on the medium modification of the reclustered large-radius jets production, in particular, the hardest parton splitting of these jets in Pb+Pb relative to that in p+p collisions. Jet propagation and jet-induced medium excitation in the hot-dense medium is investigated within the Linear Boltzmann Transport (LBT) model.

The nuclear modification factor of the large radius jets evaluated as a function of jet transverse momentum is a little smaller than the value of inclusive  $R = 0.4$  jets due to its complex structures. The large radius jet constituents are reclustered with the  $k_t$  algorithm to obtain the splitting scale  $\sqrt{d_{12}}$ , which characterizes the transverse momentum scale for the hardest splitting in the jet. The jet yields evaluated as a function of the splitting scale  $\sqrt{d_{12}}$  of the hardest parton splitting is overall suppressed in Pb+Pb collisions relative to p+p collisions due to the reduction of jets yields.

A significant evolution of  $R_{AA}$  with  $\sqrt{d_{12}}$  is observed at small  $\sqrt{d_{12}}$  values, indicating a significant difference in the quenching of large-radius jets having single sub-jet and those with more complex substructure. We find that large radius jets with small splitting angle and small fragmentation function are less suppressed. A detailed analysis shows that jet substructures are significantly modified due to the interactions between jet parton and therm medium constituents.

**Primary author:** Mr ZHANG, Shan-liang (SCNU)

**Co-authors:** Prof. ZHANG, Ben-Wei (Central China Normal University); YANG, Meng-Quan (CCNU)

**Presenter:** Mr ZHANG, Shan-liang (SCNU)