## The 14th Workshop on QCD Phase Transition and Relativistic Heavy-Ion Physics (QPT 2021)



Contribution ID: 18

Type: not specified

## Measurement of proton-Xi correlation function in Au+Au collisions at 200GeV at RHIC-STAR

The study of baryon-baryon interactions is important to understand existence of strangelets and various exotic hadrons, and for modeling of astronomical objects such as neutron stars. A detailed knowledge of nucleon-nucleon (NN) potentials exists in literature, however very little is known about interactions between anti-nucleons. Similarly lack of scattering data for hyperon-nucleon (YN) systems makes it difficult to construct YN potentials. In heavy-ion collisions, a large number of baryons are produced in each nucleus-nucleus collision, which allows us to study the NN and YN interactions. Measurements of two-particle correlation function are used to study the space-time dynamics of the source created in heavy-ion collisions. At low relative momentum, the two-particle correlations are effected by the Final State Interactions (FSI), making it possible to measure FSI between nucleon and multi-strange baryon  $\Xi$ . The first measurement of proton- $\Xi$  correlation function from the STAR experiment at RHIC for the Au+Au collisions at  $\sqrt{s_NN} = 200$  GeV will be presented.

**Primary author:** 米, 柯 (Central China Normal University) **Presenter:** 米, 柯 (Central China Normal University)