



Light Hyper-Nuclei Directed Flow in $\sqrt{s_{NN}} = 3$ GeV Au+Au Collisions at RHIC

胡晨露 (Chenlu Hu) for STAR Collaboration




Institute of Modern Physics, CAS

July 27, 2021




- 1) Motivation
- 2) STAR Detector System for Fixed-target runs
- 3) Results of Hyper-nuclei (${}^3_{\Lambda}\text{H}$ & ${}^4_{\Lambda}\text{H}$) from 3 GeV
Au+Au Collisions
- 4) Summary



- 1) Hyper-nucleus provides opportunity for studying hyperon-nucleon (YN) interactions. Important for understanding inner structure of compact stars
- 2) In high-energy heavy-ion collisions (HICs), hyper-nuclei are abundantly produced in high baryon density region. Ideal environment for studying properties of hyper-nuclei
 - Lifetime
 - Hyper-nuclei production mechanism in HICs
 - Others(decay branching ratio, binding energy, etc)


A. Andronic et al., Phys. Lett. **B697**, 203(2011)
 J. Steinheimer et al., Phys. Lett. **B714**, 85(2012)



RHIC Beam Energy BES-II in 2019-2021:




- Fixed Target Run extends collision energy down to : $\sqrt{s_{NN}} = 3 - 7.2 \text{ GeV}$ corresponding to chemical potential: $750 \geq \mu_B \geq 420 \text{ MeV}$

STAR TPC Particle Identification




2018 STAR FXT 3 GeV data set;
260M minimum biased events

- 1) PID of p , d , t , ${}^3\text{He}$, ${}^4\text{He}$, π^- are made based on the dE/dx vs p/q distribution and particles are selected by $\ln \sigma$ method;
- 2) Hyper-nuclei reconstruction channels:




Concept:

- Mother and daughter particles have the same state vector and are treated in the same way
- Reconstruction of decay chains
- **Kalman Filter (KF) based**
- Geometry independent and Vectorized



Simulated AuAu collision at 25 AGeV

S. Gorbunov and I. Kisel, Reconstruction of decayed particles based on the Kalman filter. CBM-SOFT-note-2007-003, 7 May 2007




KF Particle package shows a high quality of the reconstructed particles, high efficiencies, and high signal to background ratios


KF Particle Finder — M. Zyzak, "Online selection of short-lived particles on many-core computer architectures in the CBM experiment at FAIR," Dissertation thesis, Goethe University of Frankfurt, 2016, <http://publikationen.uni-frankfurt.de/frontdoor/index/index/docId/41428>

KF Particle provides a direct approach to physics analysis: ALICE, CBM, sPHENIX and STAR


$\Lambda \rightarrow p + \pi^-$




${}^3_\Lambda\text{H} \rightarrow p + d + \pi^-$



${}^3_\Lambda\text{H} \rightarrow \pi^- + {}^3\text{He}$




${}^4_\Lambda\text{H} \rightarrow \pi^- + {}^4\text{He}$




➤ KFParticle package used for Λ , ${}^3_\Lambda\text{H}$ and ${}^4_\Lambda\text{H}$ reconstructions

Phase space



Acceptance and Efficiency



$$\frac{d^2 N}{p_T dp_T d\varphi} = \frac{1}{2\pi} \frac{dN}{p_T dp_T} \left\{ 1 + \sum_{n=1}^{\infty} 2v_n(p_T) \cos[n(\varphi - \Psi_R)] \right\}$$

– v_1 Directed flow; – v_2 Elliptic flow ...

- 1) Fixed Target $\sqrt{s_{NN}} = 3$ GeV Au+Au collision;
 $y_{\text{target}} \approx -1.045$
- 2) Charged tracks measured by TPC used for centrality definition;



- 1st order event plane angle measured by Event Plane Detector (EPD)

- Event-plane resolution determination:


$$R_1 = \langle \cos(\Psi_1 - \Psi_r) \rangle = \frac{\sqrt{\pi}}{2\sqrt{2}} \chi_1 \exp\left(-\frac{\chi_1^2}{4}\right) \left[I_0\left(\frac{\chi_1^2}{4}\right) + I_1\left(\frac{\chi_1^2}{4}\right) \right]$$

$$R_2 = \langle \cos(2(\Psi_1 - \Psi_r)) \rangle = \frac{\sqrt{\pi}}{2\sqrt{2}} \chi_1 \exp\left(-\frac{\chi_1^2}{4}\right) \left[I_{\frac{1}{2}}\left(\frac{\chi_1^2}{4}\right) + I_{\frac{3}{2}}\left(\frac{\chi_1^2}{4}\right) \right]$$

- The event plane resolution is in the range of 40 – 75% for the mid-centrality region 5-40% 3 GeV Au+Au collisions




${}^4_{\Lambda}\text{H}$ p_T : (1.2, 3.0) GeV/c; y : (-1.0, 0.0); Centrality: 5-40%



p_1 is observed v_1


Fit function: $f(\phi - \psi_1) = p_0 * (1 + v_1 * 2 * \cos(\phi - \psi_1))$

$\sqrt{s_{NN}} = 3$ GeV Au+Au Collisions at RHIC



- 1) **First observation** of hyper-nuclei collectivity v_1 in high-energy nuclear collisions, EP resolution and efficiency corrections applied;
- 2) Like the cases for light nuclei, hyper-nuclei v_1 seems to follow the mass number scaling within uncertainties \rightarrow


Coalescence is a dominant process for mid-rapidity hyper-nuclei formation in the collisions



1) Within statistical uncertainties, the slopes of v_1 for hyper-nuclei ${}^3_{\Lambda}\text{H}$ and ${}^4_{\Lambda}\text{H}$ seem following the mass number scaling in the 5-40% 3 GeV Au+Au collisions;

→ Coalescence is a dominant process for hyper-nuclei formation in the collisions

- 1) Light hyper-nuclei ${}^3_{\Lambda}\text{H}$ and ${}^4_{\Lambda}\text{H}$ are reconstructed from 3 GeV Au+Au collisions at RHIC; Largest ${}^3_{\Lambda}\text{H}$ and ${}^4_{\Lambda}\text{H}$ data samples even being observed.
- 2) First measurements of ${}^3_{\Lambda}\text{H}$ and ${}^4_{\Lambda}\text{H}$ directed flow (v_1) from 5 – 40% centrality.
- 3) dv_1/dy slopes of hyper-nuclei ${}^3_{\Lambda}\text{H}$ and ${}^4_{\Lambda}\text{H}$ seem to flow a mass number scaling.
This result implies that Coalescence is a dominant process for hyper-nuclei formation in such collisions.



Thank you very much for your
attention!

Acknowledgements:

Yuri Fisyak, Ivan Kisel, Iouri Vassiliev, Maksym Zyzak