

# Axi-Higgs Cosmology

## Cosmology with a changing Higgs VEV and an Axionic Solution

Lingfeng Li (HKUST)





Based on [2102.11257] and [2105.01631]

w/ Leo WH Fung, Tao Liu, Hoang Nhan Luu, Yu-Cheng Qiu and S.-H. Henry Tye

Jul. 20, 2021, 15th Workshop on TeV Physics, Beijing

# Tension in $\Lambda$ CDM


- ▶ From CMB and local measurements:  $H_{0,\text{P18}} = 67.36 \pm 0.54$  km/s/Mpc [Aghanim et al., 2020] vs.  $H_{0,\text{late}} = 73.3 \pm 0.8$  km/s/Mpc [Verde et al., 2019] from  $z \lesssim 2$ .
- ▶ From Big Bang Nucleosynthesis (BBN): the abundance ratio  ${}^7\text{Li}/\text{H} \times 10^{10}$ :  $1.6 \pm 0.3$  (observed) vs.  $5.6 \pm 0.3$  (theoretical) [Zyla et al., 2020, Pitrou et al., 2018, Iliadis and Coc, 2020].
- ▶ The weak lensing measurement of  $S_8$  together with the clustering parameter  $\sigma_8$  [Troxel et al., 2018] yields a value smaller  $S_{8,\text{DES}} = 0.773^{+0.026}_{-0.020}$  than that given by the CMB- $\Lambda$ CDM value:  $S_{8,\text{CMB}} = 0.832 \pm 0.013$ .
- ▶ Recently measurement of isotropic cosmic birefringence (ICB) based on the cross-power (parity-violating)  $C_l^{EB}$  data in CMB [Minami and Komatsu, 2020], deviate from 0 by  $\sim 2.4\sigma$ .



[Verde et al., 2019, Pitrou et al., 2018, Handley and Lemos, 2019, Minami and Komatsu, 2020]

# Overview: Changing VEV by an Axion

Tension in  $H_0$  can be alleviated if  $m_e$  is  $\mathcal{O}(1\%)$  heavier when  $z \sim 1100$  (recombination) [Ade et al., 2015, Hart and Chluba, 2020]



- ▶ The Bohr radius  $\downarrow$  as  $m_e \uparrow$
- ▶ Earlier “freeze out” of recombination.
- ▶ During BBN,  $m_q \uparrow$  makes  ${}^7\text{Li} \downarrow$ .

A slowly changing field that moves  $v \propto m_e$ : the axion.

- ▶ Introduces a time scale  $\gtrsim \mathcal{O}(10^6)$  yrs,  $m_a \lesssim 3 \times 10^{-29}$  eV.
- ▶ A light scalar suppresses structure formation ( $\sigma_8$  tension resolved).
- ▶ The  $aF^{\mu\nu}\tilde{F}_{\mu\nu}$  coupling leading to non-zero ICB [Fujita et al., 2020]

# CMB: Basic Formula

The angular sound horizon  $\theta_* = \frac{r_*}{D_*}$  provides sensitivity to  $H_0$  :

$$r_* = \int_{z_*}^{\infty} dz \frac{c_s(z)}{H(z)} \propto \int_{z_*}^{\infty} dz \left/ \sqrt{3 \left[ 1 + \frac{3\omega_b}{4\omega_\gamma(1+z)} \right] [\omega_r(1+z)^4 + \omega_m(1+z)^3 + \omega_\Lambda]} \right. \quad (1)$$

$$D_* = \int_0^{z_*} dz \frac{1}{H(z)} \propto \int_0^{z_*} dz \left/ \sqrt{\omega_r(1+z)^4 + \omega_m(1+z)^3 + \omega_\Lambda} \right. , \quad (2)$$


## Earlier recombination $\Rightarrow$ higher $H_0$

- ▶ For  $r_*$ , major contribution comes from integration near  $z^*$ , earlier  $z_*$  reduces  $r^*$ .
- ▶  $D_*$  hardly feels the change as it is dominated by late time expansion.  
 $\Rightarrow \frac{\partial \log(r_*)}{\partial \log(z_*)} \simeq -0.66$  ,  $\frac{\partial \log(D_*)}{\partial \log(z_*)} \sim -10^{-2}$  .
- ▶ Larger cosmological constant is possible.

# CMB Preference

The two most important (so far) constraint that we decide to match

$$(r_d h)_{\text{BAO}} = (99.95 \pm 1.20) \text{ Mpc} , \quad (\theta_*)_{\text{CMB}} = (1.04110 \pm 0.00031) \times 10^{-2} .$$



Relative to Planck18 fit values:

- ▶  $H_0 \sim 71 \pm 1 \text{ km/s/Mpc}$  when  $\delta v \sim 4\%$  .
- ▶  $H_0 \sim 69 \pm 1 \text{ km/s/Mpc}$  when  $\delta v \sim 1\%$  .
- ▶  $\omega_c$  increases by 1 – 4% .

# Axion as the Modulator of VEV

The minimal coupling ( $\phi = \text{Higgs}$ ,  $a = \text{axion}$ ):


$$V(\phi, a) \sim V_{\text{SM}} = \mu^2 |\phi|^2 + \frac{\kappa}{4} |\phi|^4,$$

$$\mu^2 \mapsto \mu^2 \left(1 + \text{const} \times \frac{a^2}{M_{\text{PL}}^2}\right), \quad \kappa \mapsto \kappa \left(1 + \text{const}' \times \frac{a^2}{M_{\text{PL}}^2}\right),$$

$$W \supset X(m_s^2 G(A) - \kappa K(A) H_u H_d) \rightarrow V_\phi = |m_s^2 G(a) - \kappa K(a) \phi^\dagger \phi|^2.$$


“Misalignment” with  $m_a \lesssim 10^{-29} \text{eV}$ :

- ▶ Higgs always in the bottom.
- ▶  $a \rightarrow$  sub-component of DM.
- ▶  $x \equiv \rho(\text{axion DM})/\rho(\text{all matter})$ .
- ▶  $x \sim \mathcal{O}(1\%)$  when  $f_a \sim M_{\text{PL}}$ .




# $\sigma_8/S_8$ Tension Resolved

The dilemma of early recombination solution of  $H_0$  tension:  
more CDM to fit CMB  $\Rightarrow$  higher  $\sigma_8$  [Jedamzik et al., 2020]



$$\frac{\mathcal{P}_m^x(k)}{\mathcal{P}_m^0(k)} = \left( \frac{k_J(z_{\text{eq}})}{k} \right)^{10 - 2\sqrt{25 - 24x}} \quad (3)$$

The axion quantum pressure and its potential force suppress the structural formation smaller than the Jeans scale.  
[Marsh and Ferreira, 2010, Kobayashi et al., 2017].






# BBN: An Alternative VEV

Heuristically, the neutron to proton number ratio  $n/p$  can be written as [Hall et al., 2014]:

$$\frac{n}{p} \sim e^{-\frac{\Delta m_{np}}{T_{np}}} e^{-\frac{t_D}{\tau_n}}$$

- ▶ Fermi constant  $G_F \propto v^{-2}$  that affect weak interactions. Larger  $m_W \Rightarrow$  earlier  $n \leftrightarrow p$  freeze out & longer  $\tau_n$ .
- ▶ Electron mass  $m_e \propto v$ . Similar effect.
- ▶ Increasing the isospin-breaking  $\Delta m_q \equiv m_d - m_u \propto v$  contributes to larger  $\Delta m_{np} \equiv m_n - m_p \Rightarrow$  later  $n \leftrightarrow p$  freeze out & shorter  $\tau_n$ .
- ▶ Averaged light quark mass  $\bar{m}_q \equiv (m_u + m_d)/2 \propto v$ . The change of  $\bar{m}_q$  may significantly influence the rates of strong/nuclear interactions. Essentially changes  $m_\pi$ , **making nuclei unstable**.

# Compatibility with BBN and ${}^7\text{Li}$ Problem



Heavy nuclei more fragile as  $m_q \uparrow \Rightarrow m_\pi \uparrow$ ,  ${}^7\text{Li}$  production harder.

[Flambaum and Wiringa, 2007]

$\begin{matrix} \text{Y} \\ \backslash \\ \text{X} \end{matrix}$	$m_W$	$m_e$	$\Delta m_q$	$\bar{m}_q$	$\eta$
$Y_p$	2.9	0.40	-5.9	-1.0	0.039
$D/H$	1.6	0.59	-5.3	10	-1.6
${}^7\text{Li}/H$	1.7	-0.04	-5.3	-60	2.1

[Dent et al., 2007, Cheoun et al., 2011,

Mori and Kusakabe, 2019]

►  $\delta v \equiv (v_{\text{BBN}} - v_0)/v_0 \simeq 1\%$

►  $\delta\eta \simeq \delta\omega_b \simeq 1 - 3\%$

Overlaps w/ CMB preference!

# Isotropic Cosmic Birefringence

If the axion has a coupling with the EM gauge field as

$$\frac{1}{32\pi^2} \frac{a}{f_a} F \tilde{F} ,$$

the initial phase of axion breaks parity and rotate the linearly polarized CMB [Harari and Sikivie, 1992]:

$$\beta = \frac{1}{16\pi^2 f_a} \int_{t_{\text{ini}}}^{t_0} dt \dot{a} = \frac{1}{16\pi^2 f_a} \left[ a(t_0) - a(t_{\text{ini}}) \right] . \quad (4)$$


$$\beta \sim -\frac{1}{16\pi^2} \frac{a_{\text{ini}}}{f_a} = 0.35 \pm 0.14(\text{degree}), \quad \Rightarrow \quad \frac{a_{\text{ini}}}{f_a} \simeq 1.0 \pm 0.3 . \quad (5)$$

[Minami and Komatsu, 2020]


# Observational Limits: Atomic Clocks

$d(\delta\nu)/dt$  limit from atomic clocks [Lange et al., 2021]:

$$\left. \frac{d(\delta\nu)}{dt} \right|_{t_0} \simeq \left. \frac{d(\delta\mu)}{dt} \right|_{t_0} = - (0.08 \pm 0.36) \times 10^{-16} \text{ yr}^{-1}$$





$$m_a \gtrsim 2 \times 10^{-30} \text{ eV}$$







$\mathcal{O}(10^{-18}) \text{ yr}^{-1}$  precision is needed  $\Rightarrow$  future improvements of atomic (nuclear) clocks.






Discovery potential within 1-2 decades.

# Summary



- ▶ A higher Higgs VEV when  $z \gtrsim 1100$ :
  - ▶ Earlier recombination, larger  $H_0$ .
  - ▶ Compatible with BBN (and  ${}^7\text{Li}$ )!
- ▶ An axionic solution:
  - ▶ Alleviate tension in  $\sigma_8/S_8$ , even with more DM.
  - ▶ Naturally explaining the non-zero ICB.
  - ▶ Great accessibility by different observations.

-  Ade, P. A. R. et al. (2015).  
Planck intermediate results - XXIV. Constraints on variations in fundamental constants.  
*Astron. Astrophys.*, 580:A22.
-  Aghanim, N. et al. (2020).  
Planck 2018 results. VI. Cosmological parameters.  
*Astron. Astrophys.*, 641:A6.
-  Ali-Haïmoud, Y. and Hirata, C. M. (2010).  
Ultrafast effective multilevel atom method for primordial hydrogen recombination.  
*Physical Review D*, 82(6).
-  Cheoun, M.-K., Kajino, T., Kusakabe, M., and Mathews, G. J. (2011).  
Time Dependent Quark Masses and Big Bang Nucleosynthesis Revisited.  
*Phys. Rev. D*, 84:043001.

-  Dent, T., Stern, S., and Wetterich, C. (2007).  
Primordial nucleosynthesis as a probe of fundamental physics parameters.  
*Phys. Rev. D*, 76:063513.
-  Flambaum, V. and Wiringa, R. B. (2007).  
Dependence of nuclear binding on hadronic mass variation.  
*Phys. Rev. C*, 76:054002.
-  Fujita, T., Murai, K., Nakatsuka, H., and Tsujikawa, S. (2020).  
Detection of isotropic cosmic birefringence and its implications for axion-like particles including dark energy.
-  Hall, L. J., Pinner, D., and Ruderman, J. T. (2014).  
The Weak Scale from BBN.  
*JHEP*, 12:134.
-  Handley, W. and Lemos, P. (2019).

Quantifying tensions in cosmological parameters:  
Interpreting the DES evidence ratio.

*Phys. Rev. D*, 100(4):043504.



Harari, D. and Sikivie, P. (1992).

Effects of a Nambu-Goldstone boson on the polarization of radio galaxies and the cosmic microwave background.

*Phys. Lett. B*, 289:67–72.



Hart, L. and Chluba, J. (2020).

Updated fundamental constant constraints from Planck 2018 data and possible relations to the Hubble tension.

*Mon. Not. Roy. Astron. Soc.*, 493(3):3255–3263.



Iliadis, C. and Coc, A. (2020).

Thermonuclear reaction rates and primordial nucleosynthesis.

*Astrophys. J.*, 901(2):127.




Jedamzik, K., Pogosian, L., and Zhao, G.-B. (2020).




Why reducing the cosmic sound horizon can not fully resolve the Hubble tension.

 Kobayashi, T., Murgia, R., De Simone, A., Iršič, V., and Viel, M. (2017).





Lyman- $\alpha$  constraints on ultralight scalar dark matter:  
Implications for the early and late universe.  
*Phys. Rev. D*, 96(12):123514.

 Lange, R., Huntemann, N., Rahm, J. M., Sanner, C., Shao, H., Lipphardt, B., Tamm, C., Weyers, S., and Peik, E. (2021).

Improved limits for violations of local position invariance  
from atomic clock comparisons.  
*Phys. Rev. Lett.*, 126(1):011102.

 Marsh, D. J. E. and Ferreira, P. G. (2010).  
Ultra-Light Scalar Fields and the Growth of Structure in the  
Universe.

*Phys. Rev. D*, 82:103528.

-  Minami, Y. and Komatsu, E. (2020).  
New Extraction of the Cosmic Birefringence from the Planck 2018 Polarization Data.  
*Phys. Rev. Lett.*, 125(22):221301.
-  Mori, K. and Kusakabe, M. (2019).  
Roles of  ${}^7\text{Be}(n, p){}^7\text{Li}$  resonances in big bang nucleosynthesis with time-dependent quark mass and Li reduction by a heavy quark mass.  
*Phys. Rev. D*, 99(8):083013.
-  Pitrou, C., Coc, A., Uzan, J.-P., and Vangioni, E. (2018).  
Precision big bang nucleosynthesis with improved Helium-4 predictions.  
*Phys. Rept.*, 754:1–66.
-  Troxel, M. A. et al. (2018).

Dark Energy Survey Year 1 results: Cosmological constraints from cosmic shear.

*Phys. Rev. D*, 98(4):043528.



Verde, L., Treu, T., and Riess, A. (2019).

Tensions between the Early and the Late Universe.

*Nature Astron.*, 3:891.



Zyla, P. et al. (2020).

Review of Particle Physics.

*PTEP*, 2020(8):083C01.