

# **Higgs potential and BSM opportunity 希格斯物理研讨会**

Friday, August 27, 2021 - Tuesday, August 31, 2021

南京大学（鼓楼校区）线上会议 Nanjing University (Gulou Campus)  
at ZOOM

## **Book of Abstracts**



# Contents

Opening . . . . .	1
TBD . . . . .	1
TBD . . . . .	1
Probing electroweak phase transition with multi-TeV muon colliders and gravitational waves . . . . .	1
Selected results in the 3HDMs . . . . .	1
Probing the electroweak symmetry breaking with Higgs production at the LHC . . . . .	2
Affleck-Dine Leptogenesis from Higgs Inflation . . . . .	2
Measurement of the CP property of the Higgs to top Yukawa coupling . . . . .	2
Probing the B+L violation process with the observation of cosmic magnetic field . . . . .	3
Search for heavy Higgs decaying to two tau leptons with the ATLAS detector using pp collision at $\sqrt{s} = 13$ TeV . . . . .	3
Probing Higgs boson properties with boosted objects at the CMS experiment . . . . .	3
Search for Resonant and Non-resonant VHH Production with ATLAS detector . . . . .	4
Probe BSM physics with the interaction between the Higgs boson and top quark . . . . .	4
All order nonfactorizable jet veto effects in Higgs boson production . . . . .	4
Search for Higgs boson pair production in $\tau\tau$ final state . . . . .	5
Primordial black holes from a cosmic phase transition: The collapse of Fermi-balls . . . . .	5
Search for doubly and singly charged Higgs bosons decaying into vector bosons in multi- lepton final states with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV . . . . .	6
New Physics bounded from above: 2HDM confronting strong 1st order electroweak phase transition . . . . .	6
电弱相变中引力波信号的精确计算 . . . . .	6
Measurements of Higgs boson production using decays to two c-quarks with the ATLAS detector . . . . .	7

Combined measurements of Higgs boson production and decay and constraints on the Higgs boson self-coupling from the combination of single-Higgs and double-Higgs production analyses performed with the ATLAS experiment . . . . .	7
Higgs Alignment and Novel CP-Violating Observables in 2HDM . . . . .	8
Measurements of Higgs boson decaying into tau leptons using 139 fb at the ATLAS experiment . . . . .	8
Next-to-leading order corrections for $gg \rightarrow ZH$ with top quark mass dependence . . . . .	8
Fiducial and differential cross-section measurements in the di-photon channel using full Run2 dataset at ATLAS . . . . .	9
Search for Higgs boson pair production in the $b\bar{b}\tau\tau$ decay channel using 13 TeV pp collision data from the ATLAS detector . . . . .	9
Boosted technique for Higgs measurement and New Physics Searches . . . . .	10
Measuring Higgs Boson Self-couplings with $2 \rightarrow 3$ VBS Processes . . . . .	10
Effective picture of cosmic bubble expansion . . . . .	10
Long-lived Particle(LLP) searching at the future collider . . . . .	11
Higgs Alignment and Novel CP-Violating Observables in 2HDM . . . . .	11
Selected results in the 3HDMs . . . . .	11
Boosted technique for Higgs measurement and New Physics Searches . . . . .	11
Affleck-Dine Leptogenesis from Higgs Inflation . . . . .	11
Search for doubly and singly charged Higgs bosons decaying into vector bosons in multi-lepton final states with the ATLAS detector using proton-proton collisions at 13 TeV . . . . .	12
New Physics bounded from above: 2HDM confronting strong 1st order electroweak phase transition . . . . .	12
Search for heavy Higgs decaying to two tau leptons with the ATLAS detector using pp collision at 13 TeV . . . . .	12
Search for Resonant and Non-resonant VHH Production with ATLAS detector . . . . .	12
Probe BSM physics with the interaction between the Higgs boson and top quark . . . . .	12
Searching for Long-lived Particles at future lepton collider . . . . .	12
Composite Higgs and Dark Matter . . . . .	13
Higgs potential and phase transition . . . . .	13
Non-resonant di-Higgs process at LHC . . . . .	13
BSM resonance search for HH (SH, SS) at LHC . . . . .	13
Lunch time . . . . .	13

Higgs Overview . . . . .	13
Precise theoretical predictions for Higgs processes . . . . .	13
Precise Higgs measurements at LHC . . . . .	13
Search for rare Higgs processes . . . . .	14
Introduction to the standard model and electro-weak symmetry breaking . . . . .	14
Extension of standard model, e.g. 2HDM . . . . .	14
Experimental Higgs physics, e.g. discovery, precise measurement, BSM Higgs searches . . . . .	14
The multilepton anomalies at the LHC and the prospect of new physics at the EW scale . . . . .	14
Searches for light scalar and charged scalar . . . . .	14
Heavy neutral scalar searches . . . . .	15
BSM Higgs couplings . . . . .	15
Higgs bosons in new physics models . . . . .	15
BSM Higgs searches in final states with top quarks . . . . .	15
Opening . . . . .	15
Speech from NJU . . . . .	15
Physics Highlights at CEPC . . . . .	15
会议通知 . . . . .	16
Openning . . . . .	16
Welcome from NJU . . . . .	16
School opening . . . . .	16
Close . . . . .	16



36

## Opening

37

TBD

38

TBD

Di-Higgs/Scalar / 39

## Probing electroweak phase transition with multi-TeV muon colliders and gravitational waves

**Authors:** Ke-pan Xie<sup>1</sup>; Wei Liu<sup>2</sup>

<sup>1</sup> *Seoul National University*

<sup>2</sup> *Nanjing University of Science and Technology*

**Corresponding Author:** wei.liu@njust.edu.cn

We study the complementarity of the proposed multi-TeV muon colliders and the near-future gravitational wave (GW) detectors to the first order electroweak phase transition (FOEWPT), taking the real scalar extended Standard Model as the representative model. A detailed collider simulation shows the FOEWPT parameter space can be greatly probed via the the vector boson fusion production of the singlet, and its subsequent decay to the di-Higgs or di-boson channels. Especially, almost all the parameter space yielding detectable GW signals can be probed by the muon colliders. Therefore, if we could detect stochastic GWs in the future, a muon collider could provide a hopeful crosscheck to identify their origin. On the other hand, there is considerable parameter space that escapes GW detections but is within the reach of the muon colliders. The precision measurements of Higgs couplings could also probe the FOEWPT parameter space efficiently.

BSM Higgs sector / 40

## Selected results in the 3HDMs

**Author:** Igor Ivanov<sup>1</sup>

<sup>1</sup> *SYSU, School of Physics and Astronomy*

**Corresponding Author:** ivanov@mail.sysu.edu.cn

New Physics can manifest itself at colliders in the form of a non-minimal Higgs sector. In particular, the Higgs doublets can also come in generations leading to rich phenomenology. In this talk, I will give a brief overview of the three-Higgs-doublet models (3HDM) focusing on novel model-building opportunities, which go beyond the well studied 2HDM case. I will highlight some recent results related to novel symmetries and their consequences.

**Summary:**

New Physics can manifest itself at colliders in the form of a non-minimal Higgs sector. In particular, the Higgs doublets can also come in generations leading to rich phenomenology. In this talk, I will give a brief overview of the three-Higgs-doublet models (3HDM) focusing on novel model-building opportunities, which go beyond the well studied 2HDM case. I will highlight some recent results related to novel symmetries and their consequences.

**Higgs precise measurement / 41**

## Probing the electroweak symmetry breaking with Higgs production at the LHC

**Author:** Bin Yan<sup>1</sup>

**Co-author:** Ke-Pan Xie<sup>2</sup>

<sup>1</sup> LANL

<sup>2</sup> Seoul National University

**Corresponding Author:** binyan@lanl.gov

The electroweak symmetry breaking (EWSB) mechanism is still an undecided question in particle physics. We propose to utilize the single top quark and Higgs associated production ( $th$ ),  $Zh$  production via gluon fusion at the LHC to probe the couplings between the Higgs and the gauge bosons and further to test the EWSB. We demonstrate that the  $th$  and  $gg \rightarrow Zh$  productions are sensitive to the relative sign of couplings ( $ht\bar{t}$ ,  $hWW$ ) and ( $ht\bar{t}$ ,  $hZZ$ ), respectively. We find that the relative sign between  $hWW$  and  $hZZ$  couplings could be fully determined after combining the present measurements from  $gg \rightarrow h$ ,  $t\bar{t}h$  and the  $th$ ,  $Zh$  channels at the 13 TeV LHC, and this conclusion is not sensitive to the possible new physics contribution induced by  $Zt\bar{t}$  couplings in the  $gg \rightarrow Zh$  production.

**BSM Higgs sector / 42**

## Affleck-Dine Leptogenesis from Higgs Inflation

**Author:** Chengcheng Han<sup>1</sup>

<sup>1</sup> Sun Yat-sen University

I will present a simple extension of the Standard model to explain the origin of inflation, baryon asymmetry and neutrino masses at the same time.

**Higgs precise measurement / 43**



## Measurement of the CP property of the Higgs to top Yukawa coupling

**Author:** 洪洮杨<sup>1</sup>

<sup>1</sup> *Lawrence Berkeley National Laboratory*

**Corresponding Author:** hongtao.yang@cern.ch

Di-Higgs/Scalar / 45

## Probing the B+L violation process with the observation of cosmic magnetic field

**Author:** Ligong Bian<sup>1</sup>

<sup>1</sup> *Chongqing University*

**Corresponding Author:** lgbycl@cqu.edu.cn

We numerically investigate the B+L violation process by performing three-dimensional lattice simulations of a unified scenario of first-order phase transitions and the sphaleron generation. The simulation results indicate that the Chern-Simons number changes along with the helical magnetic field production when the sphaleron decay occurs. Based on these numerical results, we then propose a novel method to probe the baryon asymmetry generation of the Universe, which is a general consequence of the electroweak sphaleron process, through the astronomical observation of corresponding helical magnetic fields.

BSM Higgs sector / 46

## Search for heavy Higgs decaying to two tau leptons with the ATLAS detector using pp collision at $\sqrt{s} = 13$ TeV

**Author:** 瀚非叶<sup>1</sup>

**Co-authors:** Lei Zhang<sup>1</sup>; Xiaozhong Huang<sup>2</sup>

<sup>1</sup> *Nanjing University*

<sup>2</sup> *NJU*

This talk will cover the latest result of searching for heavy Higgs decaying to two tau leptons with the ATLAS full Run2 datasets using pp collision at  $\sqrt{s} = 13$  TeV. This search is performed over the mass range of 0.2-2.5 TeV and the two  $\tau$  leptons decay is considered, where at least one  $\tau$  decays leptonically. This channel is sensitive at high  $\tan\beta$  value according to hMSSM and other 2HDMs, where  $\tan\beta$  is the ratio of the vacuum expectation values of the two Higgs. The results of this search show that most of the parameter space of hMSSM is excluded with 95% CL.

Di-Higgs/Scalar / 47

## Probing Higgs boson properties with boosted objects at the CMS experiment

**Author:** Huilin Qu<sup>1</sup>

<sup>1</sup> *CERN*

**Corresponding Author:** hlquphy@gmail.com

Advances in boosted jet tagging techniques, particularly with novel machine learning approaches, have led to significant performance improvements in the reconstruction and identification of the hadronic decays of the Higgs boson and provided additional handles to probe the Higgs boson properties at the LHC. In this talk, I will present recent progress in machine learning-based methods for boosted Higgs boson reconstruction and discuss their applications in recent CMS searches, e.g., the  $H \rightarrow cc$  decay and the VBF Higgs pair production.

**BSM Higgs sector / 48**

## **Search for Resonant and Non-resonant VHH Production with ATLAS detector**

**Author:** Zhongyukun XU<sup>1</sup>

<sup>1</sup> *Shandong university*

**Corresponding Author:** zhongyukun.xu@cern.ch

A search for a vector boson produced in addition to two SM Higgs bosons is studied. Three different models are considered either non-resonant production with SM-like or non-SM-like Higgs self-coupling and  $hhVV$  coupling, or resonant production in the form of either  $VH(hh)$  or  $A \rightarrow ZH(hh)$ . Three sub-channel is considered for different decay mode for vector boson to  $Zll/Wlv/Zvv$  final states. Only  $HH \rightarrow bbbb$  decay is considered for statistics and high signal efficiency.

**Higgs precise measurement / 49**

## **Probe BSM physics with the interaction between the Higgs boson and top quark**

**Author:** Hualin Mei<sup>1</sup>

<sup>1</sup> *U*

**Corresponding Author:** hualin.mei@cern.ch

The top quark is the heaviest elementary particle, studying its interaction with the Higgs boson not only helps to further understand the newly discovered Higgs boson itself, but also opens new opportunities to probe the BSM physics. This talk will summarize recent new experimental results that connect the top-Higgs interaction with searches of new physics beyond the standard model with the LHC Run 2 dataset.

**Higgs precise measurement / 50**

## **All order nonfactorizable jet veto effects in Higgs boson production**

**Author:** Dingyu Shao<sup>1</sup>

<sup>1</sup> *Fudan University*

**Corresponding Author:** shaodinyu@gmail.com

I will discuss nonfactorizable QCD corrections from central jet veto to Higgs boson production at the Large Hadron Collider. At hadron colliders, phase factors in the scattering amplitudes lead to double-logarithmic corrections starting at four-loop order. I will introduce our recent work on the first all-order structure of these “super-leading” logarithms. Our analysis shows that in certain kinematic distributions the nonfactorizable corrections can be as large as a percent making them quite comparable to their factorizable counterparts.

**Di-Higgs/Scalar / 51**

## Search for Higgs boson pair production in $\gamma\gamma$ final state

**Author:** 子航贾 <sup>1</sup>

<sup>1</sup> *Nanjing University*

**Corresponding Author:** zihang.jia@cern.ch

Searches are performed for non-resonant and resonant di-Higgs boson production in the  $\gamma\gamma$  final state. The data set used corresponds to an integrated luminosity of 139 fb<sup>-1</sup> of proton–proton collisions at a center-of-mass energy of 13 TeV recorded by the ATLAS detector at the CERN Large Hadron Collider. No excess with respect to background expectations is found and upper limits on the di-Higgs boson production cross sections are set. A 95% confidence level upper limit of 130 fb is set on the  $\gamma\gamma \rightarrow \gamma\gamma$  non-resonant production, where the expected limit is 180 fb. The observed (expected) limit corresponds to 4.1 (5.5) times the cross section predicted by the Standard Model. The observed (expected) limit on the Higgs boson trilinear coupling modifier  $\kappa_\lambda$  is extracted to be [-1.5, 6.7] ([-2.4, 7.7]) at 95% confidence level. The constraints on  $\kappa_\lambda$  are obtained over an expected background hypothesis excluding  $\gamma\gamma \rightarrow \gamma\gamma$  production. For the resonant production of a new hypothetical scalar particle  $\phi$  ( $\phi \rightarrow \gamma\gamma \rightarrow \gamma\gamma\gamma$ ), limits on the cross section  $\phi \rightarrow \gamma \rightarrow \gamma\gamma$  are presented for the narrow-width approximation as a function of  $m_\phi$  in the range 251 GeV  $\leq m_\phi \leq$  1000 GeV. The observed (expected) limits on the cross section  $\phi \rightarrow \gamma \rightarrow \gamma\gamma$  range from 610 fb to 47 fb (360–43 fb) over the considered mass range.

**Di-Higgs/Scalar / 52**

## Primordial black holes from a cosmic phase transition: The collapse of Fermi-balls

**Author:** 柯盼谢 <sup>1</sup>

**Co-author:** Kiyoharu Kawana <sup>1</sup>

<sup>1</sup> *Seoul National University*

We propose a novel primordial black hole (PBH) formation mechanism based on a first-order phase transition (FOPT). If a fermion species gains a huge mass in the true vacuum, the corresponding particles get trapped in the false vacuum as they do not have sufficient energy to penetrate the bubble wall. After the FOPT, the fermions are compressed into the false vacuum remnants to form non-topological solitons called Fermi-balls, and then collapse to PBHs due to the Yukawa attractive force. We derive the PBH mass and abundance, showing that for a  $\mathcal{O}(\text{GeV})$  FOPT the PBHs could be  $\sim 10^{17}$  g and explain all of dark matter. If the FOPT happens at higher scale, PBHs are typically overproduced and extra dilution mechanism is necessary to satisfy current constraints.

**BSM Higgs sector / 53****Search for doubly and singly charged Higgs bosons decaying into vector bosons in multi-lepton final states with the ATLAS detector using proton–proton collisions at  $\sqrt{s} = 13$  TeV****Authors:** Yanwen Liu<sup>1</sup>; hanlin Xu<sup>2</sup><sup>1</sup> *USTC*<sup>2</sup> *University of Science and Technology of China***Corresponding Author:** hanlinxuy@gmail.com

A search for charged Higgs bosons decaying into  $W^\pm W^\pm$  or  $W^\pm Z$  bosons is performed, involving experimental signatures with two leptons of the same charge, or three or four leptons with a variety of charge combinations, missing transverse momentum and jets.

A data sample of proton–proton collisions at a centre-of-mass energy of 13 TeV recorded with the ATLAS detector at the Large Hadron Collider between 2015 and 2018 is used.

The data correspond to a total integrated luminosity of  $139 \text{ fb}^{-1}$ .

The search is guided by a type-II seesaw model that extends the scalar sector of the Standard Model with a scalar triplet,

leading to a phenomenology that includes doubly and singly charged Higgs bosons.

Two scenarios are explored, corresponding to the pair production of doubly charged  $H^{\pm\pm}$  bosons, or the associated production of a doubly charged  $H^{\pm\pm}$  boson and a singly charged  $H^\pm$  boson.

No significant deviations from the Standard Model predictions are observed.

$H^{\pm\pm}$  bosons are excluded at 95% confidence level up to 350 GeV and 230 GeV for the pair and associated production modes, respectively.

**Di-Higgs/Scalar / 54****New Physics bounded from above: 2HDM confronting strong 1st order electroweak phase transition****Author:** mengchao Zhang<sup>1</sup><sup>1</sup> *itp*

I will discuss the general difficulty in New Physics search, and how to put an upper limit on New Physics scale (which can be detected or excluded in the near future) via strong 1st order electroweak phase transition.

**Di-Higgs/Scalar / 55****电弱相变中引力波信号的精确计算****Author:** 潇王<sup>1</sup><sup>1</sup> *中山大学天琴中心***Corresponding Author:** wangxiao2016@ihep.ac.cn

The energy budget of electroweak first-order phase transition is essential for the gravitational wave spectra. Most of the previous studies are based on the bag model with same sound velocity in the symmetric and broken phase. We study the energy budget and the corresponding gravitational

wave spectra beyond the bag model, where the sound velocities could be different in the symmetric and broken phase. Taking the Higgs sextic effective model as a representative model, we calculate the sound velocities in different phase, the gravitational wave spectra and the signal-to-noise ratio for different combinations of phase transition parameters beyond the bag model. We compare these new results with the ones obtained from the bag model. The proper sound velocities and phase transition parameters at the appropriate temperature are important to obtain more precise predictions.

**Higgs precise measurement / 56**

## **Measurements of Higgs boson production using decays to two c-quarks with the ATLAS detector**

**Author:** Tao Wang<sup>1</sup>

<sup>1</sup> *USTC*

**Corresponding Author:** tao.wang@mail.ustc.edu.cn

Testing the couplings of the Higgs boson to quarks is important to understand the origin of quark masses. The next most promising candidate is the decay to a pair of charm quark and antiquark.

The presentation will focus on the analysis of the associated production of the Higgs boson with a W or Z boson performed by the ATLAS Collaboration using data collected between 2015 and 2018, and will describe the analysis strategy employed to search for the  $H \rightarrow cc$  signal. As a result, the best limit set on Higgs to charm quark Yukawa coupling modifier ( $\kappa_C$ ) to date will be shown.

**Higgs precise measurement / 57**

## **Combined measurements of Higgs boson production and decay and constraints on the Higgs boson self-coupling from the combination of single-Higgs and double-Higgs production analyses performed with the ATLAS experiment**

**Author:** Kunlin RAN<sup>1</sup>

<sup>1</sup> *Beijing*

A combination of measurements of Higgs boson production cross sections and branching fractions is presented. The combination is based on the analyses of the Higgs boson decay modes  $H \rightarrow \gamma\gamma$ ,  $ZZ$ ,  $WW$ ,  $\tau\tau$ ,  $bb$ ,  $\mu\mu$ , and searches for decays into invisible final states. Up to 139 fb<sup>-1</sup> of pp collision data collected at  $\sqrt{s}=13$  TeV with the ATLAS detector are used. Combined cross section measurements are presented for the ggF, VBF, WH, ZH and ttH. The global signal strength, defined as the measured Higgs boson signal yield normalized to its SM prediction, is determined to be  $\mu=1.06 \pm 0.07$ . The combined measurement yields an observed (expected) significance for the WH of 6.3 $\sigma$  (5.2 $\sigma$ ). Measurements in kinematic regions defined within the simplified template cross section framework are also shown. The results are interpreted in terms of modifiers applied to the Standard Model couplings of the Higgs boson to other particles. No significant deviations from SM predictions are observed.

Constraints on the Higgs boson self-coupling are set by combining the single Higgs boson analyses targeting the  $\gamma\gamma$ ,  $ZZ$ ,  $WW$ ,  $\tau\tau$  and  $bb$  decay channels and the double Higgs boson analyses in the  $bbbb$ ,  $bb\tau\tau$  and  $bbyy$  decay channels. The data used in these analyses correspond to an integrated luminosity of up to 79.8 fb<sup>-1</sup> for single Higgs boson analyses and up to 36.1 fb<sup>-1</sup> for the double Higgs boson analyses. With the assumption that new physics affects only the Higgs boson

self-coupling ( $\lambda_{\text{HHH}}$ ), values outside the interval  $[-2.3, 10.3]$  are excluded at 95% CL. Results with less stringent assumptions are also provided, introducing additional coupling modifiers for the Higgs boson interactions with the other SM particles.

**BSM Higgs sector / 58**

## Higgs Alignment and Novel CP-Violating Observables in 2HDM

**Authors:** Ian Low<sup>1</sup>; Xiaoping Wang<sup>2</sup>

<sup>1</sup> Argonne National Laboratory/Northwestern University

<sup>2</sup> Beihang University

**Corresponding Author:** hcwangxiaoping@163.com

Null results from searches for new physics at the Large Hadron Collider (LHC) enforce the belief that new particles must be much heavier than the weak scale. We undertake a systematic study of the interplay between Higgs alignment and CP-violation in complex two-Higgs-doublet models (C2HDMs), which enables us to construct a CP-violating scenario where new Higgs bosons are close to the weak scale after including stringent constraints from the electric dipole moment and measurements at the LHC. In addition, we propose a smoking-gun signal of CP-violation in the Higgs-to-Higgs decay,  $(h_3 \rightarrow h_2 h_1 \rightarrow 3 h_1)$ , where  $h_3, h_2$ , and  $h_1$  are the heaviest, second heaviest and the SM-like neutral Higgs bosons, respectively. The mere presence of this decay channel is sufficient to establish CP-violation in C2HDMs. The final state with three 125 GeV Higgs bosons is distinct and provides a unique venue for new measurements at the LHC.

**Higgs precise measurement / 59**

## Measurements of Higgs boson decaying into tau leptons using 139 fb at the ATLAS experiment

**Author:** Antonio De Maria<sup>1</sup>

<sup>1</sup> Nanjing University

**Corresponding Author:** antonio.de.maria@cern.ch

The most recent results for the cross-section measurements of the Higgs boson decaying into tau leptons at the ATLAS experiment will be presented. These results are setting the base for other searches in the Higgs boson sector in the tau final state aiming to look for physics beyond the Standard Model. The analysis is performed using the 139 fb dataset recorded by the ATLAS experiment at LHC in the period 2015-2018. A description of the analysis strategy concerning the signal categorization as well as the background estimation will be given, followed by an overview and discussion of the main analysis results: the inclusive Higgs boson to tau cross-section, the cross-section measurement for each of the main Higgs boson production mode at the LHC and the measurements within the Simplified Template Cross Section (STXS).

**Higgs precise measurement / 60**

## Next-to-leading order corrections for $gg \rightarrow ZH$ with top quark mass dependence

**Author:** 国兴王<sup>1</sup>

**Co-authors:** Lilin Yang<sup>2</sup>; Yongqi XU<sup>1</sup>; xu xiaofeng<sup>3</sup>

<sup>1</sup> Peking University

<sup>2</sup> Zhejiang University

<sup>3</sup> school of physics Peking University

**Corresponding Author:** wangguoxing2015@pku.edu.cn

Recently, we present for the first time a calculation of the complete next-to-leading order corrections to the  $gg \rightarrow ZH$  process. We use the method of small mass expansion to tackle the most challenging two-loop virtual amplitude, in which the top quark mass dependence is retained throughout the calculations. We show that our method provides reliable numeric results in all kinematic regions, and present phenomenological predictions for the total and differential cross sections at the Large Hadron Collider and its future upgrades. Our results are necessary ingredients towards reducing the theoretical uncertainties of the  $pp \rightarrow ZH$  cross sections down to the percent-level, and provide important theoretical inputs for future precision experimental collider programs. In this talk, I will start with a brief introduction to the small mass expansion and then show our new results for the  $ZH$  production at the LHC.

**Higgs precise measurement / 61**

## **Fiducial and differential cross-section measurements in the di-photon channel using full Run2 dataset at ATLAS**

**Author:** Fábio Alves<sup>1</sup>

<sup>1</sup> Nanjing University

**Corresponding Author:** fabio.lucio@cern.ch

Since the discovery of a scalar particle with mass at 125 GeV in the experiments ATLAS and CMS at LHC, different measurements based on its properties have been performed and the observations nicely correspond to the Higgs boson predicted by the Standard Model of particle physics. Among these measurements, the fiducial and differential cross-section play an important role in the test of the SM predictions as well as in the probe for BSM physics contributions exploring a variety of physics observables. Given that these measurements are performed in a specific region of the phase space (fiducial region), the model dependence is reduced. In this talk, the latest results on the differential and fiducial cross-section of the Higgs boson decay in the di-photon channel with full Run2 dataset (139 fb<sup>-1</sup>) collected with ATLAS experiment will be discussed.

**Di-Higgs/Scalar / 63**

## **Search for Higgs boson pair production in the $bb\tau\tau$ decay channel using 13 TeV pp collision data from the ATLAS detector**

**Author:** 博文张<sup>1</sup>

<sup>1</sup> Nanjing University

**Corresponding Author:** bowen.zhang@cern.ch

A search for non-resonant and resonant Higgs boson pair production in  $bb\tau\tau$  final states is conducted using p-p collision data with an integrated luminosity of 139 fb<sup>-1</sup> collected at  $s=\sqrt{13}$  TeV by the ATLAS detector at the LHC. Events with at least one  $\tau$ -lepton decaying hadronically are considered.

Multivariate analysis techniques are used to extract the signals. The talk will show an overview of the physics motivation, the analysis methodology as well as the preliminary results.

**BSM Higgs sector / 64**

## Boosted technique for Higgs measurement and New Physics Searches

**Author:** Qiang Li<sup>1</sup>

<sup>1</sup> *Peking University*

**Corresponding Author:** qliphy0@pku.edu.cn

Highly Lorentz-boosted scalar particles (Higgs or Radion) decaying to  $bb$  or  $WW \rightarrow 4q$  can be reconstructed as single large-radius jets, and identified using jet substructure and deep neural networks. The techniques can be exploited for both the SM Higgs measurement and New Physics Searches. Two recent applications at the CMS experiments will be covered in this talk, including, 1) a search for standard model Higgs bosons produced with transverse momentum ( $p_T$ ) greater than 450 GeV and decaying to bottom quark-antiquark pairs, and 2) a first search for resonances decaying into a radion and a W boson in proton-proton collision data, with the radion reconstructed through its decay into two W bosons. In the future, we expect to extend these techniques to boosted  $H \rightarrow WW$  and Higgs pair studies.

Refs:

<https://arxiv.org/abs/2006.13251>

CMS-PAS-B2G-20-001

CMS-PAS-B2G-21-002

**Di-Higgs/Scalar / 65**

## Measuring Higgs Boson Self-couplings with $2 \rightarrow 3$ VBS Processes

**Author:** Junmou (俊谋) Chen (谌)<sup>1</sup>

<sup>1</sup> *KIAS*

**Corresponding Author:** rainbowend@kias.re.kr

We study the measurement of Higgs boson self-couplings through  $2 \rightarrow 3$  vector boson scattering (VBS) processes in the framework of Standard Model effective field theory (SMEFT) at both proton and lepton colliders. The SMEFT contribution to the amplitude of the  $2 \rightarrow 3$  VBS processes, taking  $WLWL \rightarrow WLWLh$  and  $WLWL \rightarrow hhh$  as examples, exhibits enhancement with the energy  $A(\text{BSM})/A(\text{SM}) \sim$  which indicates the sensitivity of these processes to the related dimension-six operators in SMEFT. Simulation of the full processes at both hadron and lepton colliders with a variety of collision energies are performed to estimate the allowed region on and . Especially we find that, with the help of exclusively choosing longitudinal polarizations in the final states and suitable cuts,  $WW$  process is as important as the more widely studied triple Higgs production ( $hhh$ ) in the measurement of Higgs self-couplings. Our analysis indicates that these processes can play important roles in the measurement of Higgs self-couplings at future 100 TeV pp colliders and muon colliders. However, their cross sections are generally tiny at low energy machines, which makes them much more challenging to explore.



**Di-Higgs/Scalar / 66****Effective picture of cosmic bubble expansion****Authors:** Huaike Guo<sup>1</sup>; huaike guo<sup>2</sup><sup>1</sup> *Institute of Theoretical Physics*<sup>2</sup> *School of Physics Peking University*

We derive an effective equation-of-motion for an expanding bubble wall in the thermal plasma with a general form of the thermal friction. The efficiency factor for gravitational waves productions from colliding bubble walls is obtained with a special interest for the strong first-order phase transition. Primary authors Prof. Rong-Gen Cai (ITP-CAS), Shao-Jiang Wang

**BSM Higgs sector / 67****Long-lived Particle(LLP) searching at the future collider****Author:** Yulei Zhang<sup>1</sup><sup>1</sup> *Shanghai Jiaotong University*

In contrast to the standard model particles, long-lived particles(LLPs) decay with a significant distance from the primary vertex. We present a novel LLPs search strategy based on machine learning (ML) and image recognition techniques utilizing detector information from future lepton colliders.

68

**Higgs Alignment and Novel CP-Violating Observables in 2HDM****Corresponding Authors:** hcwangxiaoping@buaa.edu.com, hcwangxiaoping@163.com

69

**Selected results in the 3HDMs****Corresponding Author:** ivanov@mail.sysu.edu.cn

70

**Boosted technique for Higgs measurement and New Physics Searches****Corresponding Author:** qliphy0@pku.edu.cn

71

**Affleck-Dine Leptogenesis from Higgs Inflation**

Corresponding Author: hanchch@mail.sysu.edu.cn

72

**Search for doubly and singly charged Higgs bosons decaying into vector bosons in multi-lepton final states with the ATLAS detector using proton-proton collisions at 13 TeV**

Corresponding Author: hanlinxuy@gmail.com

73

**New Physics bounded from above: 2HDM confronting strong 1st order electroweak phase transition**

Corresponding Author: mczhangchina@gmail.com

74

**Search for heavy Higgs decaying to two tau leptons with the ATLAS detector using pp collision at 13 TeV**

Corresponding Authors: hanfei.ye@cern.ch, 18108085770@163.com

75

**Search for Resonant and Non-resonant VHH Production with ATLAS detector**

Corresponding Author: zhongyukun.xu@cern.ch

76

**Probe BSM physics with the interaction between the Higgs boson and top quark**

Corresponding Author: hualin.mei@cern.ch

77

**Searching for Long-lived Particles at future lepton collider**

Di-Higgs/Scalar / 78

## **Composite Higgs and Dark Matter**

Corresponding Author: ccfspesysu@gmail.com

Di-Higgs/Scalar / 79

## **Higgs potential and phase transition**

Corresponding Author: jshu@itp.ac.cn

Di-Higgs/Scalar / 80

## **Non-resonant di-Higgs process at LHC**

Di-Higgs/Scalar / 81

## **BSM resonance search for HH (SH, SS) at LHC**

82

## **Lunch time**

Higgs precise measurement / 83

## **Higgs Overview**

Corresponding Author: liantaow@gmail.com

Higgs precise measurement / 84

## **Precise theoretical predictions for Higgs processes**

Corresponding Author: j.wang@sdu.edu.cn

Higgs precise measurement / 85

## **Precise Higgs measurements at LHC**

**Corresponding Author:** nan.lu@cern.ch

**Higgs precise measurement / 86**

## **Search for rare Higgs processes**

**Corresponding Author:** huangyp@mail.ihep.ac.cn

**Pre-School / 87**

## **Introduction to the standard model and electro-weak symmetry breaking**

**Pre-School / 88**

## **Extension of standard model, e.g. 2HDM**

**Corresponding Author:** chenning\_symmetry@nankai.edu.cn

**Pre-School / 89**

## **Experimental Higgs physics, e.g. discovery, precise measurement, BSM Higgs searches**

**Corresponding Author:** yanwen@ustc.edu.cn

**Di-Higgs/Scalar / 90**

## **The multilepton anomalies at the LHC and the prospect of new physics at the EW scale**

**Corresponding Author:** amohammed@ihep.ac.cn

A study showed that the multilepton anomalies in the LHC data could be explained by adding two real scalar bosons to the SM. This study provides a more comprehensive set of LHC results of the multilepton production. We find that combining such results leads to substantial inconsistencies between the data and SM Monte Carlo predictions. These discrepancies appear in corners of the phase-space where different SM processes dominate, indicating that the potential miss modelling of a single SM process is unlikely to explain these excesses.

**BSM Higgs sector / 91**

## **Searches for light scalar and charged scalar**

**Corresponding Author:** wangjin3@ihep.ac.cn

**BSM Higgs sector / 92**

## **Heavy neutral scalar searches**

**Corresponding Author:** lailin.xu@cern.ch

**BSM Higgs sector / 93**

## **BSM Higgs couplings**

**Corresponding Author:** qinghongcao@pku.edu.cn

**BSM Higgs sector / 94**

## **Higgs bosons in new physics models**

**Corresponding Author:** jmyang@itp.ac.cn

**BSM Higgs sector / 95**

## **BSM Higgs searches in final states with top quarks**

96

## **Opening**

97

## **Speech from NJU**

**Higgs precise measurement / 98**

## **Physics Highlights at CEPC**

**Corresponding Author:** ruanmq@ihep.ac.cn

99

### **会议通知**

**Openning / 100**

### **Openning**

**Openning / 101**

### **Welcome from NJU**

**Pre-School / 102**

### **School opening**

**BSM Higgs sector / 103**

### **Close**

**Corresponding Author:** leizhang1801@nju.edu.cn