

# A unified formulation of one-loop tensor integrals for finite-volume effects

Ze-Rui Liang

(梁泽锐·湖南大学)

第七届手征有效场论研讨会

东南大学·南京, October 15-17, 2022

Based on [[ZRL](#)] and De-Liang Yao(姚德良), arXiv: 2207.11750]

# Outline

- Introduction
- Decomposition of one-loop tensor integrals
  - Definition of loop integrals for FVC
  - Decomposition of the FVC tensor integrals
  - Evaluation of the coefficients
- Reduction of tensor coefficients
  - Center-of-mass frame
  - Passarino-Veltman reduction
- A pedagogic example of application
- Summary and outlook

# Finite Volume Corrections

- The finite volume correction (FVC) for a given quantity  $Q$  is given by

$$\delta Q = Q(L) - Q(\infty)$$

- $Q(L)$  and  $Q(\infty)$  are calculated in the finite volume and infinite volume, respectively.
- FVC are not only of theoretical interest, but also the need in the precise extraction of physical results in lattice QCD simulation.
- The Lüscher formula provides an approach to calculate FVC to masses.
  - Lüscher formula relates the finite size mass shift to an integral of a special amplitude, evaluated in the infinite volume. [M. Luscher, Commun. Math. Phys. 104, 177-20(1986)]
  - Its application to the study of the FVC to the masses, pions, nucleon and heavy mesons, has been made.  
[G. Colangelo, et al, EPJC 33, (2004)], [G. Colangelo, et al, NPB 721, (2005)], [G. Colangelo, et al, PRD 82, 034506 (2010)]
- This approach fails in generating exponential terms beyond leading order.
- A resummed version [G. Colangelo, et al, NPB 721, (2005)] or a Lüscher-formula-like asymptotic [G. Colangelo, et al, PLB 590 (2004), 258-264] expression was proposed. But the feasibility of the Lüscher formula approach is rather limited.

# Chiral perturbation theory

- At finite volume, another systematical and popular tool to evaluate FVC is the ChPT.

[J. Gasser, H. Leutwyler, PLB 184 (1987) 83], [J. Gasser, H. Leutwyler, PLB 188 (1987) 477], [J. Gasser, H. Leutwyler, NPB 307 (1988) 763]

- The Lagrangian is the same as the infinite case.
- In a cubic box, momentum is discretized where the boundary conditions are imposed.




Fig. from [Alessio Giovanni Willy Vaghi, PHD thesis, (2015)]

- We are interested in ChPT for  $p$ -regime:

$$M_\pi L \gg 1$$

# Chiral perturbation theory

- A multitude of works concerning FVC based on ChPT have been done :
  - Masses: [S. R. Beane, PRD 70, 034507 (2004)], [L. S. Geng, *et al*, PRD 84, 074024 (2011)], [L. Alvarez-Ruso, *et al*, PRD 88, 054507 (2013)], [D. L. Yao, PRD 97, 034012 (2018)], [D. Severt, *et al*, CTP. 72, 075201 (2020)]
  - Decay constants: [D. Becirevic, *et al*, PRD 69, 054010 (2004)], [L. S. Geng, *et al*, PRD 89, 113007 (2014)]
  - Nucleon electric dipole moments: [T. Akan, *et al*, PLB 736, 163-168 (2014)]
  - Scalar form factors in  $K_{\ell 3}$  semi-leptonic decay: [K. Ghorbani, *et al*, EPJC 71, 1671 (2011)]
  - FVC to forward Compton scattering off the nucleon: [J. L. de la Parra, *et al*, PRD 103, 034507 (2021)]
  - ...
- Calculations of FVC in ChPT are tedious :
  - Complexity occurs in the one-loop analyses.
  - Automation of the one-loop calculations of FVC is still unavailable.
  - Expressions of the results for a given quantity might be different in form.
- **Our work**
  - Intend to give a unified description of the one-loop tensor integrals in a finite volume.
  - Generalize tensor decomposition of the one-loop tensor integrals to the FVC case, and derive a compact formula for the tensor coefficients.
  - Investigate the feasibility of the PV reduction of the tensor integrals.

# Definition of loop integrals for FVC I

- General form of one-loop  $N$ -point rank- $P$  tensor integrals

$$T^{N,\mu_1,\dots,\mu_P} = \frac{1}{i} \int \frac{d^d k}{(2\pi)^d} \frac{k^{\mu_1} \dots k^{\mu_P}}{D_1 D_2 \dots D_N}, \quad D_j = [(k + p_{j-1})^2 - m_j^2 + i0^+]$$

with  $p_0 = 0$ ,  $j = 1, 2, \dots, N$ , and an infinitesimal imaginary part  $i0^+$ .

- The finite-volume tensor integrals

- In a cubic box of volume  $V = L^3$ , the periodic boundary conditions  $\mathbf{k}_n = \frac{2\pi\mathbf{n}}{L}$

$$\int \frac{d\mathbf{k}}{(2\pi)^3} F(\mathbf{k}) \rightarrow \frac{1}{L^3} \sum_{\mathbf{n}} F(\mathbf{k}_n), \quad \mathbf{n} \equiv (n_1, n_2, n_3) \text{ with } n_i \in \mathbb{Z}$$

- The tensor integrals at finite volume are

$$T_V^{N,\mu_1,\dots,\mu_P} = \frac{1}{i} \left( \int \frac{1}{L^3} \sum_{\mathbf{n}} \int \frac{d\mathbf{k}^0}{2\pi} \right) \frac{k^{\mu_1} \dots k^{\mu_P}}{D_1 D_2 \dots D_N} \equiv \frac{1}{i} \int_V \frac{d^d k}{(2\pi)^d} \frac{k^{\mu_1} \dots k^{\mu_P}}{D_1 D_2 \dots D_N}.$$

# Definition of loop integrals for FVC II

- Poisson summation formula

$$\frac{1}{L^3} \sum_{\mathbf{n}} F(\mathbf{k}_n) = \sum_{\mathbf{n}} \int \frac{d\mathbf{k}}{(2\pi)^3} e^{i\mathbf{l}_k \cdot \mathbf{k}} F(\mathbf{k}_n).$$

- Then the finite-volume tensor integrals are

$$T_V^{N, \mu_1, \dots, \mu_P} = \sum_{\mathbf{n}} \frac{1}{i} \int_V \frac{d^d k}{(2\pi)^d} e^{-i\mathbf{l}_k \cdot \mathbf{k}} \frac{k^{\mu_1} \dots k^{\mu_P}}{D_1 D_2 \dots D_N},$$

with a four vector  $l_k^\mu = (0, \mathbf{n}L) = n^\mu L$ .  $|\mathbf{n}| = 0$  represents the infinite-volume contribution.

- The difference between the infinite and finite cases defines the FVC, and the tensor integrals for FVC are

$$\tilde{T}^{N, \mu_1, \dots, \mu_P} = \sum_{\mathbf{n} \neq 0} \frac{1}{i} \int_V \frac{d^d k}{(2\pi)^d} e^{-i\mathbf{l}_k \cdot \mathbf{k}} \frac{k^{\mu_1} \dots k^{\mu_P}}{D_1 D_2 \dots D_N}.$$

# Decomposition of the FVC tensor

- Considering the discretization effects at finite volume, a unit space-like vector  $n^\mu = (0, \mathbf{n})$  is introduced.

$$\tilde{L}^{\mu_1 \cdots \mu_P} = \left\{ \underbrace{g \cdots g}_s p \cdots p \underbrace{n \cdots n}_r \right\}_{i_{2s+1} \cdots i_{P-2s-r}}^{\mu_1 \cdots \mu_P},$$

- $2s$  out of  $P$  indices are distributed over the metric tensors and any pair of them are symmetrical.
- the  $n$ -vectors occupy  $r$  indices from the remaining ones.
- the rest indices are assigned to the momenta
- the number of terms

$$\frac{P!}{2^s s! r! (P - 2s - r)!}$$



# Examples

- Some instructive examples

$$\{pp \cdots p\}_{i_1 i_2 \cdots i_P}^{\mu_1 \mu_2 \cdots \mu_P} = p_{i_1}^{\mu_1} p_{i_2}^{\mu_2} \cdots p_{i_P}^{\mu_P} ,$$

$$\{pn\}_{i_1}^{\mu_1 \mu_2} = p_{i_1}^{\mu_1} n^{\mu_2} + n^{\mu_1} p_{i_1}^{\mu_2} ,$$

$$\{ppn\}_{i_1 i_2}^{\mu_1 \mu_2 \mu_3} = p_{i_1}^{\mu_1} p_{i_2}^{\mu_2} n^{\mu_3} + p_{i_1}^{\mu_1} n^{\mu_2} p_{i_2}^{\mu_3} + n^{\mu_1} p_{i_1}^{\mu_2} p_{i_2}^{\mu_3} ,$$

$$\{pnn\}_{i_1}^{\mu_1 \mu_2 \mu_3} = p_{i_1}^{\mu_1} n^{\mu_2} n^{\mu_3} + n^{\mu_1} p_{i_1}^{\mu_2} n^{\mu_3} + n^{\mu_1} n^{\mu_2} p_{i_1}^{\mu_3} ,$$

$$\{gn\}_{i_1}^{\mu_1 \mu_2 \mu_3} = g^{\mu_1 \mu_2} n^{\mu_3} + g^{\mu_1 \mu_3} n^{\mu_2} + g^{\mu_2 \mu_3} n^{\mu_1} ,$$

$$\begin{aligned} \{gpn\}_{i_1}^{\mu_1 \mu_2 \mu_3 \mu_4} &= g^{\mu_1 \mu_2} (p_{i_1}^{\mu_3} n^{\mu_4} + n^{\mu_3} p_{i_1}^{\mu_4}) + g^{\mu_1 \mu_3} (p_{i_1}^{\mu_2} n^{\mu_4} + n^{\mu_2} p_{i_1}^{\mu_4}) \\ &\quad + g^{\mu_1 \mu_4} (p_{i_1}^{\mu_2} n^{\mu_3} + n^{\mu_2} p_{i_1}^{\mu_3}) + g^{\mu_2 \mu_3} (p_{i_1}^{\mu_1} n^{\mu_4} + n^{\mu_1} p_{i_1}^{\mu_4}) \\ &\quad + g^{\mu_2 \mu_4} (p_{i_1}^{\mu_1} n^{\mu_3} + n^{\mu_1} p_{i_1}^{\mu_3}) + g^{\mu_3 \mu_4} (p_{i_1}^{\mu_1} n^{\mu_2} + n^{\mu_1} p_{i_1}^{\mu_2}) , \end{aligned}$$

$$\{gg\}_{i_1}^{\mu_1 \mu_2 \mu_3 \mu_4} = g^{\mu_1 \mu_2} g^{\mu_3 \mu_4} + g^{\mu_1 \mu_3} g^{\mu_2 \mu_4} + g^{\mu_1 \mu_4} g^{\mu_2 \mu_3} .$$

# Decomposition of the FVC tensor integrals

- The one-loop tensor integrals can be decomposed into the form as

$$\tilde{T}^{N,\mu_1\cdots\mu_P} = \sum_{\mathbf{n} \neq 0} \vec{T}^{N,\mu_1\cdots\mu_P}$$

with

$$\vec{T}^{N,\mu_1\cdots\mu_P} = \sum_{s=0}^{[P/2]} \sum_{r=0}^{P-2s} \sum_{\substack{i_{2s+1}=1, \\ \dots \\ i_{P-2s-r}=1}}^{N-1} \left\{ \underbrace{g \cdots g}_s \underbrace{p \cdots p}_r \underbrace{n \cdots n}_r \right\}^{\mu_1 \cdots \mu_P}_{i_{2s+1} \cdots i_{P-2s-r}} \vec{T}^N_{\underbrace{0 \cdots 0}_{2s} i_{2s+1} \cdots i_{P-2s-r} \underbrace{N \cdots N}_r}.$$

- $[P/2]$  is the floor function.
- Tensor coefficient  $\vec{T}^N_{0 \cdots 0 i_{2s+1} \cdots i_{P-2s-r} N \cdots N}$  is invariant with respect to permutation of the subscripts  $i_j$ , i.e.  $\vec{C}_{001233} = \vec{C}_{002133}$ .
- the subscripts “N” are unique in the finite volume.

# Examples

- Decomposition of the FVC tensor integrals up to rank 3


$$\vec{T}^{N,\mu} = \sum_{i=1}^{N-1} p_i^\mu \vec{T}_i^N + n^\mu \vec{T}_N^N ,$$

$$\vec{T}^{N,\mu\nu} = g^{\mu\nu} \vec{T}_{00}^N + \sum_{i,j=1}^{N-1} p_i^\mu p_j^\nu \vec{T}_{ij}^N + \sum_{i=1}^{N-1} \{pn\}_i^{\mu\nu} \vec{T}_{iN}^N + n^\mu n^\nu \vec{T}_{NN}^N ,$$

$$\begin{aligned} \vec{T}^{N,\mu\nu\rho} = & \sum_{i=1}^{N-1} \{gp\}_i^{\mu\nu\rho} \vec{T}_{00i}^N + \{gn\}^{\mu\nu\rho} \vec{T}_{00N}^N + \sum_{i,j,k=1}^{N-1} p_i^\mu p_j^\nu p_k^\rho \vec{T}_{ijk}^N + \sum_{i,j=1}^{N-1} \{ppn\}_{ij}^{\mu\nu\rho} \vec{T}_{ijN}^N \\ & + \sum_{i=1}^{N-1} \{pnn\}_i^{\mu\nu\rho} \vec{T}_{iNN}^N + n^\mu n^\nu n^\rho \vec{T}_{NNN}^N , \end{aligned}$$

# Evaluation of the coefficients

- Technical steps:



# Evaluation of the coefficients

- In the end, the one-loop tensor integrals have the form

$$\begin{aligned} \tilde{T}^{N, \mu_1, \dots, \mu_P} = & \sum_{\mathbf{n} \neq 0} \sum_{s=0}^{[P/2]} \sum_{r=0}^{P-2s} \sum_{\substack{i_{2s+1}=1 \\ \dots \\ i_{P-2s-r}=1}}^{N-1} \underbrace{\{g \dots g\}}_s \underbrace{p \dots p}_r \underbrace{n \dots n}_r \}_{i_{2s+1}, \dots, i_{P-2s-r}}^{\mu_1 \mu_2 \dots \mu_P} \frac{(-1)^{N+P-s-r}}{(4\pi)^{d/2} 2^s} \left( \frac{iL}{2} \right)^r \\ & \times \int_0^1 dX_N X_N^{i_{2s+1}} \dots X_N^{i_{P-2s-r}} e^{iL \mathbf{k} \cdot \mathcal{P}_N} \mathcal{K}_{N-s-r-\frac{d}{2}} \left( \frac{|\mathbf{n}|^2 L^2}{4}, \mathcal{M}_N^2 \right), \end{aligned}$$

with  $\int dX_N \equiv \frac{1}{\Gamma(N)} \int_0^1 d\mathcal{X}_N = \int_0^1 dx_1 \dots \int_0^1 dx_{N-1} x_2 \dots x_{N-1}^{N-2}$ .

- A **general expression** for the coefficients reads

$$\begin{aligned} \vec{T}_{\underbrace{0 \dots 0}_{2s}}^{i_{2s+1} \dots i_{P-2s-r}} \underbrace{N \dots N}_r = & \frac{2}{(4\pi)^{d/2}} \frac{(-1)^{N+P-s-r}}{2^s} \left( \frac{iL}{2} \right)^r \int_0^1 dX_N X_N^{i_{2s+1}} \dots X_N^{i_{P-2s-r}} e^{iL \mathbf{n} \cdot \mathcal{P}_N} \\ & \times \left( \frac{|\mathbf{n}|^2 L^2}{4\mathcal{M}_N^2} \right)^{\frac{N-s-r-d/2}{2}} K_{|N-s-r-\frac{d}{2}|}(|\mathbf{n}|L\mathcal{M}_N). \end{aligned}$$

# Center-of-Mass frame

- It is convenient to compute FVC in the rest frame or in the CM frame, where the net three momentum is zero.

$$l_k \cdot p_i = 0 \iff n \cdot p_i = 0, \quad i = 1, \dots, N-1.$$

- e.g. elastic two-body forward scattering at threshold, mass renormalization in the rest frame are satisfied by this condition.
- This condition lead to the  $\tilde{L}^{\mu_1 \dots \mu_P}$  tensors with odd  $n$ -vectors vanish. And then the dependence on  $\mathbf{n}$  of the rank- $P$  tensor can be relieved

$$\sum_{\mathbf{n} \neq 0} n^{\mu_1} \dots n^{\mu_{2t}} F(n^2) = \frac{1}{2^t (d_s/2)_t} \{h \dots h\}^{\mu_1 \dots \mu_{2t}} \sum_{\mathbf{n} \neq 0} (n^2)^t F(n^2),$$

- The auxiliary tensor  $h_{\mu\nu}$  is defined as  $h_{\mu\nu} \equiv g_{\mu\nu} - \bar{h}_\mu \bar{h}_\nu = \text{diag}(0, -1, -1, -1)$  with  $\bar{h}_\mu = (1, 0, 0, 0)$ , which serves to eliminate the zero-th component of the vector.
- The rank- $P$  tensor is irrelevant of  $\mathbf{n}$ , and enable us to perform the sum over  $\mathbf{n}$  in advance.

# Tensor coefficients of FVC integrals in CM frame

- The tensor decomposition of the FVC integrals

$$\tilde{T}^{N, \mu_1 \dots \mu_P} = \sum_{s=0}^{\lfloor \frac{P}{2} \rfloor} \sum_{t=0}^{\lfloor \frac{P-2s}{2} \rfloor} \sum_{\substack{i_{2s+1}=1 \\ \dots \\ i_{P-2s-2t}=1}}^{N-1} \underbrace{\{g \dots g\}}_s \underbrace{p \dots p h \dots h}_t^{\mu_1 \mu_2 \dots \mu_P} \tilde{T}_{0 \dots 0}^N \underbrace{i_{2s+1} \dots i_{P-2s-2t}}_{2s} \underbrace{N \dots N}_{2t}.$$

- The  $\mathbf{n}$ -independent coefficients are

$$\tilde{T}_{0 \dots 0}^N \underbrace{i_{2s+1} \dots i_{P-2s-2t}}_{2s} \underbrace{N \dots N}_{2t} = \frac{1}{2^t (d_s/2)_t} \sum_{\mathbf{n} \neq 0} [(n^2)^t \tilde{T}_{0 \dots 0}^N \underbrace{i_{2s+1} \dots i_{P-2s-2t}}_{2s} \underbrace{N \dots N}_{2t}].$$

- Now the equation relies merely on  $n^2$ , then the triple sum can be replaced by a single sum  $n_s \equiv n_1^2 + n_2^2 + n_3^2$  once the multiplicity  $\vartheta(n_s)$  for a given  $n_s$  takes into account.

$$\tilde{T}_{0 \dots 0}^N \underbrace{i_{2s+1} \dots i_{P-2s-2t}}_{2s} \underbrace{N \dots N}_{2t} = \frac{(-1)^t}{2^t (d_s/2)_t} \sum_{n_s > 0} [\vartheta(n_s) n_s^t \tilde{T}_{0 \dots 0}^N \underbrace{i_{2s+1} \dots i_{P-2s-2t}}_{2s} \underbrace{N \dots N}_{2t}].$$

# PV reduction of one-point tensor integrals

- For one-point tensor integrals, they can only be contracted by the metric tensor, and then the recurrence relations

$$[(d-1) + 2(t-1)] \underbrace{\tilde{A}_{0\dots 0}}_{2s} \underbrace{1\dots 1}_{2t} + [d + 2s + 4(t-1)] \underbrace{\tilde{A}_{0\dots 0}}_{2s+2} \underbrace{1\dots 1}_{2t-2} = m_1^2 \underbrace{\tilde{A}_{0\dots 0}}_{2s} \underbrace{1\dots 1}_{2t-2} .$$

- Specifically, the relations of one-point tensor integrals are, i.e.

$$d\tilde{A}_{00} + (d-1)\tilde{A}_{11} = m_1^2 \tilde{A}_0 ,$$

$$(d+2)\tilde{A}_{0000} + (d-1)\tilde{A}_{0011} = m_1^2 \tilde{A}_{00} .$$

- All the relations can either be checked numerically or be verified by the recurrence relations of the modified Bessel functions  $K_z(Y)$ .
- All the one-loop FVC integrals can be reduced to a linear combination of  $\underbrace{\tilde{A}_{0\dots 0}}_{2s}$ .


$$\underbrace{\tilde{A}_{0\dots 0}}_{2s} \underbrace{1\dots 1}_{2t} = \sum_{i=0}^t \left\{ \frac{[m_1^2]^{t-i}}{\prod_{j=1}^t a(j)} \sum_{\substack{i_1=0 \\ \vdots \\ i_t=0}}^1 \left[ \delta_{i, \sum_{j=1}^t i_j} \prod_{j=1}^t [b(j)]^{i_j} \right] \underbrace{\tilde{A}_{0\dots 0}}_{2(s+i)} \right\} ,$$

where  $a(j) = (d-1) + 2(j-1)$ ,  $b(j) = -[d + 2s + 4(j-1)]$ , and  $\delta$  is the Kronecker delta.



# PV reduction of one-point tensor integrals


- Schematic roadmap for PV reduction of one-loop FVC tensor integrals



- Dashed lines** : represent simplification operations by the recursive use of the recurrence relations.
- The  $\tilde{A}_0$ ,  $\tilde{A}_{00}$ ,  $\tilde{A}_{0000}$ , etc, can be adopted as the tensor basis.

# PV reduction of two-point tensor integrals


- Schematic roadmap for PV reduction of two-point FVC tensor integrals



- **Dashed lines** : the number of subscripts “2” is reduced by recursively utilizing the relation deduced by contracting the  $g_{\mu\nu}$ .
- **Solid lines** : the indices “1” can be eliminated by making use of the relation obtained by contracting of the external momentum  $p_{1\mu}$ .
- Like the case for one-point integrals, the tensor coefficients only with even numbers of “0” survive.

# PV reduction of $N$ -point tensor integrals

- Schematic roadmap for PV reduction of  $N$ -point FVC tensor coefficients




- Dashed lines** : by recursively utilizing the relation deduced by contracting the  $g_{\mu\nu}$ .
- Solid lines** : by making use of the relation obtained by contracting of the external momentum  $p_j^{\mu_1}$ .
- The **boxed coefficients** are chosen as the tensor basis.

- It is a first attempt and only aim at finding out the feasibility of PV reduction and the existence of a tensor basis for the one-loop integrals at finite volume.

# The FVC of nucleon mass

- Leading one-loop Feynman diagrams contributing the nucleon mass



- The self-energy of the nucleon can be expressed as

$$\Sigma(\not{p}, \not{p}) = \sum_{\mathbf{n} \neq 0} [\mathcal{A} + \not{p}\mathcal{B} + \not{p}\mathcal{C}]$$

- $\mathcal{A}$ ,  $\mathcal{B}$  and  $\mathcal{C}$  are functions of the scalar products of the external momentum and the unit space-like vectors.
- The occurrence of the third term is due to the introduction of spatial boundary conditions of the finite volume.

# The FVC of nucleon mass

- The self-energy functions for (a)

$$\mathcal{A}_a = \frac{3g_A^2 m_N}{4F_\pi^2} \left\{ s\vec{B}_0 + 2s\vec{B}_1 + d\vec{B}_{00} + s\vec{B}_{11} + n^2\vec{B}_{22} - 2n \cdot p \left[ \vec{B}_2 + \vec{B}_{12} \right] \right\} ,$$

$$\mathcal{B}_a = \frac{3g_A^2}{4F_\pi^2} \left\{ s\vec{B}_1 + 2s\vec{B}_{11} + 2d\vec{B}_{00} + (d+2)\vec{B}_{001} + s\vec{B}_{111} + n^2(2\vec{B}_{22} + \vec{B}_{122}) \right. \\ \left. - 2n \cdot p \left[ \vec{B}_2 + 2\vec{B}_{12} + \vec{B}_{112} \right] \right\} ,$$

$$\mathcal{C}_a = \frac{3g_A^2}{4F_\pi^2} \left\{ s\vec{B}_2 - (d+2)\vec{B}_{002} - s\vec{B}_{112} - n^2\vec{B}_{222} + 2n \cdot p\vec{B}_{122} \right\} ,$$

- $s = p^2$ .
- $g_A$  is the axial coupling constant,  $F_\pi$  is the pion decay constant, and  $m_N$  denotes the nucleon mass in the chiral limit.

# The FVC of nucleon mass

- In the CM frame, one has  $\bar{u}(p)\not{p}u(p) = 0$ . And the self-energy functions for (a) can be simplified to

$$\mathcal{A}_a = \frac{3g_A^2 m_N}{4F_\pi^2} \left\{ s\tilde{B}_0 + 2s\tilde{B}_1 + d\tilde{B}_{00} + s\tilde{B}_{11} + (d-1)\tilde{B}_{22} \right\} ,$$

$$\mathcal{B}_a = \frac{3g_A^2}{4F_\pi^2} \left\{ s\tilde{B}_1 + 2s\tilde{B}_{11} + 2d\tilde{B}_{00} + (d+2)\tilde{B}_{001} + s\tilde{B}_{111} + (d-1) \left[ 2\tilde{B}_{22} + \tilde{B}_{122} \right] \right\} .$$

- The form is by making use of PV reduction

$$\mathcal{A}_a(L) = \frac{3g_A^2 m_N}{4F_\pi^2} \left\{ \tilde{A}_0(m_N^2; L) + M_\pi^2 \tilde{B}_0(m_N^2, m_N^2, M_\pi^2; L) \right\} ,$$

$$\mathcal{B}_a(L) = \frac{1}{m_N} \mathcal{A}_a(L) .$$

where  $M_\pi$  is the pion mass and  $L$  is the size of the spatial cubic box.

# The FVC of nucleon mass

- The self-energy functions for (b)

$$\mathcal{A}_b(L) = -\frac{h_A^2}{3F_\pi^2 m_\Delta} \left\{ (m_\Delta^2 - m_N^2 + 3M_\pi^2) \tilde{A}_0(M_\pi^2; L) - (m_\Delta^2 + m_N^2 - M_\pi^2) \tilde{A}_0(m_\Delta^2; L) \right. \\ \left. + \lambda(m_\Delta^2, m_N^2, M_\pi^2) \tilde{B}_0(m_N^2, m_\Delta^2, M_\pi^2; L) \right\},$$

$$\mathcal{B}_b(L) = \frac{h_A^2}{6F_\pi^2 m_\Delta^2 m_N^2} \left\{ \lambda(m_\Delta^2, m_N^2, M_\pi^2) \tilde{A}_0(m_\Delta^2; L) - [(m_\Delta^2 - M_\pi^2)^2 - m_N^4 + 4m_N^2 M_\pi^2] \tilde{A}_0(M_\pi^2; L) \right. \\ \left. + 4m_N^2 [\tilde{A}_{00}(m_\Delta^2; L) - \tilde{A}_{00}(M_\pi^2; L)] \right. \\ \left. + \lambda(m_\Delta^2, m_N^2, M_\pi^2) (m_\Delta^2 + m_N^2 - M_\pi^2) \tilde{B}_0(m_N^2, m_\Delta^2, M_\pi^2; L) \right\},$$

- Källén function  $\lambda(a, b, c) = a^2 + b^2 + c^2 - 2ab - 2ac - 2bc$ .
- $h_A$  is the coupling constant of the  $\pi N \Delta$  interaction, and  $m_\Delta$  is the mass of the  $\Delta$  resonance in the chiral limit.


# A pedagogic example of application

- The expression of the FVC on the nucleon mass

$$m_N^{\text{FVC}}(L) = [\mathcal{A}(L) + m_N \mathcal{B}(L)]$$

with  $\mathcal{A}(L) = \mathcal{A}_a(L) + \mathcal{A}_b(L)$  and  $\mathcal{B}(L) = \mathcal{B}_a(L) + \mathcal{B}_b(L)$ .

- FVC to the nucleon mass




- The validity of the PV reduction for the FVC tensor coefficients is explicitly verified.
- The result of diagram (a) is identical to the one given in Ref. [L. Alvarez-Ruso, *et al*, PRD 88, 054507 (2013)].
- The contributions of the nucleon and delta loops are comparable with each other, which implies the importance of the  $\Delta$  resonance in the estimation of FVC to the nucleon mass.



# A pedagogic example of application

- The  $L$ -dependence of the nucleon mass with different pion mass.



- For a given finite size  $L$ , the larger the pion mass is, the smaller the FVC become.
- The effect of FVC on the nucleon mass becomes negligible when  $M_\pi L \gtrsim 3$ .

# Summary and Outlook

- A systematical formulation of one-loop tensor integrals for FVC has been advocated.
  - A compact formula for the tensor coefficients in the decomposition has been derived, which is suitable for numerical computations.
  - In the CM frame, the tensor coefficients can be simplified by means of PV reduction.
  - An example is given to illustrate the application of our formulation.
- 
- The formulation pave a path for efficient computations of FVC. (e.g. can be readily implemented in FeynCalc.)
  - Chiral extrapolation of Lattice QCD results with FVC and precise extraction of physical quantities. (e.g. doubly charmed baryons and Goldstone bosons)
  - Generalize to two-loop integrals.

Many thanks for your attention!