



A non-minimal SU(6) GUT and high-quality Axion

南开大学物理科学学院
陈宁

2021.07.14, @西藏高能物理论坛

based on [2106.00223](#)
with 刘宇统, 滕召隆

Background: GUT

Topics	Authors	Publications	Date
Fourth color unification	Pati, Salam	[5]	1974
SU(5) GUT	Georgi, Glashow	[6]	1974
SO(10) GUT	Fritzsch, Minkowski	[7]	1975
Peccei-Quinn mechanism	Peccei, Quinn	[8]	1977
Seesaw mechanism	Yanagida	[9]	1979
	Gell-Mann, Ramond, Slansky	[10]	1979
KSVZ axion	Kim	[11]	1979
	Shifman, Vainshtein, Zakharov	[12]	1980
SU(N + 4) global symmetry	Dimopoulos, Raby, Susskind	[13]	1980
SUSY SU(5) GUT	Dimopoulos, Georgi	[14]	1981
DFSZ axion	Zhitnitsky	[15]	1980
	Dine, Fischler, Srednicki	[16]	1981
Axion in SU(5) GUT	Wise, Georgi, Glashow	[17]	1981
Leptogenesis	Fukugita, Yanagida	[18]	1986
...			

Table 1: Collections of the major theoretical advances in the GUT and some related BSM physical issues listed in chronological order.

Background: GUT

- * Over 40 yrs since the $SU(5)$ and $SO(10)$, there is convincing evidence that the BSM new physics should be put forth to address:
 - (1) **Strong CP problem, e.g., the Peccei-Quinn mechanism**
 - (2) neutrino masses through the seesaw mechanism
 - (3) Baryon asymmetry through the baryogenesis/leptogenesis
 - (4) Dark matter

- * The simplest $SU(5)$, $SO(10)$ and their varieties, do not seem to include all necessary ingredients for BSM. One often needs to put new physical ingredients into the $SU(5)$ & $SO(10)$ GUTs by hand.

Background: GUT

- * In the *minimal* $SU(5)$ GUT, Georgi & Glashow assumed: “as few leptons (fermions) as possible, no unobserved leptons (fermions) ...”
- * Our assumption: *a successful GUT can address as many BSM issues as possible, with the minimal set of fields unless otherwise necessary.*
- * We consider the PQ-quality for the strong CP problem to start with, and we get more than that (see next).

Background: Strong CP

- * The strong CP problem, a topological term for the QCD vacuum

$$\mathcal{L}_\theta = \theta \frac{\alpha_{3c}}{8\pi} G_{\mu\nu}^a \tilde{G}^{\mu\nu a}, \text{ and experimentally from the neutron}$$

EDM: $|\bar{\theta}| \lesssim 10^{-10}$, with $\bar{\theta} = \theta + \arg \det M_q$, very different from the $\mathcal{O}(1)$ expectation of θ parameter.

- * PQ mechanism: to replace θ by a periodic pseudo-scalar field $a \rightarrow a + 2\pi f_a$, f_a is known as the axion decay constant. There is a classical window of $10^8 \text{ GeV} \lesssim f_a \lesssim 10^{12} \text{ GeV}$.

- * Axion induced potential: $V_{\text{QCD}} = \Lambda_{\text{QCD}}^4 [1 - \cos(a/f_a)]$.

Background: Strong CP

- * Invisible axion models such as Kim-Shifman-Vainshtein-Zakharov (KSVZ) and Dine-Fischler-Srednicki-Zhitnisky (DFSZ), a comes as a phase from a complex scalar field

$$\Phi = \frac{1}{\sqrt{2}}(v_a + \rho_a)\exp(ia/f_a).$$

- * PQ quality: $U(1)_{PQ}$ symmetry (expressed in terms of Φ) is global and put in by hand in KSVZ/DFSZ models, and the gravity does not respect global symmetries. ['92 Barr, Seckel, Kamionkowski, March-Russell, Holman, Hsu, Kephart, Kolb, Watkins, Widrow.]

Background: Strong CP

* A general operator of $\mathcal{O}_{PQ}^{d=2m+n} = k \frac{|\Phi|^{2m} \Phi^n}{M_{\text{pl}}^{2m+n-4}} + H.c.$ ['92

Kamionkowski, March-Russell] $\Delta PQ = n$ with $PQ(\Phi) = 1$.

* The $\mathcal{O}_{PQ}^{d=2m+n}$ shifts the V_{QCD} minima $|\bar{\theta}| = |\langle a \rangle / f_a| \lesssim 10^{-10}$

* if $|k| \sim 10^{-2}$ and $2m + n = 5$, $\Rightarrow f_a \lesssim 10 \text{ GeV}$, ruled out,
else if $f_a \sim 10^{12} \text{ GeV}$ and $2m + n = 5$, $\Rightarrow |k| \lesssim 10^{-55}$, very
fine-tuned.

* NB, the renormalizable operators with $2m + n \leq 4$ are even
worse in PQ-quality. This suggests to consider the underlying
gauge symmetry of the Φ .

Global Symmetries

- * The usual wisdom of a high-quality PQ is to have the $U(1)_{PQ}$ as an emergent global symmetry.
- * The QCD has the global symmetries of $\mathcal{G}_{\text{global}} = SU(3)_L \otimes SU(3)_R \otimes U(1)_V \otimes U(1)_A$, while QCD is vectorial.
- * The chiral gauge theory w.o. Unification: to put another confining sector with the SM, e.g. $SU(5) \otimes \mathcal{G}_{\text{SM}}$ by Gavela, Ibe, Quilez, and Yanagida [1812.08174].
- * Dimopoulos-Raby-Susskind (1980) studied a strongly-interacting theory: an anomaly-free $SU(N + 4)$ chiral gauge theory with N anti-fundamental fermions and one rank-2 anti-symmetric fermion, and it has $\mathcal{G}_{\text{global}} = SU(N) \otimes U(1)$, $N \geq 2$.

Our result:

- * We start from $SU(6)$, and identify $\mathcal{G}_{\text{global}} = SU(2)_F \otimes U(1)_{\text{PQ}}$.
- * Our finding is that a *non-minimal* $SU(6)$ GUT with its minimal fermion && Higgs setup can lead to:
 - (1) Automatic high-quality PQ symmetry breaking @ $10^8 \text{ GeV} \lesssim f_a \lesssim 10^{10} \text{ GeV}$, with an extended symmetry of $\mathcal{G}_{331} = SU(3)_c \otimes SU(3)_L \otimes U(1)_N$
 - (2) Automatic KSVZ vector-like quarks $m_D \sim f_a$, with fixed electric charge of $-1/3$.
 - (3) A cosmological-safe axion model, no DW formation
 - (4) Automatic Type-I seesaw mechanism with sterile neutrino mass at $\sim f_a$
 - (5) Automatic Type-II 2HDM at the EW scale

The $SU(6)$ model

The SU(6) model

- * The minimal anomaly-free SU(6) has fermions of: $2 \times \bar{\mathbf{6}}_F \oplus \mathbf{15}_F$
- * How to break the SU(6) to the $\mathcal{G}_{\text{SM}} = \text{SU}(3)_c \otimes \text{SU}(2)_L \otimes \text{U}(1)_Y$? The Georgi-Glashow SU(5) is a subgroup of the SU(6), while the direct breaking to the SU(5) is unwelcome. The sequential breaking of $\text{SU}(5) \rightarrow \mathcal{G}_{\text{SM}}$ leads to the proton decays with lower mass scale, hence faster decay rate.
- * Alternative pattern is: $\text{SU}(6) \rightarrow \mathcal{G}_{331} \rightarrow \mathcal{G}_{\text{SM}}$, with $\mathcal{G}_{331} = \text{SU}(3)_c \otimes \text{SU}(3)_L \otimes \text{U}(1)_N$. This is achievable with an adjoint Higgs of $\mathbf{35}_H$ at the GUT scale (1974 Ling-Fong Li).

The $SU(6)$ Higgs sector

- * An adjoint Higgs of 35_H at the GUT scale.
- * There is a brute-force method: to perform the tensor products of all $SU(6)$ fermions $\bar{6} \otimes \bar{6} = \bar{15} \oplus \bar{21}$, $\bar{6} \otimes 15 = 6 \oplus 84$, and $15 \otimes 15 = \bar{15} \oplus \bar{105} \oplus \bar{105}'$, and include all possible Higgs fields to form gauge-invariant Yukawa couplings.
- * Physical requirements: all SM Yukawa couplings should be reproduced $\Rightarrow 6_H$ (for d^i and ℓ^i) and 15_H (for u^i)
- * Two $6_H^{I,II}$ are needed to respect the $SU(2)_F$.
- * A 21_H is introduced for the sterile neutrino masses.

The SU(6) Higgs sector

- * minimal Higgs sector: $\mathbf{6}_H^{\alpha=I,II}$, $\mathbf{15}_H$, $\mathbf{21}_H$, $\mathbf{35}_H$
- * Hierarchies of Higgs VEVs: $\langle \mathbf{35}_H \rangle \sim \Lambda_{\text{GUT}}$,
 $\langle \mathbf{6}_H^{II} \rangle = v_3$, $\langle \mathbf{21}_H \rangle = v_6$, $v_3 \sim v_6 \sim v_{331}$
 $\langle \mathbf{6}_H^I \rangle = v_d = v_{\text{EW}} \sin \beta$, $\langle \mathbf{15}_H \rangle = v_u = v_{\text{EW}} \cos \beta$
 $\Lambda_{\text{GUT}} \gg v_{331} \gg v_{\text{EW}} = (\sqrt{2} G_F)^{-1/2} \simeq 246 \text{ GeV}$
- * $\mathbf{6}_H^{II}$ and $\mathbf{21}_H$ are responsible for the $\mathcal{G}_{331} \rightarrow \mathcal{G}_{\text{SM}}$ breaking.
- * Two Higgs doublets from $\mathbf{6}_H^I$ and $\mathbf{15}_H$ are responsible for the EWSB.
- * One bonus: free from the μ -problem in the SUSY extension, since one cannot form $W \supset \mu \mathbf{6}_H^{I*} \mathbf{15}_H$ term.

The SU(6) fermions

SU(6)	\mathcal{G}_{331}	\mathcal{G}_{SM}
$\bar{\mathbf{6}}_{\text{F}}^{\text{I}}$	$(\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\text{F}}^{\text{I}}$ $(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\text{F}}^{\text{I}}$	$(\bar{\mathbf{3}}, \mathbf{1}, +\frac{2}{3})_{\text{F}}^{\text{I}} : \underline{d_R^c}$ $(\mathbf{1}, \mathbf{2}, -1)_{\text{F}}^{\text{I}} : \underline{(e_L, -\nu_L)}$ $(\mathbf{1}, \mathbf{1}, 0)_{\text{F}}^{\text{I}} : \underline{N}$
$\bar{\mathbf{6}}_{\text{F}}^{\text{II}}$	$(\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_{\text{F}}^{\text{II}}$ $(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_{\text{F}}^{\text{II}}$	$(\bar{\mathbf{3}}, \mathbf{1}, +\frac{2}{3})_{\text{F}}^{\text{II}} : \underline{D_R^c}$ $(\mathbf{1}, \mathbf{2}, -1)_{\text{F}}^{\text{II}} : \underline{(e'_L, -\nu'_L)}$ $(\mathbf{1}, \mathbf{1}, 0)_{\text{F}}^{\text{II}} : \underline{N'}$
$\mathbf{15}_{\text{F}}$	$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{2}{3})_{\text{F}}$ $(\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})_{\text{F}}$ $(\mathbf{3}, \mathbf{3}, 0)_{\text{F}}$	$(\bar{\mathbf{3}}, \mathbf{1}, -\frac{4}{3})_{\text{F}} : \underline{u_R^c}$ $(\mathbf{1}, \mathbf{2}, +1)_{\text{F}} : \underline{(\nu_R'^c, e_R'^c)}$ $(\mathbf{1}, \mathbf{1}, +2)_{\text{F}} : \underline{e_R^c}$ $(\mathbf{3}, \mathbf{2}, +\frac{1}{3})_{\text{F}} : \underline{(u_L, d_L)}$ $(\mathbf{3}, \mathbf{1}, -\frac{2}{3})_{\text{F}} : \underline{D_L}$

The SU(6) Yukawa

* The most general Yukawa coupling:

$$15_F \bar{6}_F^\alpha 6_H^{\alpha*} + 15_F 15_F 15_H + \bar{6}_F^\alpha (i\sigma_2)_{\alpha\beta} \bar{6}_F^\beta 21_H + H.c.$$

* The PQ charge and a discrete \mathbb{Z}_4 symmetry (for PQ quality):

	$\bar{6}_F^\alpha$	15_F	6_H^α	15_H	21_H	35_H
$U(1)_{PQ}$	1	1	2	-2	-2	0
$SU(2)_F$	\square	1	\square	1	1	1
\mathbb{Z}_4	1	$\frac{1}{2}$	$\frac{3}{2}$	-1	-2	0

At the UV the global $U(1)_{PQ}[SU(6)]^2$ anomaly: $N_{SU(6)} = 9$

The $SU(6)$ Axion

The SU(6) Axion

* The physical axion field comes from: $\mathbf{6}_H^{\text{II}} \supset (\mathbf{1}, \mathbf{3}, +\frac{1}{3})_H \supset \frac{v_3}{\sqrt{2}} \exp(ia_3/v_3)$

and $\mathbf{21}_H \supset (\mathbf{1}, \mathbf{6}, +\frac{2}{3})_H \supset \frac{v_6}{\sqrt{2}} \exp(ia_6/v_6)$

* To impose an orthogonality condition between the $U(1)_{\text{PQ}}$

$J_{\text{PQ}}^\mu = q_3 v_3 (\partial^\mu a_3) + q_6 v_6 (\partial^\mu a_6)$ and the $U(1)_N$ $J_N^\mu = \frac{1}{3} v_3 (\partial^\mu a_3) + \frac{2}{3} v_6 (\partial^\mu a_6)$

currents. Physical charge: $q \equiv c_1 \text{PQ} + c_2 N$.

* 't Hooft global anomaly matching: $N_{\text{SU}(3)_c} = N_{\text{SU}(6)} \Rightarrow c_1 = 1$.

* $a_{\text{phys}} = \cos \phi a_3 + \sin \phi a_6$, $\tan \phi = \frac{v_3}{2v_6}$.

* Axion decay const: $9v_{331}^{-2} = \frac{1}{4}v_6^{-2} + v_3^{-2}$ and $f_a = v_{331}/18$.

The PQ quality

- * The leading PQ-breaking operator respecting the $SU(2)_F$ and \mathbb{Z}_4 :

$$\mathcal{O}_{PQ}^{d=6} = \left[\epsilon_{\alpha\beta} \mathbf{6}_H^\alpha \mathbf{6}_H^\beta \mathbf{15}_H \right]^2$$

$$\supset \left[\epsilon_{\alpha\beta} \epsilon_{abc} (\mathbf{1}, \mathbf{3}, +\frac{1}{3})^{a,\alpha} (\mathbf{1}, \mathbf{3}, +\frac{1}{3})^{b,\beta} (\mathbf{1}, \mathbf{3}, -\frac{2}{3})^c \right]^2$$

if no \mathbb{Z}_4 : $\mathcal{O}_{PQ}^{d=3} = \epsilon_{\alpha\beta} \epsilon_{abc} (\mathbf{1}, \mathbf{3}, +\frac{1}{3})^{a,\alpha} (\mathbf{1}, \mathbf{3}, +\frac{1}{3})^{b,\beta} (\mathbf{1}, \mathbf{3}, -\frac{2}{3})^c$ is

dangerous in PQ-quality.


- * Axion effective potential:

$$V = -\Lambda_{\text{QCD}}^4 \cos\left(\frac{a_{\text{phys}}}{f_a}\right) + \frac{|k| (v_u v_d v_3)^2}{4M_{\text{pl}}^2} \cos\left(\frac{a_{\text{phys}}}{6f_a} + \delta\right)$$

$$* \quad |\bar{\theta}| \equiv \left| \frac{\langle a_{\text{phys}} \rangle}{f_a} \right| \lesssim 10^{-10} \Rightarrow f_a \lesssim \frac{3.7 \times 10^7 \cos \phi}{|k \sin(\delta)|^{1/2}} \left(\tan \beta + \frac{1}{\tan \beta} \right) \text{ GeV}$$

This is solely determined by the symmetry consideration in a GUT!


The PQ quality



$$10^8 \text{ GeV} \lesssim f_a \lesssim 10^{10} \text{ GeV}$$

$$m_a = 5.70 \left(\frac{10^{12} \text{ GeV}}{f_a} \right) \mu\text{eV} \sim (10^{-4}, 10^{-2}) \text{ eV}$$

The axion searches



$$g_{a\gamma\gamma} = \left(\frac{E}{N_{\text{SU}(3)_c}} - 1.92 \right) \left(\frac{1.14 \times 10^{-3} \text{ GeV}}{f_a} \right) \text{ GeV}^{-1}$$

$$\text{U}(1)_{\text{PQ}}[\text{U}(1)_{\text{em}}]^2 \text{ anomaly factor : } E = \sum_f \text{PQ}_f \dim(\mathcal{C}_f) \text{Tr} q_f^2 = 24$$

The Axion domain walls

- * Back to the axion effective potential:

$$V = -\Lambda_{\text{QCD}}^4 \cos\left(\frac{a_{\text{phys}}}{f_a}\right) + \frac{|k| (v_u v_d v_3)^2}{4M_{\text{pl}}^2} \cos\left(\frac{a_{\text{phys}}}{6f_a} + \delta\right)$$

- * The $\cos\left(\frac{a_{\text{phys}}}{f_a}\right)$ term is periodic and has degenerate minima, this leads to the DWs.

- * DWs are problematic in cosmology, with the energy density

$\rho_{\text{DW}} \sim \sigma/t$. The energy densities for radiation/matter:

$\rho_{\text{rad}} \propto t^{-2}$, $\rho_{\text{matt}} \propto t^{-3/2}$. DWs can overtake the Universe once they are formed.

The Axion domain walls

* The second PQ-breaking term acts as the biased term to collapse the DWs. [Vilenkin ('81), Gelmini, Gleiser, Kolb, ('89), Larsson, Sarkar, White ('96)]

* To have DWs collapse before formation: $t_{\text{dec}} < t_{\text{form}}$.

* In our case:

$$t_{\text{form}} \sim 10^2 \text{ sec} \left(\frac{10^{13} \text{ GeV}}{v_{331}} \right) \sim \mathcal{O}(10^4) - \mathcal{O}(10^6) \text{ sec}$$

$$t_{\text{dec}} \approx \frac{\sigma_{\text{DW}}}{v_{331}^4} \sim 10^{-66} \text{ sec} \left(\frac{M_{\text{pl}} v_{331}}{v_u v_d} \right)^2 \left(\frac{10^{13} \text{ GeV}}{v_{331}} \right)^3$$
$$\sim \mathcal{O}(10^{-8}) - \mathcal{O}(10^{-6}) \text{ sec}$$

The $SU(6)$ Unification

The SU(6) unification

* The gauge couplings: $(\alpha_{3c}, \alpha_{3L}, \alpha_N)$ for the \mathcal{G}_{331} , and $(\alpha_{3c}, \alpha_{2L}, \alpha_Y)$ for the \mathcal{G}_{SM} . Use $\alpha_1 = \frac{4}{3}\alpha_N$ for the \mathcal{G}_{331} embedding into the SU(6).

* The one-loop RGEs of the SU(6):

$$\alpha_i^{-1}(\mu_2) = \alpha_i^{-1}(\mu_1) - \frac{b_i^{(1)}}{2\pi} \log\left(\frac{\mu_2}{\mu_1}\right)$$

* The matching conditions: $\alpha_{3L}^{-1}(v_{331}) = \alpha_{2L}^{-1}(v_{331})$,

$$\alpha_1^{-1}(v_{331}) = -\frac{1}{4}\alpha_{2L}^{-1}(v_{331}) + \frac{3}{4}\alpha_Y^{-1}(v_{331}).$$

The SU(6) unification

* non-SUSY: $b_i^{(1)} = -\frac{11}{3}C_2(\mathcal{G}_i) + \frac{2}{3} \sum_f T(\mathcal{R}_f^i) + \frac{1}{3} \sum_s T(\mathcal{R}_s^i)$

SUSY: $b_i^{(1)} = -3C_2(\mathcal{G}_i) + \sum_\chi T(\mathcal{R}_\chi^i)$

* SUSY extension: $\mathbf{21}_H$ super-multiplet is anomalous, we include a $\overline{\mathbf{21}}_H$.

* non-SUSY: $m_Z \leq \mu \leq v_{331} : (b_{\text{SU}(3)_c}^{(1)}, b_{\text{SU}(2)_L}^{(1)}, b_{\text{U}(1)_Y}^{(1)}) = (-7, -3, 7)$


$v_{331} \leq \mu \leq \Lambda_{\text{GUT}} : (b_{\text{SU}(3)_c}^{(1)}, b_{\text{SU}(3)_L}^{(1)}, b_{\text{U}(1)_1}^{(1)}) = (-5, -\frac{11}{3}, \frac{43}{6})$

* SUSY:

$m_Z \leq \mu \leq v_{331} : (b_{\text{SU}(3)_c}^{(1)}, b_{\text{SU}(2)_L}^{(1)}, b_{\text{U}(1)_Y}^{(1)}) = (-3, 1, 11)$

$v_{331} \leq \mu \leq \Lambda_{\text{GUT}} : (b_{\text{SU}(3)_c}^{(1)}, b_{\text{SU}(3)_L}^{(1)}, b_{\text{U}(1)_1}^{(1)}) = (0, \frac{13}{2}, \frac{29}{2})$

The SU(6) unification



$\alpha_{3c}(m_Z)$, $\alpha_{em}(m_Z)$, $\sin^2 \theta_W(m_Z)$ as inputs

The SU(6) unification

- * To impose the unification condition at the UV:

$$\alpha_{3c}^{-1}(\Lambda_{\text{GUT}}) = \alpha_{3L}^{-1}(\Lambda_{\text{GUT}}) = \alpha_1^{-1}(\Lambda_{\text{GUT}}) = \alpha_{\text{GUT}}^{-1}(\Lambda_{\text{GUT}})$$

- * Benchmark $v_{331} = 5.7 \times 10^{10}$ GeV, we find:

$$\Lambda_{\text{GUT}} \approx 7.8 \times 10^{15} \text{ GeV}, \quad \alpha_{\text{GUT}}^{-1}(\Lambda_{\text{GUT}}) = 18.14$$

$$\sin^2 \theta_W(m_Z) = 0.22923$$

$$\text{PDG: } \sin^2 \theta_W(m_Z) = 0.23117$$

- * Proton lifetime:

$$\tau[p \rightarrow e^+ \pi^0] \sim 10^{36} \text{ yrs} \left(\frac{\alpha_{\text{GUT}}^{-1}}{35} \right)^2 \left(\frac{\Lambda_{\text{GUT}}}{10^{16} \text{ GeV}} \right)^4$$

$$\approx 9.8 \times 10^{34} \text{ yrs}$$

$$\text{SK limit: } \tau_p \gtrsim 2.4 \times 10^{34} \text{ yrs}$$

Summary

- * We show a *non-minimal* $SU(6)$ GUT model with the minimal setup to achieve a high-quality axion by identifying the $U(1)_{PQ}$ as the Abelian component of the emergent global symmetries.
- * The axion decay constant: $10^8 \text{ GeV} \lesssim f_a \lesssim 10^{10} \text{ GeV}$ w.o. much fine-tuning of the EFT parameter.
- * The GUT spectrum contains vector-like KSVZ D -quarks, and heavy leptons & singlet neutrinos, type-I seesaw.
- * Safe from the cosmological constraints.

Backups

The SU(6) Yukawa

* At the $\mathcal{G}_{331} \rightarrow \mathcal{G}_{SM}$ breaking:

$$\mathbf{15}_F \bar{\mathbf{6}}_F^{\text{II}} \mathbf{6}_H^{\text{II}*} + H.c. \supset$$

$$(\mathbf{3}, \mathbf{3}, 0)_F \otimes (\bar{\mathbf{3}}, \mathbf{1}, +\frac{1}{3})_F^{\text{II}} \otimes (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_H^{\text{II}} + H.c.$$

$$(\mathbf{1}, \bar{\mathbf{3}}, +\frac{2}{3})_F \otimes (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_F^{\text{II}} \otimes (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_H^{\text{II}} + H.c.$$

$$\Rightarrow m_D \sim m_{e'} \sim m_{\nu'} \simeq \mathcal{O}(v_{331})$$

D -hadron lifetime: $\tau_D \sim m_D^{-1} \sim \mathcal{O}(10^{-36}) - \mathcal{O}(10^{-34})$ sec, Vs. the BBN constraint of $\tau_Q \lesssim 10^{-2}$ sec.

* $\bar{\mathbf{6}}_F^{\text{I}} \bar{\mathbf{6}}_F^{\text{II}} \mathbf{21}_H + H.c. \supset$

$$(\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_F^{\text{I}} \otimes (\mathbf{1}, \bar{\mathbf{3}}, -\frac{1}{3})_F^{\text{II}} \otimes (\mathbf{1}, \mathbf{6}, +\frac{2}{3})_H + H.c.$$

$$\Rightarrow m_{N, N'} \simeq \mathcal{O}(v_{331})$$

The SU(6) Yukawa

$$* \mathbf{15}_F \bar{\mathbf{6}}_F^I \mathbf{6}_H^{I*} + H.c. \supset$$

$$\left(\mathbf{3}, \mathbf{2}, +\frac{1}{3}\right)_F \otimes \left(\bar{\mathbf{3}}, \mathbf{1}, +\frac{2}{3}\right)_F^I \otimes \left(\mathbf{1}, \mathbf{2}, -1\right)_H^I + H.c.$$

$$\left(\mathbf{1}, \mathbf{2}, -1\right)_F \otimes \left(\mathbf{1}, \mathbf{1}, +2\right)_F^I \otimes \left(\mathbf{1}, \mathbf{2}, -1\right)_H^I + H.c.$$

$$\Rightarrow m_{d,\ell} \simeq \mathcal{O}(v_{EW})$$

$$\mathbf{15}_F \mathbf{15}_F \mathbf{15}_H + H.c. \supset$$

$$\left(\mathbf{3}, \mathbf{2}, +\frac{1}{3}\right)_F \otimes \left(\bar{\mathbf{3}}, \mathbf{1}, -\frac{4}{3}\right)_F^I \otimes \left(\mathbf{1}, \mathbf{2}, +1\right)_H + H.c.$$

$$\Rightarrow m_u \simeq \mathcal{O}(v_{EW})$$

The EW EFT = type-II 2HDM

The SU(6) Yukawa

$$* \bar{\mathbf{6}}_{\mathbf{F}}^{[\text{I}]} \bar{\mathbf{6}}_{\mathbf{F}}^{[\text{II}]} \mathbf{15}_{\mathbf{H}} + H.c. \supset$$

$$(\mathbf{1}, \mathbf{2}, -1)_{\mathbf{F}}^{[\text{I}]} \otimes (\mathbf{1}, \mathbf{1}, 0)_{\mathbf{F}}^{[\text{II}]} \otimes (\mathbf{1}, \mathbf{2}, +1)_{\mathbf{H}} + H.c.$$

$$\Rightarrow m_{\nu N'} \sim m_{\nu' N} \simeq \mathcal{O}(v_{\text{EW}})$$

$$* \text{Type-I seesaw: } (\nu, N') \mathcal{M} (\nu, N')^T \Rightarrow m_{\nu} \sim \frac{v_{\text{EW}}^2}{v_{331}}, m_{N'} \sim v_{331}$$

The SU(6) fermions

- * To obtain the spectrum: $N = \frac{1}{3} \text{diag}(-\mathbb{1}_3, +\mathbb{1}_3)$ for the SU(6) fundamental, and $Y = \text{diag}(\frac{1}{3} + 2N, \frac{1}{3} + 2N, -\frac{2}{3} + 2N)$ for the SU(3)_L fundamental.
- * In the SU(5) GUT: $\bar{\mathbf{5}}_{\text{F}} = d_R^c \oplus \ell_L$, $\mathbf{10}_{\text{F}} = q_L \oplus u_R^c \oplus e_R^c$
- * In the SU(6) GUT: $\bar{\mathbf{6}}_{\text{F}}^{\text{I}} \supset d_R^c \oplus \ell_L$, $\mathbf{15}_{\text{F}} \supset q_L \oplus u_R^c \oplus e_R^c$
- * Additional SU(6) fermions are vectorial: (D_L, D_R^c) , (N, N') , $(\ell'_L, \ell'_R{}^c)$