





Search for the invisible decays  
of the Higgs boson at 13 TeV  
(CMS experiment)

Vukasin Milosevic (IHEP Beijing)



THE 7TH CHINA LHC PHYSICS WORKSHOP  
(CLHCP2021)  
25-28.10.2021.



# Why the interest for the invisibly decaying Higgs?


- ◆ The "crown jewel" of the experimental particle physics:
  - ◆ Higgs boson was discovered by **ATLAS** and **CMS experiments** at CERN in 2012
  - ◆ All of the following measurements of its properties have been **consistent** with the **Standard Model (SM)**
  - ◆ Large uncertainties of these measurements can allow for physics beyond the SM



Why the interest in the invisible final state?

- ◆ According to the **SM**, the probability of  $\text{Br}(H \rightarrow 4\nu) \sim 0.1\%$ 
  - ◆ **Can represent a good way of testing for BSM physics!**
  - ◆ **Higgs boson** could be a **mediator between SM and DM** sector
  - ◆ Detection would require it to recoil against a visible system




Eur. Phys. J. C 79 (2019) 421




# Why the interest for the invisibly decaying Higgs?

- ◆ Higgs boson can take a role of **a mediator** between SM and DM particles:
  - ◆ Detection requires for the **Higgs to recoil against a visible system**
  - ◆ **Large** missing transverse energy ( $E_{T,miss}$ )

- ◆ **qqH**: Higgs boson is produced in a vector boson fusion topology (VBF)
- ◆ **VH**: Higgs boson production with a vector boson
- ◆ **ggH**: Higgs boson produced via gluon fusion.






# Where were we up until now? Early Run 2 combination


- ◆ The first combination measurement using Run 2 data was published **using the 2016 dataset**
- ◆ No significant deviation from the SM was reported:
  - ◆ The result of the measurement is expressed as the **95% CL upper limit on the  $B(H \rightarrow \text{inv.})$**
  - ◆ **This publication also included a first combination of Run 1 and 2015+2016 data**
    - ◆ Setting the  $B(H \rightarrow \text{inv.})$  limit to be at **0.19 (0.15)** for the observed (expected) value

## Physics Letters B 793 2019

35.9 fb<sup>-1</sup> (13 TeV)



4.9 fb<sup>-1</sup> (7 TeV) + 19.7 fb<sup>-1</sup> (8 TeV) + 38.2 fb<sup>-1</sup> (13 TeV)



# The Run 2 analysis strategy: Introduction

◆ **VBF production mode of the Higgs boson has a characteristic signature:**

- ◆ **Two jets** with a **large geometrical separation**
- ◆ **High dijet invariant mass** (a good way to control S/B)
- ◆ Represents a channel with the largest sensitivity

◆ **Main backgrounds:**



- ◆ **QCD and EWK produced V+jets** (where  $V = W/Z$ )
  - ◆ Irreducible when  $Z \rightarrow \nu\nu$  and  $W \rightarrow l\nu$
  - ◆ With the charged lepton being missed in the detection


◆ **Estimated through dedicated control regions in data (CR):**

- ◆ Z or W boson associated with the same dijet topology
- ◆ Resulting in four CRs separated by lepton flavour ( $e/\mu$ )

◆ **QCD multi jet processes - data driven estimation**


◆ **Contributions expected from diboson and top processes are estimated using simulation**






# The Run 2 analysis strategy: Two trigger approach


- ◆ Previous analysis strategy relied on purely  $E_{T,miss}$  trigger algorithms
  - ◆ **VBF topology targeting cuts were applied at the offline stage**
  - ◆ Imposed a high  $E_{T,miss}$  requirement:  $E_{T,miss} > 250$  GeV
    - ◆ **Froming the high- $E_{T,miss}$  (MTR) analysis category**
- ◆ The recent upgrades of the Level-1 trigger enabled complex variable manipulation at the first triggering stage:
  - ◆ **Brought in the possibility to target VBF topology**
  - ◆ **New VBF H L1 algorithm explored selection requirements ( $m_{jj}, p_T^{j1/2}$ )**
  - ◆ A follow up path at the second (HLT) stage:
    - ◆ Matched the selection logic of the L1 seed
    - ◆ Imposed  $E_{T,miss}$  cuts in order to reduce rate/timing
- ◆ **These additions led to a formation of a low- $E_{T,miss}$  analysis category (VTR)**
  - ◆ For  $160 < E_{T,miss} < 250$  GeV, where the VBF trigger performs better than the generic  $E_{T,miss}$  ones



# The Run 2 analysis strategy: Two trigger approach

- ◆ Previous analysis strategy relied on purely  $E_{T,miss}$  trigger algorithms
  - ◆ **VBF topology targeting cuts were applied at the offline stage**
  - ◆ Imposed a high  $E_{T,miss}$  requirement:  $E_{T,miss} > 250$  GeV
    - ◆ **Froming the high- $E_{T,miss}$  (MTR) analysis category**
  
- ◆ The recent upgrades of the Level-1 trigger enabled complex variable manipulation at the first triggering stage:
  - ◆ **Brought in the possibility to target VBF topology**
  - ◆ **New VBF H L1 algorithm explored selection requirements ( $m_{jj}, p_T^{j1/2}$ )**
  - ◆ A follow up path at the second (HLT) stage:
    - ◆ Matched the selection logic of the L1 seed
    - ◆ Imposed  $E_{T,miss}$  cuts in order to reduce rate/timing
  
- ◆ **These additions led to a formation of a low- $E_{T,miss}$  analysis category (VTR)**
  - ◆ For  $160 < E_{T,miss} < 250$  GeV, where the VBF trigger performs better than the generic  $E_{T,miss}$  ones





# Data Quality issues in Run 2 data

◆ During the 2017/18 data taking period, there were several detector related issues affecting this analysis:

## ◆ The HEM problem:


- ◆ A section of the HCAL endcap calorimeter was not functional during part of the 2018 era
- ◆ Inability to properly identify electrons / photons in the region  $\eta < -1.39$  and  $-1.6 < \phi < -0.9$ 
  - ◆ Mitigated by including specific selection criteria on electrons
- ◆ A high source of MET in SR in affected phi slice due to the lost tracks
  - ◆ Mitigated by placing a removal selection requirement the affected MET phi region

## ◆ HF noise:


- ◆ Appearance of jet “horns” (large data to MC discrepancy) for  $|\eta| \sim 3.0$
- ◆ HF jet shape variable selection introduced in order to battle it
- ◆ Required a data driven estimation of the multijet HF noise by inverting one selection requirement
  - ◆ Creating a noise enriched region, while estimating it through the use of a transfer factor

# The Run 2 analysis strategy: QCD multijet estimation


- ◆ A data-driven estimate is performed using events in which the  $E_{T,miss}$  arises from mismeasured jets:
  - ◆ A QCD multijet enriched region (CR) is formed by inverting one of the selection requirements
  - ◆ The low  $X = \min \Delta \phi(j, E_{T,miss})$  is used to define QCD CR
- ◆ Two steps are taken in order to obtain the QCD multijet contribution in the SR:
  - ◆ Shape of the dijet mass and its SR normalisation




## Normalisation



## Shape





# VBF analysis: Full Run 2 measurement - Results

◆ The VBF H(invisible) measurement using full Run 2 data - new result ([CMS-PAS-HIG-20-003](#))

◆ Improvements to the analysis strategy:

◆ Addition of new VBF H(invisible) topology targeting triggers

◆ Creating of a new, low  $E_T^{miss}$ , analysis category


◆ Addition of another ( $\gamma$ ) control region

◆ Helping with statistical precision of Z(l) CRs

◆ Brought ~20% gain in terms of signal sensitivity (when compared to 2016 strategy)


◆ No significant deviation from the SM was reported and the observed (expected) 95% CL upper limit was placed at:

◆  $B(H \rightarrow inv) = 0.17 (0.11)$



**CMS-PAS-HIG-20-003**

Category	Observed	Expected	1- $\sigma$ interval	2- $\sigma$ interval
2016	0.38	0.28	[0.20 – 0.40]	[0.15 – 0.53]
MTR 2017	0.25	0.19	[0.14 – 0.28]	[0.10 – 0.40]
MTR 2018	0.24	0.15	[0.11 – 0.22]	[0.08 – 0.31]
MTR 2017 2018	0.17	0.13	[0.09 – 0.18]	[0.07 – 0.25]
VTR 2017	0.57	0.45	[0.32 – 0.66]	[0.24 – 0.94]
VTR 2018	0.44	0.34	[0.24 – 0.49]	[0.18 – 0.69]
all 2017	0.24	0.18	[0.13 – 0.26]	[0.09 – 0.37]
all 2018	0.25	0.15	[0.10 – 0.21]	[0.08 – 0.29]
all 2017 2018	0.18	0.12	[0.08 – 0.17]	[0.06 – 0.23]
Run2	0.17	0.11	[0.08 – 0.15]	[0.06 – 0.21]



# VBF analysis: Full Run 2 measurement - Results

## ◆ The VBF H(invisible) measurement using full Run 2 data - new result ([CMS-PAS-HIG-20-003](#))

### ◆ Improvements to the analysis strategy:

#### ◆ Addition of new VBF H(invisible) topology targeting triggers

##### ◆ Creating of a new, low $E_T^{miss}$ , analysis category

#### ◆ Addition of another ( $\gamma$ ) control region

##### ◆ Helping with statistical precision of Z(l) CRs

#### ◆ Brought ~20% gain in terms of signal sensitivity (when compared to 2016 strategy)


## ◆ No significant deviation from the SM was reported and the observed (expected) 95% CL upper limit was placed at:

◆  $B(H \rightarrow inv) = 0.17 (0.11)$

## ◆ Reinterpretation of the results in terms of Higgs portal models:

### ◆ 90% CL upper limits on the spin-independent DM-nucleon scattering cross section

### ◆ Assuming a scalar or fermion DM candidate



# Summary

◆ These slides have summarised the recent studies of the invisible decays of the Higgs boson produced in a VBF topology from the CMS Collaboration:

◆ **First combination:**

◆ Focused on the **Run 1 + early Run 2 measurements**

◆ Sets a limit on  $B(H \rightarrow \text{inv})$  at 0.19 (0.15) for the observed (expected) value

◆ **Measurements using the full Run 2 dataset:**


◆ **Z(l)H(invisible):**  $B(H \rightarrow \text{inv}) = 0.29$  (0.25)

◆ **Mono V/mono Jet:**  $B(H \rightarrow \text{inv}) = 0.28$  (0.25)

◆ **VBF H(invisible):**  $B(H \rightarrow \text{inv}) = 0.17$  (0.11)


◆ **The last missing piece is the ttH full Run 2 search**

◆ Currently being prepared with spring conferences as its goal



**Thank you for your time!**

# BACKUP



# Selection requirements

Observable	MTR	VTR
Choice of pair	leading- $p_T$	leading- $M_{jj}$
Leading (subleading) jet	$p_T > 80$ (40) GeV, $ \eta  < 4.7$	$p_T > 140$ (70) GeV, $ \eta  < 4.7$
$p_T^{\text{miss}}$	$> 250$ GeV	$160 < p_T^{\text{miss}} \leq 250$
$\min\Delta\phi(\vec{p}_T^{\text{miss}}, \vec{p}_T^{\text{jet}})$	$> 0.5$ rad	$> 1.8$ rad
$ \Delta\phi_{jj} $	$< 1.5$ rad	$< 1.8$ rad
$M_{jj}$	$> 200$ GeV	$> 900$ GeV
$ p_T^{\text{miss}} - \text{calo}p_T^{\text{miss}}  / p_T^{\text{miss}}$		$< 0.5$
Leading/subleading jets $ \eta  < 2.5$		NHEF $< 0.8$ , CHEF $> 0.1$
HF-noise jet candidates		0 (see Table ??)
$\tau_h$ candidates		$N_{\tau_h} = 0$ with $p_T > 20$ GeV, $ \eta  < 2.3$
b quark jet		$N_{\text{jet}} = 0$ with $p_T > 20$ GeV, DeepCSV Medium
$\eta_{j1} \times \eta_{j2}$		$< 0$
$ \Delta\eta_{jj} $		$> 1$
Muons (electrons)		$N_{\mu,e} = 0$ with $p_T > 10$ GeV, $ \eta  < 2.4$ (2.5)
Photons		$N_{\gamma} = 0$ with $p_T > 15$ GeV, $ \eta  < 2.5$

# Uncertainties

Source of uncertainty	Ratios	Uncertainty vs. $M_{jj}$
Theoretical uncertainties		
Ren. scale V+jets (VBF)	$Z_{SR}/W_{SR}$	7.5%
Ren. scale V+jets (strong)	$Z_{SR}/W_{SR}$	8.2%
Fac. scale V+jets (VBF)	$Z_{SR}/W_{SR}$	1.5%
Fac. scale V+jets (strong)	$Z_{SR}/W_{SR}$	1.3%
PDF V+jets (strong)	$Z_{SR}/W_{SR}$	0%
PDF V+jets (VBF)	$Z_{SR}/W_{SR}$	0%
NLO EWK corr. V+jets (strong)	$Z_{SR}/W_{SR}$	0.5%
Ren. scale $\gamma$ +jets (VBF)	$Z_{SR}/\gamma_{CR}$	6–10%
Ren. scale $\gamma$ +jets (strong)	$Z_{SR}/\gamma_{CR}$	6–10%
Fac. scale $\gamma$ +jets (VBF)	$Z_{SR}/\gamma_{CR}$	2.5%
Fac. scale $\gamma$ +jets (strong)	$Z_{SR}/\gamma_{CR}$	2.5%
PDF $\gamma$ +jets (strong)	$Z_{SR}/\gamma_{CR}$	2.5%
PDF $\gamma$ +jets (VBF)	$Z_{SR}/\gamma_{CR}$	2.5%
NLO EWK corr. $\gamma$ +jets	$Z_{SR}/\gamma_{CR}$	3%
Experimental uncertainties		
Muon id. eff.	$Z_{CR}/Z_{SR}, W_{CR}/W_{SR}$	$\approx 0.5\%$ (per lepton)
Muon iso. eff.	$Z_{CR}/Z_{SR}, W_{CR}/W_{SR}$	$\approx 0.1\%$ (per lepton)
Electron reco. eff.	$Z_{CR}/Z_{SR}, W_{CR}/W_{SR}$	$\approx 0.5\%$ (per lepton)
Electron id. eff.	$Z_{CR}/Z_{SR}, W_{CR}/W_{SR}$	$\approx 1\%$ (per lepton)
Photon id. eff.	$Z_{SR}/\gamma$	5%
Muon veto	$Z_{SR}/W_{SR}, W_{CR}/W_{SR}$	$\approx 0.5\%$
Electron veto (reco)	$Z_{SR}/W_{SR}, W_{CR}/W_{SR}$	$\approx 1.5$ (1)% for VBF (strong)
Electron veto (id)	$Z_{SR}/W_{SR}, W_{CR}/W_{SR}$	$\approx 2.5$ (2)% for VBF (strong)
$\tau$ veto	$Z_{SR}/W_{SR}, W_{CR}/W_{SR}$	$\approx 1\%$
Electron trigger	$Z_{CR}/Z_{SR}, W_{CR}/W_{SR}$	$\approx 1\%$
$p_T^{\text{miss}}$ trigger	$Z_{CR}/Z_{SR}, W_{CR}/W_{SR}$	$\approx 2\%$
Photon trigger	$Z_{SR}/\gamma$	1%
<hr/>		
Jet energy scale	$Z_{SR}/W_{SR}$	1–2%
	$W_{CR}/W_{SR}$	1.0–1.5%
	$Z_{CR}/Z_{\nu\nu}$	1%
	$Z_{SR}/\gamma$	3%
<hr/>		
Jet energy resolution	$Z_{SR}/W_{SR}$	1.0–2.5%
	$W_{CR}/W_{SR}$	1.0–1.5%
	$Z_{CR}/Z_{SR}$	1%
	$Z_{SR}/\gamma$	1–4%

# Uncertainty breakdown

Group of systematic uncertainties	Observed impact on $\mathcal{B}(\text{H} \rightarrow \text{inv})$	Expected impact on $\mathcal{B}(\text{H} \rightarrow \text{inv})$
Theory	+0.026 -0.025	$\pm 0.024$
MC stat.	+0.024 -0.023	+0.023 -0.024
Triggers	+0.021 -0.022	$\pm 0.021$
Leptons/photons/b	+0.012 -0.011	+0.010 -0.011
QCD multijet mismodeling	$\pm 0.013$	$\pm 0.014$
Jet calibration	+0.010 -0.007	$\pm 0.007$
Lumi/PU	$\pm 0.005$	+0.004 -0.005
Other systematic uncertainties	+0.013 -0.010	$\pm 0.010$
Stat.	$\pm 0.029$	$\pm 0.030$