

Measurements of the Higgs boson inclusive and differential fiducial cross-sections in the diphoton decay channel with pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector



Gangcheng Lu¹, on behalf of the ATLAS collaboration

¹IHEP, CAS


CLHCP 2021, November 25st

Motivation


Among the possible studies of the properties of the Higgs boson, the measurement of its production cross-section in fiducial regions defined by detector acceptance minimizes the physics assumptions that is needed for extrapolation to the full phase space.

Besides, the measured cross-sections are not split by production process in favor of a production-inclusive measurement, hence further minimizing SM assumptions.

Despite the small branching ratio of the Higgs boson decay to two photons of $(2.27 \pm 0.01) \times 10^{-3}$, the excellent photon reconstruction and identification efficiency of the ATLAS detector enable extraction of the Higgs boson signal.




Fiducial phase space sensitive to specific Higgs production modes




Subsets of the diphoton baseline fiducial region are defined to provide phase-space regions sensitive to particular Higgs production modes

- ◆ VBF-enhanced: sensitive to VBF
- ◆ $N_{\text{lepton}} \geq 1$: sensitive to VH, ttH and tH
- ◆ High E_T^{miss} : sensitive to VH, ttH and BSM effects
- ◆ ttH-enhanced: sensitive to ttH and tH

Inclusive fiducial cross-section measurements



Matrix-inversion unfolding




Fiducial region	Measured [fb]	SM prediction [fb]	95% CL, Upper-limit [fb]	p-value
Diphoton	67	64	-	69%
VBF-enhanced	1.8	1.53	0.10	64%
$N_{\text{lepton}} \geq 1$	0.81	0.59	0.03	36%
High E_T^{miss}	0.28	0.302	0.017	93%
ttH-enhanced	0.53	0.60	0.05	79%
Total	132	126	7	-

Differential fiducial cross-section measurements


Multiple observables chosen in the measurement of differential and fiducial cross-sections

All differential measurements are limited by statistical uncertainties

Besides default MC, the measurements are compatible with many theoretical predictions (such as MATRIX+RadISH, RadISH+NNLOjet, ResBos2, GoSam).




Limits on the b - and c -quark Yukawa coupling



Probe κ_b and κ_c indirectly through the measured $p_T^{\gamma\gamma}$ spectrum, not limited by tagging efficiency of jets originating from b - and c -quarks.

Two fitting strategy studied:

- ◆ Only consider shape (shape)
- ◆ Consider also the normalization of cross-section times branching ratio (shape+XS+BR)



κ_b limits are comparable with direct searches^[4], while constraints on κ_c improve (no upper limits on κ_c in direct searches^[5])


fit-setup	κ_c	Observed 95% CL	Expected 95% CL
Shape-only	κ_c [-9.9, 13.3] κ_b [-2.7, 7.9]	[-10.0, 15.1] [-2.6, 7.6]	
Shape+Normalisation (with Branching ratio variations)	κ_c [-3.9, 3.9] κ_b [-1.3, -1.0] \cup [0.9, 1.3]	[-3.1, 3.1] [-1.2, -0.9] \cup [0.8, 1.2]	

Effective Field Theory (EFT) interpretation

In EFT approach, an effective Lagrangian is defined by \mathcal{L}_{SM} supplemented by additional dimension-6 operators:

$$\mathcal{L}_{\text{EFT}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{c_i}{\Lambda^2} O_i^{(6)}$$

Limits on the Wilson coefficients are obtained using a simultaneous fit to 5 measured cross-sections and their correlations: $p_T^{\gamma\gamma}$, N_{jets} , m_{jj} , $\Delta\phi_{jj}$ and $p_T^{\gamma\gamma}$



References

- [1] Measurements of the Higgs boson production and decay rates and coupling strengths using pp collision data at $\sqrt{s} = 7$ and 8 TeV in the ATLAS experiment, [arXiv:1507.04548](https://arxiv.org/abs/1507.04548)
- [2] Measurements of Higgs boson properties in the diphoton decay channel with 36fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector, [Phys.Rev.D 98 \(2018\) 052005](https://doi.org/10.1103/PhysRevD.98.052005)
- [3] Measurements of inclusive and differential cross sections in the $H \rightarrow ZZ^* \rightarrow 4l$ decay channel in pp collision at $\sqrt{s} = 13$ TeV with the ATLAS detector, [JHEP 10 \(2017\) 132](https://doi.org/10.1007/JHEP10(2017)132)
- [4] Combined measurements of Higgs boson couplings in proton-proton collisions at $\sqrt{s} = 13$ TeV, [Eur. Phys. J. C 79 \(2019\) 421](https://doi.org/10.1140/epjc/s10050-019-7121-0)
- [5] A search for the standard model Higgs boson decaying to charm quarks, [JHEP 03 \(2020\) 131](https://doi.org/10.1007/JHEP03(2020)131)



中国科学院高能物理研究所
Institute of High Energy Physics
Chinese Academy of Sciences