

The 10th International Workshop on Chiral Dynamics



Report of Contributions

Contribution ID: 0

Type: **not specified**

ChPT and BSM dark mesons

We plan to present the phenomenology and experimental searches of the dark pions, which are the lightest hadrons in a hidden sector confining gauge theory. Such a scenario arises in many extensions of the Standard Model (SM). We consider that the leading interactions between the light hidden sector quarks and the SM particles come from the mixing of the light hidden quarks with heavy electroweak doublet states through Higgs Yukawa couplings, so that the leading portals are the Z and Higgs bosons.

Summary

Based on arXiv:1906.02198 and an upcoming project.

Primary author: LI, Lingfeng (Brown University)

Presenter: LI, Lingfeng (Brown University)

Contribution ID: 1

Type: **Parallel-Few-Body Physics**

Fast & rigorous constraints on chiral three-nucleon forces from few-body observables

We explore the constraints on the three-nucleon force (3NF) of chiral effective field theory (χ EFT) that are provided by bound-state observables in the $A = 3$ and $A = 4$ sectors. Our statistically rigorous analysis incorporates experimental error, computational method uncertainty, and the uncertainty due to truncation of the χ EFT expansion at next-to-next-to-leading order. A consistent solution for the ^3H binding energy, the ^4He binding energy and radius, and the ^3H β -decay rate can only be obtained if χ EFT truncation errors are included in the analysis. All of these except the β -decay rate give essentially degenerate constraints on the 3NF low-energy constants, so it is crucial for estimating these parameters. We use eigenvector continuation for fast and accurate emulation of No-Core Shell Model calculations of the considered few-nucleon observables. This facilitates sampling of the posterior probability distribution, allowing us to also determine the distributions of the hyperparameters that quantify the truncation error. We find a χ EFT expansion parameter of $Q = 0.33 \pm 0.06$ for these observables.

Primary author: Prof. PHILLIPS, Daniel (Ohio University)

Co-authors: Prof. EKSTRÖM, Andreas (Chalmers University of Technology); Prof. FORSSÉN, Christian (Chalmers University of Technology); Prof. SVENSSON, Isak (Chalmers University of Technology); Dr MELENDEZ, Jordan (The Ohio State University); Prof. FURNSTAHL, Richard (The Ohio State University); Prof. WESOŁOWSKI, Sarah (Salisbury University)

Presenter: Prof. PHILLIPS, Daniel (Ohio University)

Contribution ID: 2

Type: **not specified**

Theoretical aspects of virtual states, bound states, and resonances in Hadron physics from Friedrichs model

Friedrichs model is a solvable model which is suitable to study the properties of the states that appear in scatterings. I will first review some theoretical aspects of the virtual state, bound state and resonances solutions for the Friedrichs model. Then some applications of this model to the hadron physics will be presented.

Primary author: 肖, 志广 (四川大学)

Co-author: Dr ZHOU, Zhi-Yong (Southeast University, China)

Presenter: 肖, 志广 (四川大学)

Contribution ID: 3

Type: **Parallel-Goldstone Boson**

Theoretical analysis of the doubly radiative decays

$$\eta^{(\prime)} \rightarrow \pi^0 \gamma \gamma \text{ and } \eta' \rightarrow \eta \gamma \gamma$$

The scalar and vector meson exchange contributions to the doubly radiative decays

$$\eta^{(\prime)} \rightarrow \pi^0 \gamma \gamma \text{ and } \eta' \rightarrow \eta \gamma \gamma$$

are analysed within the Linear Sigma Model and Vector Meson Dominance frameworks, respectively.

Predictions for the diphoton invariant mass spectra and the associated integrated branching ratios are given and compared with current available experimental data.

While a satisfactory description of the shape of the $\eta \rightarrow \pi^0 \gamma \gamma$ and $\eta' \rightarrow \pi^0 \gamma \gamma$ decay spectra is obtained, thus supporting the validity of the approach, the corresponding branching ratios cannot be reproduced simultaneously. A first theoretical prediction for the recently measured $\eta' \rightarrow \eta \gamma \gamma$ by the BESIII collaboration is also presented.

Primary author: Dr GONZÁLEZ-SOLÍS, Sergi (Indiana University)

Co-authors: Mr ROYO, Emilio (IFAE, Autonomous University of Barcelona); ESCRIBANO, Rafel (Universitat Autònoma de Barcelona); Dr JORA, Renate (National Institute of Physics and Nuclear Engineering, Romania)

Presenter: Dr GONZÁLEZ-SOLÍS, Sergi (Indiana University)

Contribution ID: 4

Type: **Parallel-Goldstone Boson**

A theoretical analysis of the semileptonic decays

$$\eta^{(\prime)} \rightarrow \pi^0 l^+ l^- \text{ and } \eta' \rightarrow \eta l^+ l^-$$

A theoretical analysis of the C -conserving semileptonic decays $\eta^{(\prime)} \rightarrow \pi^0 l^+ l^-$ and $\eta' \rightarrow \eta l^+ l^-$ ($l = e$ or μ) is carried out within the framework of the Vector Meson Dominance model. A phenomenological model is then used to parametrise the VMD couplings and numerical values are obtained from an optimisation fit. The signature of CP -violating operators from the SMEFT on experimental observables is also investigated and quantified for the $l = \mu$ case.

Primary authors: ROYO, Emilio (U); ESCRIBANO, Rafel (Universitat Autònoma de Barcelona)

Co-author: Dr SANCHEZ-PUERTAS, Pablo (Institut de Física d'Altes Energies (IFAE) and Barcelona Institute of Science and Technology (BIST))

Presenter: ROYO, Emilio (U)

Contribution ID: 5

Type: **Parallel-Goldstone Boson**

Dilaton chiral perturbation theory and applications

We review dilaton chiral perturbation theory (dChPT), the effective low-energy theory for the light sector of near-conformal, confining theories. dChPT provides a systematic expansion in both the fermion mass and the distance to the conformal window. It accounts for the pions and the light scalar, the approximate Nambu-Goldstone bosons for chiral and scale symmetry, respectively. A unique feature of dChPT is the existence of a large-mass regime in which the theory exhibits approximate hyperscaling, while the expansion nevertheless remains systematic.

We discuss applications to lattice data, presenting successes as well as directions for future work.

Primary author: Dr SHAMIR, Yigal (Tel Aviv University)

Co-author: Prof. GOLTERMAN, Maarten (San Francisco State University)

Presenter: Dr SHAMIR, Yigal (Tel Aviv University)

Contribution ID: 6

Type: **Parallel-Hadron Structure**

Chiral EFT of nucleons and pions in the presence of external gravitational field

Effective chiral Lagrangian of nucleons and pions in external gravitational field and the corresponding energy-momentum tensor will be considered. Gravitational form factors of the nucleon and their relation to internal forces will be discussed.

Primary author: Mr ALHARAZIN, Herzallah (Theoretical Hadron Physics)

Presenter: Mr ALHARAZIN, Herzallah (Theoretical Hadron Physics)

Contribution ID: 7

Type: **Parallel-Goldstone Boson**

Short-distance constraints in hadronic-light-by-light for the muon $g - 2$

We have made recent progress in studying the short-distance properties of the hadronic light-by-light contribution to the muon $g - 2$. The intermediate and short-distance part is a major contributor to the error of the theoretical prediction, see the white paper [arxiv:2006.04822, Physics Reports 887 (2020) 1-166]. We have recently shown that the massless quark-loop is the first term in a systematic expansion at short-distance [arxiv:1908.03331, Phys.Lett. B798 (2019) 134994]. This result already helped in the white paper in bringing down the error. Since then we have shown that both nonperturbative [arxiv:2008.13487, JHEP 10 (2020) 203] and the perturbative corrections [arxiv:2101.09169, JHEP 04 (2021) 240] are under control. The talk will describe these developments and how they fit in the total theoretical prediction for the muon $g - 2$.

Primary authors: RODRÍGUEZ, Antonio (IFIC (CSIC-Univ Valencia)); BIJNENS, Johan (L); Dr HERMANSSON TRUEDSSON, Nils (University of Berne)

Presenter: BIJNENS, Johan (L)

Contribution ID: 8

Type: **Parallel-Goldstone Boson**

NNLO Positivity Bounds on Chiral Perturbation Theory for a General Number of Flavours

We present positivity bounds, derived from the principles of analyticity, unitarity and crossing symmetry, that constrain the low-energy constants of chiral perturbation theory.

Bounds are produced for 2, 3 or more flavours with equal meson masses, up to and including next-to-next-to-leading order, using the second and higher derivatives of the amplitude. We enhance the bounds by using the most general isospin combinations possible (or higher-flavour counterparts thereof) and by analytically integrating the low-energy range of the amplitude. In addition, we present a powerful and general mathematical framework for efficiently managing large numbers of positivity bounds.

Primary authors: Dr ALVAREZ, Benjamin (Aix Marseille University); BIJNENS, Johan (L); SJÖ, Mattias (L)

Presenter: SJÖ, Mattias (L)

Contribution ID: 9

Type: **Parallel-Goldstone Boson**

Three-pion scattering in the Chiral Perturbation Theory

Within the framework of the massive $O(N)$ nonlinear sigma model extended to the next-to-leading order in the chiral counting (for $N = 3$ corresponding to the two(-quark)-flavor Chiral Perturbation Theory), we calculate the relativistic six-pion scattering amplitude at low energy up to and including terms $\mathcal{O}(p^4)$. Results for the pion mass, decay constant and the four-pion amplitude in the case of N (meson) flavors at $\mathcal{O}(p^4)$ are also presented.

Primary authors: BIJNENS, Johan (Lund University); HUSEK, Tomáš (Lund University (SE))

Presenter: HUSEK, Tomáš (Lund University (SE))

Contribution ID: 10

Type: **Parallel-Hadron Structure**

On QCD contribution to vacuum energy

In the framework of a two-loop order calculation for an effective field theory of scalar, vector and fermion fields interacting with the metric field we show that for the cosmological constant term which is fixed by the condition of vanishing vacuum energy, the graviton remains massless and there exists a self-consistent effective field theory of general relativity defined on a flat Minkowski background.

Next, using this result we address the issues of fine tuning of the strong interaction contribution to the vacuum energy and the compatibility of chiral symmetry in the light quark sector with the consistency of the effective field theory of general relativity in a flat Minkovski background.

Primary author: GEGELIA, Jambul (R)

Co-author: Prof. MEISSNER, Ulf-G. (Univ. Bonn & FZ Juelich)

Presenter: GEGELIA, Jambul (R)

Contribution ID: 11

Type: **Parallel-Goldstone Boson**

Dispersive analysis of the Primakoff reaction

$$\gamma K \rightarrow K \pi$$

We provide a dispersion-theoretical representation of the reaction amplitudes $\gamma K \rightarrow K \pi$ in all charge channels, based on modern pion-kaon P -wave phase shift input. Crossed-channel singularities are fixed from phenomenology as far as possible. We demonstrate how the subtraction constants can be matched to a low-energy theorem and radiative couplings of the $K^*(892)$ resonances, thereby providing a model-independent framework for future analyses of high-precision kaon Primakoff data.

Primary author: STAMEN, Dominik (B)

Co-authors: Dr KUBIS, Bastian (Bonn University); DAX, Maximilian (Bonn University)

Presenter: STAMEN, Dominik (B)

Contribution ID: 12

Type: **Parallel-Few-Body Physics**

Extracting few-body matrix elements from lattice QCD

In this talk I will report on a recent lattice QCD determination of few-body matrix elements made by the NPLQCD Collaboration. One of them is the fraction of the longitudinal momentum of ^3He that is carried by the isovector combination of u and d quarks. The ratio of this combination to that in the constituent nucleons is extracted in a calculation with quark masses corresponding to a pion mass of 800 MeV, and then extrapolated to the physical point, showing that it is consistent with, and significantly more precise than, determinations from global nuclear parton distribution function fits. The other quantity studied is the tritium axial charge, related to the Gamow-Teller matrix element, which is also computed with heavier-than-physical quark masses and then extrapolated and compared with the phenomenological value. The importance of this last study is that it demonstrates that QCD can explain the quenching of the triton axial charge.

Primary author: ILLA, Marc (U)

Presenter: ILLA, Marc (U)

Contribution ID: 13

Type: **not specified**

Most charming dibaryon near unitarity from lattice QCD

We present a first study on a pair of triply charmed baryons, $\Omega_{ccc}\Omega_{ccc}$ in the 1S_0 channel (the most charming dibaryon), on the basis of the HAL QCD method.

The calculations are performed on the $(2+1)$ -flavor lattice QCD configurations with nearly physical light-quark masses and physical charm-quark mass.

We show that the system with the Coulomb repulsion taking into account the charge form factor of Ω_{ccc} leads to the scattering length $a_0^C \simeq -19$ fm and the effective range $r_{\text{eff}}^C \simeq 0.45$ fm, which indicates $\Omega_{ccc}\Omega_{ccc}$ is located in the unitary regime.

Primary author: LYU, Yan (school of physics Peking University)

Co-authors: TONG, Hui (school of physics Peking University); AOKI, Sinya (YITP, Kyoto University); Dr MIYAMOTO, Takaya (RIKEN); Prof. DOI, Takumi (RIKEN); Dr SUGIURA, Takuya (RIKEN); Prof. HATSUDA, Tetsuo (RIKEN); Prof. 孟, 杰 (北京大学)

Presenter: LYU, Yan (school of physics Peking University)

Contribution ID: 14

Type: **Parallel-Hadron Structure**

Structure-dependent electromagnetic finite-size effects

In this talk we present a relativistic and model-independent method to analytically derive electromagnetic finite-size effects beyond the point-like approximation. Structure-dependence appears in terms of physical form-factors and derivatives thereof. The values of these physical quantities can be taken either from experimental measurements or auxiliary lattice calculations. We first apply our method to the meson mass, and then to leptonic decays of pions and kaons. Knowledge of the latter allows for improved numerical control in extractions of the CKM-matrix element ratio $|V_{us}/V_{ud}|$ from lattice QCD+QED.

Primary author: Dr HERMANSSON-TRUEDSSON, Nils (University of Bern)

Co-authors: Dr PORTELLI, Antonin (University of Edinburgh); Dr DI CARLO, Matteo (University of Edinburgh); Dr HANSEN, Maxwell T. (University of Edinburgh)

Presenter: Dr HERMANSSON-TRUEDSSON, Nils (University of Bern)

Contribution ID: 15

Type: **Parallel-Few-Body Physics**

Nuclear structure corrections in muonic atoms from chiral effective field theory

Precision spectroscopic measurement in muonic atoms require precision theoretical calculations. While quantum electrodynamics effects are very well known, nuclear structure corrections are presently the largest source of uncertainty and consequently the bottle-neck for fully exploiting the experimental precision in extracting nuclear radii.

Utilizing techniques and methods developed in few-body nuclear physics, we have been able to provide the so far most precise determination of nuclear structure corrections to the Lamb shift.

I will present our recent calculations for light muonic atoms, where we use chiral effective field theory potentials and perform a study

of the uncertainties coming from the order-by-order chiral expansion.

Primary author: BACCA, Sonia (Johannes Gutenberg University)

Co-author: LI MULI, Simone (Johannes Gutenberg University)

Presenters: LI MULI, Simone (Johannes Gutenberg University); BACCA, Sonia (Johannes Gutenberg University)

Contribution ID: 16

Type: **not specified**

Strangeness $S=-3$ and -4 baryon-baryon interactions in chiral effective field theory

The Jülich-Bonn-Munich Collaboration has applied chiral effective field theory to investigate the baryon-baryon interaction involving hyperons. These studies, performed so far up to next-to-leading order (NLO) in the chiral expansion, have shown that for the strangeness $S = -1$ (ΛN , ΣN) and $S = -2$ ($\Lambda\Lambda$, ΞN) sectors a consistent and satisfactory description of the available scattering data and experimental constraints can be achieved within the assumption of broken $SU(3)$ flavor symmetry. In addition, applications of the resulting potentials in bound-state calculations for light hypernuclei led to results close to the empirical values.

In the present contribution we discuss a possible extension of this approach to strangeness $S = -3$ and $S = -4$ baryon-baryon systems where empirical information is rather scarce. Specifically we address the question in how far measurements of two-body correlation functions in heavy-ion collisions and/or in high energetic proton-proton scattering can be used to pin down the interaction in channels like $\Xi\Lambda$ or $\Xi\Xi$, at least on a semi-quantitative level.

Primary author: Dr HAIDENBAUER, Johann (Forschungszentrum Juelich GmbH)

Co-author: Prof. MEISSNER, Ulf-G. (Univ. Bonn & FZ Juelich)

Presenter: Dr HAIDENBAUER, Johann (Forschungszentrum Juelich GmbH)

Contribution ID: 17

Type: **Plenary**

Precision calculations of charge radii of light nuclei

We present a high-precision calculation of the deuteron, triton, ^3He and ^4He charge form factors based on the latest two- and three-nucleon forces, and charge density operators derived up through the fifth order in the chiral effective field theory.

We predict the values of the structure radius and the quadrupole moment of the deuteron, the ^4He charge radius, and the isoscalar combination of the 3N charge radii $(r^2(^3\text{H}) + 2r^2(^3\text{He}))/3$. A comprehensive and systematic analysis of various sources of uncertainty in all our predictions is performed.

Using the predicted value for the deuteron structure radius together with the very accurate atomic data for the difference of the deuteron and proton charge radii we extract the charge radius of the neutron.

Finally, using the predicted isoscalar combination of the 3N charge radii and preliminary experimental data on the ^3He charge radius we estimate the charge radius of the triton.

Primary authors: NOGGA, Andreas (Forschungszentrum Jülich); FILIN, Arseniy (Ruhr University Bochum); KÖRBER, Christopher (Ruhr University Bochum); MÖLLER, Daniel (Ruhr University Bochum); EPELBAUM, Evgeny (Ruhr University Bochum); KREBS, Hermann (Ruhr University Bochum); REINERT, Patrick (Ruhr University Bochum); BARU, Vadim (Ruhr University Bochum)

Presenter: FILIN, Arseniy (Ruhr University Bochum)

Contribution ID: 18

Type: **Parallel-Few-Body Physics**

Explicit renormalization of the nucleon-nucleon interaction in chiral EFT and non-perturbative effects.

Nucleon-nucleon interaction

is studied within chiral effective field theory with a finite cutoff at next-to-leading order in the chiral expansion.

The leading order interaction is resummed non-perturbatively, whereas the next-to-leading-order terms are taken into account in a perturbative manner. Explicit renormalizability of such a scheme is proven in certain important cases.

In particular, it is verified whether the power-counting breaking terms originating from the integration regions with large momenta can be absorbed by the renormalization of the low energy constants.

The importance of non-perturbative effects is analyzed in detail.

Primary author: Dr GASPARYAN, Ashot (Ruhr University of Bochum)

Co-author: Prof. EPELBAUM, Evgeny (Ruhr University of Bochum)

Presenter: Dr GASPARYAN, Ashot (Ruhr University of Bochum)

Contribution ID: 19

Type: **Parallel-Goldstone Boson**

Patterns of C - and CP -violation in hadronic η and η' three-body decays

We construct hadronic amplitudes for the three-body decays $\eta^{(\prime)} \rightarrow \pi^+\pi^-\pi^0$ and $\eta' \rightarrow \eta\pi^+\pi^-$ in a non-perturbative fashion, allowing for C - and CP -violating asymmetries in the $\pi^+\pi^-$ distributions.

These amplitudes are consistent with the constraints of analyticity and unitarity. We find that the currently most accurate Dalitz-plot distributions taken by the KLOE-2 and BESIII collaborations confine the patterns of these asymmetries to a relative per mille and per cent level, respectively. Our dispersive representation allows us to extract the individual coupling strengths of the C - and CP -violating contributions arising from effective isoscalar and isotensor operators in $\eta^{(\prime)} \rightarrow \pi^+\pi^-\pi^0$ and an effective isovector operator in $\eta' \rightarrow \eta\pi^+\pi^-$, while the strongly different sensitivities to these operators can be understood from chiral power counting arguments.

Primary author: AKDAG, Hakan (Bonn University)

Presenter: AKDAG, Hakan (Bonn University)

Contribution ID: 20

Type: **Plenary**

Study of deuteron-dark matter scattering within the χ EFT framework

In this work we studied the deuteron-WIMP scattering, assuming the latter to be a Dirac particle. In order to describe the WIMP-nucleon interaction we used the χ EFT framework, an EFT based on chiral symmetry which has pions and nucleons as effective degrees of freedom.

The Lagrangian interaction terms have been obtained by placing the WIMP as an external source in the QCD Lagrangian. Therefore, considering only interactions invariant under parity, charge conjugation and time-reversal symmetries, we have examined five WIMP and quarks interaction types (scalar, pseudoscalar, vector, axial and tensor).

We have applied this program to study the nuclear response to DM scattering up to second chiral order and finally to calculate the rate of WIMP scattering in a target of liquid deuterium. With respect to other calculations in the literature, we have included systematically the contributions of two-body currents and treated quark-WIMP vertex of tensor type. We plan to apply this formalism to study also helium-DM scattering since liquid helium has been proposed as a possible target for a new direct detection experiment.

Primary author: FILANDRI, Elena (U)

Co-author: Prof. VIVIANI, Michele (University of Pisa & INFN)

Presenter: FILANDRI, Elena (U)

Contribution ID: 21

Type: **not specified**

Two-particle scattering from finite-volume quantization conditions using the plane wave basis

We propose an alternative approach to Lüscher's formula for extracting two-body scattering phase shifts from finite volume spectra with no reliance on the partial wave expansion. We use an effective-field-theory-based Hamiltonian method in the plane wave basis and decompose the corresponding matrix elements of operators into irreducible representations of the relevant point groups. The proposed approach allows one to benefit from the knowledge of the long-range interaction and avoids complications from partial wave mixing in a finite volume. We consider spin-singlet channels in the two-nucleon system and pion-pion scattering in the ρ -meson channel in the rest and moving frames to illustrate the method for non-relativistic and relativistic systems, respectively. For the two-nucleon system, the long-range interaction due to the one-pion exchange is found to make the single-channel Lüscher formula unreliable at the physical pion mass. For S-wave dominated states, the single-channel Lüscher method suffers from significant finite-volume artifacts for a $L = 3\text{ fm}$ box, but it works well for boxes with $L > 5\text{ fm}$. However, for P-wave dominated states, significant partial wave mixing effects prevent the application of the single-channel Lüscher formula regardless of the box size (except for the near-threshold region). Using a toy model to generate synthetic data for finite-volume energies, we show that our effective-field-theory-based approach in the plane wave basis is capable of a reliable extraction of the phase shifts. For pion-pion scattering, we employ a phenomenological model to fit lattice QCD results at the physical pion mass. The extracted P-wave phase shifts are found to be in a good agreement with the experimental results.

Primary author: MENG, Lu (Ruhr University Bochum)

Co-author: Prof. EPELBAUM, Evgeny (Ruhr University Bochum)

Presenter: MENG, Lu (Ruhr University Bochum)

Contribution ID: 22

Type: **Parallel-Hadron Structure**

New insights into the nucleon form factor and proton radius from dispersive analysis

We present a combined analysis of the electromagnetic form factors of the nucleon in the space- and timelike regions using dispersion theory. Our framework provides a consistent description of the experimental data over the full range of momentum transfer, in line with the strictures from analyticity and unitarity. The statistical uncertainties of the extracted form factors are estimated using the bootstrap method, while systematic errors are determined from variations of the spectral functions. We also perform a high-precision extraction of the nucleon radii and find good agreement with previous analyses of spacelike data alone. For the proton charge radius, we find

$$r_E^p = 0.840^{+0.003+0.002}_{-0.002-0.002} \text{ fm},$$

where the first error is statistical and the second one is systematic.

The Zemach radius and third moment are in agreement with Lamb shift measurements and hyperfine splittings.

The combined data set of space- and timelike data disfavors a zero crossing of $\mu_p G_E^p / G_M^p$ in the spacelike region.

Finally, we discuss the status and perspectives of modulus and phase of the form factors in the timelike region in the context of future experiments as well as the onset of perturbative QCD.

Primary author: Dr LIN, Yonghui (HISKP, Bonn University)

Co-authors: Prof. HAMMER, Hans-Werner (Technical University of Darmstadt); Prof. MEISSNER, Ulf-G. (Bonn University)

Presenter: Dr LIN, Yonghui (HISKP, Bonn University)

Contribution ID: 23

Type: **not specified**

Neutron-antineutron oscillation and the deuteron lifetime

A framework based on chiral effective field theory to treat nuclear physics of baryon-number violation by two units is discussed. Neutron-antineutron oscillation and its relation to the deuteron lifetime is used as an application to illustrate this framework. The emphasis is given to how a consistent power counting is built and what statements can be drawn out of it.

Primary author: Dr LONG, Bingwei (Sichuan University)

Presenter: Dr LONG, Bingwei (Sichuan University)

Contribution ID: 24

Type: **Parallel-Goldstone Boson**

Dispersive approach to strong three-body decays of η and η' mesons

I present a dispersion-theoretical analysis of the strong three-body decays in the η and η' sector. Based on analyticity and unitarity the dispersion relations resum the leading $\pi\pi$ - and $\pi\eta$ -rescattering effects to all orders, with the resulting amplitudes being solely dependent on the respective two-body scattering phase shifts. Due to the small available phase space only three decay channels are physically allowed: the two isospin breaking transitions $\eta/\eta' \rightarrow 3\pi$ and the isospin conserving transition $\eta' \rightarrow \pi\pi\eta$. Combining our dispersive representation of $\eta \rightarrow 3\pi$ with ChPT constraints, a fit to the high-statistics A2 and KLOE-2 Dalitz-plot distributions enables us to extract the light-quark mass double ratio Q . We update our previous analysis of $\eta' \rightarrow \pi\pi\eta$ by performing fits to new high-statistics measurements from A2 and BESIII. For the process $\eta' \rightarrow 3\pi$, we include both elastic two-pion rescattering corrections to the isospin-breaking decay amplitude as well as the inelastic effect due to an isospin-conserving decay $\eta' \rightarrow \pi\pi\eta$ and subsequent isospin-breaking $\pi\eta \rightarrow \pi\pi$ rescattering. In this way, the inelastic contribution to the unitarity relation connects all three decays. The resulting dispersive description of $\eta' \rightarrow 3\pi$ is fitted to the first available measurements of the Dalitz-plot distributions from BESIII.

Primary author: ISKEN, Tobias (G)

Co-authors: KUPSC, Andrzej (Uppsala University); KUBIS, Bastian (Bonn University); STOFFER, Peter (Zürich University)

Presenter: ISKEN, Tobias (G)

Contribution ID: 25

Type: **Parallel-Goldstone Boson**

A review on theoretical calculations of π^0 lifetime

The neutral pi meson has a prominent role among all hadrons as being the lightest of them. The most important decay mode is the process into two photons, and it saturates the π^0 decay width by almost 99%. This is a fundamental process also from the theoretical point of view, connected with the famous chiral anomaly. The overview of the theoretical calculations will be presented together with a comparison with experimental measurements. Impact on other processes will be briefly covered.

Primary author: KAMPF, Karol (C)**Presenter:** KAMPF, Karol (C)

Contribution ID: 26

Type: **Parallel-Goldstone Boson**

Review of light quark mass ratio determination via $\eta \rightarrow 3\pi$ decay

In this talk, we will show how $\eta \rightarrow 3\pi$ is the golden channel for determining the light quark mass ratio from experiment. We will review the different extractions and uncertainty assessment. The results will be compared with lattice QCD determinations and the opportunities for further improvement will be discussed.

Primary author: PASSEMAR, Emilie (Indiana University/JLab)

Presenter: PASSEMAR, Emilie (Indiana University/JLab)

Contribution ID: 27

Type: **not specified**

Two-loop calculation of the nucleon self-energy

The nucleon-self energy

is calculated in SU(2) covariant chiral perturbation theory to study the pion mass dependence of the nucleon mass up to chiral order $O(q^6)$, i.e., including all two-loop diagrams.

Applying an algorithm from Tarasov (1997), the mathematical expressions of the diagrams were expressed by a small set of (scalar) master integrals.

These master integrals are solved as a chiral expansion in d dimensions, using the strategy of regions to differentiate the infrared singular and regular part.

Extended on-mass-shell renormalization is applied, making the renormalized expressions consistent with the power counting.

Primary author: CONRAD, Nils Dirk (I)

Presenter: CONRAD, Nils Dirk (I)

Contribution ID: 28

Type: **Plenary**

Nuclear forces in a manifestly Lorentz-invariant formulation of chiral EFT

We develop a systematic approach for chiral nuclear forces by applying time-ordered perturbation theory to manifestly Lorentz-invariant formulation of chiral effective field theory. The effective potential and the scattering equation (Kadyshevsky equation) are obtained within the same framework. Restricting the non-perturbative treatment to the (non-singular) leading order potential and assuming the validity of perturbation theory for higher-order interactions, one can systematically remove all divergences from the amplitude and, therefore, employ arbitrarily large values of the cutoff. Alternatively, the full effective potential can be treated non-perturbatively by taking the cutoff of the order of the hard scale. The milder UV behavior then offers a larger flexibility regarding admissible cutoff values.

Along this line, we have studied chiral two-body force up to next-to-next-to-leading order and achieved a rather good description of phase shifts and deuteron properties, although it is computationally more demanding as compared to its non-relativistic counterpart. We expect that our approach should lead to a better description of systems with larger numbers of nucleons.

Primary author: Dr REN, Xiu-Lei (Johannes Gutenberg-Universität Mainz)

Co-authors: Prof. EPELBAUM, Evgeny (Ruhr-Universität Bochum); Dr GEGELIA, Jambul (Ruhr-Universität Bochum)

Presenter: Dr REN, Xiu-Lei (Johannes Gutenberg-Universität Mainz)

Contribution ID: 29

Type: **not specified**

Analysis of Tcc including chiral dynamics and three-body cut

A coupled-channel approach is applied to the charged tetraquark state T_{cc} recently discovered by the LHCb Collaboration with special attention paid to the three-body dynamics due to the finite life time of the D^* . The low-energy expansion of the D^*D scattering amplitude is performed and the low-energy parameters (the scattering length and the effective range) are extracted. The compositeness parameter of the T_{cc} is found to be close to unity, which implies that the T_{cc} is a hadronic molecule, generated by the interactions in the $D^{*+}D^0$ and $D^{*0}D^+$ channels. With help of heavy-quark spin symmetry, an isoscalar D^*D^* molecular partner of the T_{cc} with $J^P = 1^+$ is predicted under the assumption that the $DD^*-D^*D^*$ coupled-channel effects can be neglected.

Primary author: Dr DU, Meng-Lin (IFIC, Valencia U.)

Co-authors: Prof. NEFEDIEV, Alexey (P.N. Lebedev Physical Institute of the Russian Academy of Sciences); FILIN, Arseniy (University Bochum); Prof. HANHART, Christoph (Forschungszentrum Juelich); Prof. GUO, Feng-Kun (ITP, Chinese Academy of Science); Prof. NIEVES, Juan (IFIC, Valencia U.); Prof. WANG, Qian (South China Normal University.); BARU, Vadim (University Bochum); DONG, Xiang-Kun (ITP, Chinese Academy of Science)

Presenter: Dr DU, Meng-Lin (IFIC, Valencia U.)

Contribution ID: 30

Type: **not specified**

Pion-mass dependence of $\gamma^{(*)}\pi \rightarrow \pi\pi$ and $\pi\pi \rightarrow \pi\pi$

Recent years have witnessed drastic progress in the computation of hadronic scattering via lattice QCD. Nevertheless, computations are often performed at unphysically high pion masses. Here, we extrapolate lattice results of the anomalous $\gamma^{(*)}\pi \rightarrow \pi\pi$ scattering to the physical pion mass via a dispersive framework, extract the associated chiral anomaly and the radiative coupling of the ρ -resonance to $\pi\gamma$. In addition, we confront unitarized 1- and 2-loop ChPT with $\pi\pi \rightarrow \pi\pi$ data, determine the characteristics of the ρ and assess the chiral convergence. The talk is based on arXiv:2110.11372 and arXiv:2009.04479.

Primary author: NIEHUS, Malwin (HISKP)

Co-authors: Dr KUBIS, Bastian (Bonn University); Dr RUIZ DE ELVIRA, Jacobo (HISKP-Bonn); HOFERICHTER, Martin (University of Bern)

Presenter: NIEHUS, Malwin (HISKP)

Contribution ID: 31

Type: **Parallel-Few-Body Physics**

Review of the experimental activity at RIKEN to explore the three-nucleon interactions

Understanding the nuclear properties from bare nuclear forces is one of the main topics in nuclear physics. The importance of three-nucleon forces (3NFs), which appear when more than two nucleons interact, has been indicated in various nuclear phenomena, such as few-nucleon scattering, nuclei binding energies, and state equation of nuclear matter.

Nucleon-deuteron (Nd) scattering, where numerically exact solutions of the corresponding Faddeev equations for any 2N- and 3N-forces are feasible, offers a good opportunity to study dynamical aspects of 3NFs, that are momentum, spin, and isospin dependences. It provides not only cross sections but also a variety of spin observables at different incident nucleon energies.

Signatures of 3NF effects in the Nd elastic scattering was pointed out for the first time by Wita{\l}a et al., in 1998 [1]. Clear signals from 3NFs were found around the cross section minimum occurring at the center of mass angle around 120 degrees for incident energies above 70 MeV/nucleon. Since then, we have performed the measurements of elastic deuteron-proton scattering with a RIKEN polarized deuteron beam, providing precise data of the cross section (70 and 135 MeV/nucleon), all deuteron analyzing powers (70, 100, 135, 190, 250, 300 MeV/nucleon), and polarization transfer coefficients (135 MeV/nucleon).

In the workshop, I review the experimental activity at RIKEN to explore the three-nucleon interactions and touch upon our future plan.

[1] H. Wita{\l}a et al., Phys. Rev. Lett. 81, 1183 (1998).

Primary author: SEKIGUCHI, Kimiko (Tohoku University)

Presenter: SEKIGUCHI, Kimiko (Tohoku University)

Contribution ID: 32

Type: **Parallel-Few-Body Physics**

Renormalization of CP-violating nuclear forces

Electric dipole moments of nuclei, diamagnetic atoms, and certain molecules are induced by CP-violating nuclear forces. Naive dimensional analysis predicts these forces to be dominated by long-range one-pionexchange processes with short-range forces entering only at next-to-next-to-leading order in the chiral expansion. Based on renormalization arguments we argue that a consistent picture of CP-violating nuclear forces requires a leading-order short-distance operator contributing to $1S0 - 3P0$ transitions due to the attractive and singular nature of the strong tensor force in the $3P0$ channel. The short-distance operator leads to $O(1)$ corrections to static and oscillating, relevant for axion searches, electric dipole moments. We discuss strategies how the finite part of the associated low-energy constant can be determined in the case of CP violation from the QCD θ^- term by the connection to charge-symmetry violation in nuclear systems.

Primary author: Mr SHAIN PORUVELIL, sachin (University of Massachusetts, Amherst)

Co-authors: Dr GNECH, Alex (Jefferson Lab, USA); Prof. JORDY, de Vries (University of Amsterdam, The Netherlands and Nikhef, The Netherlands)

Presenter: Mr SHAIN PORUVELIL, sachin (University of Massachusetts, Amherst)

Contribution ID: 33

Type: **Parallel-Goldstone Boson**

Measurements of charged and neutral pion electromagnetic polarizabilities at GlueX

Electromagnetic polarizabilities are fundamental properties of composite systems such as molecules, atoms, nuclei and hadrons. Measurements of hadron polarizabilities can test effective field theories, dispersion theories, and lattice calculations, with the charged pion polarizability providing a test of fundamental symmetries at leading order. Significant progress has been made in measurements of nucleon polarizabilities, with uncertainties at $\approx \pm 0.4 \times 10^{-4} e fm^3$ for the proton. However experimental constraints on the charged and neutral pion polarizabilities (CPP and NPP) are much weaker, $\approx \pm 2 \times 10^{-4} e fm^3$ for the π^+ and no measurement for the π^0 . The CPP and NPP experiments at GlueX will utilize a new technique to measure pion polarizability, Primakoff photo-production of $\pi^+\pi^-$ and $\pi^0\pi^0$ pairs on a high Z target. Details of the experimental setup and technique will be presented in the talk, including the commissioning of a muon detection system constructed for the measurement. The CPP and NPP experiments are currently scheduled to run at JLab in mid-2022.

Primary author: MISKIMEN, Rory (U)

Presenter: MISKIMEN, Rory (U)

Contribution ID: 34

Type: **Parallel-Goldstone Boson**

OPE of Green functions and its phenomenological applications

In this talk, we deliver an overview of our most recent results on the operator product expansion of the three-point correlators of chiral currents. Moreover, we present some phenomenological applications of these results within the context of the odd-intrinsic parity sector, such as the decays $\rho \rightarrow \gamma\pi$, $\omega \rightarrow \gamma\pi$ and the $g - 2$

Primary author: Mr KADAVY, Tomas (Charles University)

Co-authors: Dr NOVOTNY, Jiri (Charles University); Dr KAMPF, Karol (Charles University)

Presenter: Mr KADAVY, Tomas (Charles University)

Contribution ID: 35

Type: **Parallel-Few-Body Physics**

Preliminary results for elastic nucleon-pion scattering amplitudes from lattice QCD

The prospects and difficulties of computing nucleon-pion scattering amplitudes from lattice QCD simulations are illustrated with high-statistics results on a single ensemble of gauge field configurations with dynamical up, down, and strange quarks and a pion mass $m_\pi = 200\text{MeV}$. The stochastic-LapH approach to quark propagation enables an efficient computation of all required correlation functions, and a good statistical precision is achieved for the $I = 3/2$ amplitudes. The $I = 1/2$ channel is considerably more difficult, complicating direct lattice determination of both scattering lengths.

Primary author: BULAVA, John (D)

Presenter: BULAVA, John (D)

Contribution ID: 36

Type: **Parallel-Hadron Structure**

Low-energy Kbar N interaction from high-energy nuclear collisions

The two-particle momentum correlation function from high-energy nuclear collisions is beginning to be used to study hadron-hadron interaction. In this talk, the K^-p correlation function is discussed with employing the realistic KbarN-piSigma-piLambda coupled-channel potential based on the chiral SU(3) dynamics. With the reasonable source function parameters, the theoretical calculations well reproduce the recent ALICE data from various source sizes. The coupled-channel effect and source size dependence of the correlation function are investigated in detail. Finally, the application of femtoscopy to the other hadron systems are discussed.

Primary author: KAMIYA, Yuki (Helmholtz-Institut für Strahlen- und Kernphysik)

Presenter: KAMIYA, Yuki (Helmholtz-Institut für Strahlen- und Kernphysik)

Contribution ID: 37

Type: **Parallel-Hadron Structure**

Determining the nucleon mass and sigma term from lattice QCD

We report a preliminary, percent-level determination of the nucleon mass M_N and a roughly 5%-level determination of the sigma term $\sigma_{\pi N}$ from lattice QCD. We find that our M_N extrapolation to the physical point agrees with the PDG average. Next we review the significance of $\sigma_{\pi N}$ for direct dark matter searches, and we explore the sensitivity of this observable over choice of chiral models. For our lattice calculations, we employ Möbius domain wall fermions on $N_F = 2 + 1 + 1$ dynamically, highly-improved staggered quark fermions. We include five pion masses, ranging from 130 MeV to 350 MeV; four lattice spacings, ranging from 0.06 fm to 0.15 fm; and multiple lattice volumes.

Primary author: Mr MILLER, Nolan (University of North Carolina, Chapel Hill)

Co-authors: Dr MEYER, Aaron (Brookhaven National Laboratory); Dr NICHOLSON, Amy (University of North Carolina, Chapel Hill); Dr WALKER-LOUD, André (Lawrence Berkeley National Laboratory); Dr DRISCHLER, Christian (University of California, Berkeley); Dr KÖRBER, Christopher (Ruhr-University Bochum); Dr HOWARTH, Dean (Lawrence Livermore National Laboratory); Mr BRADLEY, Grant (Brown University); Dr MONGE, Henry (University of Costa Rica); Mr LAZAROW, Malcolm (University of California, Berkeley)

Presenter: Mr MILLER, Nolan (University of North Carolina, Chapel Hill)

Contribution ID: 38

Type: **not specified**

The radiative decay width measurement of the η -meson at GlueX.

The PrimEx-eta experiment at Jefferson Lab is aiming to measure the radiative decay width of the η -meson with a 3.2% precision. This projected accuracy will ameliorate the calculation of all η -meson partial decay widths and in particular the hadronic contribution to the muon magnetic moment from lattice QCD. It will provide critical input to determine the η - η' mixing angle and the light quark mass ratio model independently. The first run was performed with the GlueX experimental setup in Hall D in 2019 and the second run in fall 2021. We will discuss the status of the experiment and how the radiative decay width is extracted via the Primakoff effect from the η -meson photoproduction off a helium nucleus. We will also discuss the measurement of Compton scattering off an atomic electron, which is used to control experimental systematics, including the detection efficiency, the luminosity, and the measurement stability over time.

Primary author: JAEGLER, Igal (J)

Presenter: JAEGLER, Igal (J)

Contribution ID: 39

Type: **Parallel-Few-Body Physics**

Fluctuations and phases in baryonic matter

The phase structure of baryonic matter is investigated with focus on the role of fluctuations beyond the mean-field approximation. The prototype test case studied is the chiral nucleon-meson model, with added comments on the chiral quark-meson model. Applications to nuclear matter include the liquid-gas phase transition. Extensions to high baryon densities are performed for both nuclear and neutron matter. The role of vacuum fluctuations is systematically explored. It is pointed out that such fluctuations tend to stabilize the hadronic phase characterized by spontaneously broken chiral symmetry, shifting the chiral restoration transition to very high densities. This stabilization effect is shown to be further enhanced by additional dynamical fluctuations treated with functional renormalisation group methods.

Primary author: BRANDES, Len (Technical University of Munich)

Presenter: BRANDES, Len (Technical University of Munich)

Contribution ID: 40

Type: **not specified**

Measurement of the neutral pion transition form factor at low Q^2 at Jefferson Lab

The neutral pion transition form factor (TFF) plays an important role in tests of low energy QCD, and in the determination of the hadronic-light-by-light (HLbL) scattering contribution to the muon anomalous magnetic moment, $(g-2)$. Several measurements of this form factor exist in the large space-like Q^2 region, but the low Q^2 space-like region remains largely unexplored. This talk will present the details and impact of the proposed precision measurement of the TFF to be carried out in Hall B of Jefferson Lab. This experiment will use a 10.5 GeV electron beam, silicon target, and a low background beamline in Hall B to measure the Primakoff cross-section for neutral pion electro-production. These data will be used to extract the pion TFF in the Q^2 range of 0.001-0.1 GeV^2 , and constrain calculations for the HLbL contribution to a muon $(g-2)$ at low Q^2 . The author acknowledges support from D.O.E. grant DE-FG02-88ER40415

Primary author: LARIN, Ilya (Jefferson Lab)

Presenter: LARIN, Ilya (Jefferson Lab)

Contribution ID: 41

Type: **Parallel-Few-Body Physics**

Deuteron VVCS and nuclear structure effects in muonic deuterium at N3LO in pionless EFT

We calculate the forward unpolarised doubly-virtual Compton scattering (VVCS) off the deuteron in the framework of pionless effective field theory, up to next-to-next-to-next-to-leading order (N3LO) for the longitudinal and next-to-leading order (NLO) for the transverse amplitude. The charge elastic form factor of the deuteron, obtained from the residue of the longitudinal VVCS amplitude, is used to extract the value of the single unknown two-nucleon one-photon contact coupling that enters the longitudinal amplitude at N3LO.

Using the obtained deuteron VVCS amplitude as a high-precision model-independent input, we examine the two-photon-exchange (TPE) corrections to the Lamb shift of muonic deuterium, and find substantial differences with the recent dispersive evaluations. Namely, the elastic contribution appears to be larger by several standard deviations, thus ameliorating the current discrepancy between theory and experiment on the size of TPE effects. We investigate the correlation between the values of the deuteron charge and Friar radii and argue that it can be used to judge on the quality of a parametrisation of the deuteron charge elastic form factor. We also study other related effects, such as the TPE contribution to the proton-deuteron isotope shift and the spin-independent deuteron generalised polarizabilities.

Primary author: LENSKY, Vadim (JGU, Mainz)

Co-authors: Dr HILLER BLIN, Astrid (JLab); Dr HAGELSTEIN, Franziska (PSI, Villigen); Dr PAS-CALUTSA, Vladimir (JGU, Mainz)

Presenter: LENSKY, Vadim (JGU, Mainz)

Contribution ID: 42

Type: **Parallel-Few-Body Physics**

A1(1260) from lattice QCD

We report the first determination of universal parameters of the axial $a_1(1260)$ resonance from lattice QCD [arXiv:2107.03973 [hep-lat] in press at PRL]. Three-body quantization condition (FVU) is generalized and utilized to extract infinite-volume information from lattice eigenenergies. Subsequently, pole position and residua of the three-body resonance are extracted on the second Riemann sheet. Results are compared to the phenomenological ones.

Primary author: Dr MAI, Maxim (GWU)

Co-authors: ALEXANDRU, Andrei (GWU); CULVER, Chris (U. of Liverpool); SADASIVAN, Daniel (Ave Maria University); LEE, Frank (GWU); DORING, Michael (GWU); BRETT, Ruairi (GWU)

Presenter: Dr MAI, Maxim (GWU)

Contribution ID: 43

Type: **Parallel-Goldstone Boson**

pi0-eta-eta' mixing from V->Pgamma and P->Vgamma decays

An enhanced phenomenological model that includes isospin-symmetry breaking is presented in this letter.

The model is then used in a number of statistical fits to the most recent experimental data for the radiative transitions

$VP\gamma$ ($V = \rho, K^*, \omega, \phi$ and $P = \pi, K, \eta, \eta'$)

and estimations for the mixing angles amongst the three pseudoscalar states with vanishing third-component of isospin are obtained.

The quality of the performed fits is good, e.g. $\chi^2_{\min}/\text{d.o.f} = 1.9$.

The current experimental uncertainties allow for isospin-symmetry violations with a confidence level of approximately 2.5σ .

Primary author: ESCRIBANO, Rafel (Universitat Autònoma de Barcelona)

Co-author: ROYO, Emilio (Universitat Autònoma de Barcelona (UAB) and Institut de Física d'Altes Energies (IFAE))

Presenter: ESCRIBANO, Rafel (Universitat Autònoma de Barcelona)

Contribution ID: 44

Type: **Parallel-Goldstone Boson**

A dispersive analysis of low energy pion photo- and electroproduction

A dispersive representation based on unitarity and analyticity is used to study the low energy $\gamma N \rightarrow \pi N$ and $\gamma^* N \rightarrow \pi N$ partial wave amplitudes.

Final state interactions of the πN system are critical to this analysis.

The left-hand cut contribution is estimated by invoking $\mathcal{O}(p^2)$ baryon chiral perturbation theory results,

while the right-hand cut contribution responsible for final state interaction effects is taken into account via an Omnes formalism with elastic phase shifts as input.

It is found that a good numerical fit can be achieved with only one subtraction parameter, and the experimental data of the multipole amplitudes E_0^+, S_0^+ in the energy region below the $\Delta(1232)$ are well described when the photon virtuality $Q^2 \leq 0.1 \text{ GeV}^2$.

Furthermore, we extend the partial wave amplitudes to the second Riemann sheet to extract the couplings of the subthreshold resonance $N^*(890)$.

The values of residue of the multipole amplitudes E_0^+, S_0^+ are almost the same as that of the $N^*(1535)$ resonance, indicating that $N^*(890)$ strongly couples to the πN system.

Primary author: CAO, Xiong-Hui (Peking University)

Co-authors: Prof. ZHENG, Han-Qing (Sichuan University); MA, Yao (Hangzhou Institute for Advanced Study)

Presenter: CAO, Xiong-Hui (Peking University)

Contribution ID: 45

Type: **Parallel-Hadron Structure**

The Pole Counting Rule and X, Y, Z States

The pole counting rule, is a powerful and model-independent method to distinguish a confining state from a hadronic molecule. It has been applied to the explorations of $X(6900)$, $X_1(2900)$ as well as $Z_c(3900)$, $X(3872)$, $X(4660)$, etc. For $X(6900)$, both a confining state and a molecular state are possible, because lacking of enough data. For $X_1(2900)$, the analysis shows that it should be a $\bar{D}_1 K$ molecule, with $J^P = 1^-$ and an iso-singlet interpretation is much more favorable. Finally, it is noted that almost all X, Y, Z particles with exotic quantum numbers can be interpreted as hadronic molecules. The $X(3872)$ is, however, more like a charmonium, since it has a $\bar{c}c$ quantum number.

Primary author: Mr CHEN, Hao (Peking University)

Co-authors: Prof. ZHENG, Han-Qing (Sichuan University); Dr QI, Hong-Rong (Tsinghua University)

Presenter: Mr CHEN, Hao (Peking University)

Contribution ID: 46

Type: **Parallel-Goldstone Boson**

The N/D study on the singularity structure of πN scattering amplitudes

The N/D method is used to study the S_{11} channel low energy πN scattering amplitude. The inputs of left cuts are obtained from various phenomenological models at tree level. With the aid of the production representation, the total phase shifts can be decomposed into different contributions, and it further reveals that the existence of subthreshold resonance $N^*(890)$ doesn't depend on the details of the dynamical inputs. Additionally, it is found that there exist virtual states in partial waves, which are induced by the u channel nucleon exchanges. These virtual states accumulate at the end point of the u channel segment cut. The end point is hence the essential singularity of the full amplitude on the second sheet of complex s plane.

Primary author: Mr LI, Qu-Zhi (Peking University)

Co-authors: Prof. ZHENG, Han-Qing (SiChuan University); Dr MA, Yao (Hangzhou Institute for Advanced Study)

Presenter: Mr LI, Qu-Zhi (Peking University)

Contribution ID: 47

Type: **Parallel-Few-Body Physics**

Fluctuations and phases in baryonic matter

The phase structure of baryonic matter is investigated with focus on the role of fluctuations beyond the mean-field approximation. The prototype test case studied is the chiral nucleon-meson model, with added comments on the chiral quark-meson model. Applications to nuclear matter include the liquid-gas phase transition. Extensions to high baryon densities are performed for both nuclear and neutron matter. The role of vacuum fluctuations is systematically explored. It is pointed out that such fluctuations tend to stabilize the hadronic phase characterized by spontaneously broken chiral symmetry, shifting the chiral restoration transition to very high densities. This stabilization effect is shown to be further enhanced by additional dynamical fluctuations treated with functional renormalisation group methods.

Primary author: BRANDES, Len (Technical University of Munich)

Presenter: BRANDES, Len (Technical University of Munich)

Contribution ID: 48

Type: **not specified**

The JLab Eta Factory (JEF) experiment

The new experiment, JLab Eta Factory (JEF), in the experimental Hall D at Jefferson Lab will extend the physics potential of the GlueX detector beyond the main spectroscopy program and perform precision measurements of various $\eta^{(\prime)}$ decays with emphasis on rare neutral modes. The physics program of the experiment spans from precision tests of low-energy QCD to search of gauge bosons in the mass range below 1 GeV coupling the SM sector to the dark sector. Photoproduction of highly boosted $\eta^{(\prime)}$ mesons using a tagged photon beam, good detection of recoil proton and multi-photon final states will allow to suppress background and collect high-statistics data sample of η mesons. All these provide many advantages over other $\eta^{(\prime)}$ experiments. The JEF experiment requires to upgrade the inner part of the forward lead glass calorimeter of the GlueX detector with high-granularity, high-resolution lead tungstate PbWO₄ scintillating crystals. The calorimeter insert is currently under construction at Jefferson Lab. The detector will be ready to take data in 2024. An overview of the JEF project will be presented.

Primary author: SOMOV, Alexander (Jefferson Lab)

Presenter: SOMOV, Alexander (Jefferson Lab)

Contribution ID: 49

Type: **Parallel-Few-Body Physics**

Nucleon-level Effective Theory of $\mu \rightarrow e$ Conversion

The Mu2E and COMET $\mu \rightarrow e$ collaborations plan to advance branching ratio sensitivities by four orders of magnitude, further constraining new sources of charged lepton flavor violation (CLFV). We formulate a non-relativistic nucleon-level effective theory for this process, in order to clarify what can and cannot be learned about CLFV operator coefficients from elastic $\mu \rightarrow e$ conversion. Utilizing state-of-the-art shell model wave functions, we derive bounds on operator coefficients from existing $\mu \rightarrow e$ conversion and $\mu \rightarrow e \gamma$ results, and estimate the improvement in these bounds that will be possible if Mu2E, COMET, and MEG II reach their design goals. In the conversion process, we employ a treatment of the lepton Coulomb physics that is very accurate, yet yields transparent results and preserves connections to standard-model processes like β decay and μ capture. The formulation provides a bridge between the nuclear physics needed in form factor evaluations and the particle physics needed to relate low-energy constraints from $\mu \rightarrow e$ conversion to UV sources of CLFV.

Primary author: RULE, Evan (U)**Presenter:** RULE, Evan (U)

Contribution ID: 50

Type: **Parallel-Goldstone Boson**

Dispersive determination of low energy πK interactions

The determination of low energy πK scattering has been subject to debate during many years. The precise and robust extraction of these processes is relevant for both experimentalist and phenomenologists. In particular, the low energy expansion of the partial waves offers rich information for chiral perturbation theory practitioners and can be compared to modern lattice QCD calculations. Furthermore, the long debated $\kappa/K_0^*(700)$ resonance lives in the vicinity of the $I = 1/2$ πK scalar threshold. In this talk we present a dispersive, model-independent determination of these low energy interactions, including the low energy expansion parameters, and compare them to previous determinations. We first use a large set of dispersion relations as constraints on combined fits to $\pi K \rightarrow \pi K$ and $\pi\pi \rightarrow K\bar{K}$ data. Then we implement different sum rules to extract these low energy parameters, producing a final set of data driven dispersive results.

Primary author: Dr RODAS, ARKAITZ (College of William and Mary)

Co-author: Prof. PELAEZ, Jose R. (Universidad Complutense)

Presenter: Dr RODAS, ARKAITZ (College of William and Mary)

Contribution ID: 51

Type: **Parallel-Hadron Structure**

Overview of TMD studies

Transverse momentum dependent (TMD) parton distributions provide three-dimensional imaging of hadrons in the momentum space and the correlation with spins. The precise measurement of TMDs is one of the main goals of many JLab12 experiments and future EIC and EicC programs. It has become an important and very active area of hadron physics from both experimental and theoretical aspects. In this talk, I will give an overview of recent progress, present status and future outlook in TMD studies.

Primary author: LIU, Tianbo (Shandong University)

Presenter: LIU, Tianbo (Shandong University)

Contribution ID: 52

Type: **Parallel-Few-Body Physics**

Constraints on the Λ -neutron interaction from charge symmetry breaking of $A = 4$ hypernuclei

Charge symmetry breaking (CSB) of the Λ -nucleon interaction has been well established by the experimentally known difference of the Λ separation energies of the mirror hypernuclei ${}^4_{\Lambda}\text{He}$ and ${}^4_{\Lambda}\text{H}$ [1]. At the same time, accurate predictions for these quantities are possible based on solutions of Faddeev-Yakubovsky equations[2].

In this contribution, we employ chiral hyperon-nucleon interactions including the leading CSB contributions to constrain the Λ -neutron interaction. To this aim, we determine the strength of the two arising CSB contact terms by a fit to the differences of the separation energies of these hypernuclei in the 0^+ and 1^+ states, respectively, and then predict Λn scattering lengths[3]. Based on two version of the hyperon-nucleon interaction at next-to-leading order and using different momentum cutoffs, we also estimate uncertainties of these predictions. The impact of the possible changes of the experimental input is discussed in view of recently improved experimental results for the separation energies of ${}^4_{\Lambda}\text{He}$ and ${}^4_{\Lambda}\text{H}$.

References

- [1] A. Nogga, H. Kamada and W. Glöckle, Phys. Rev. Lett. 88 (2002), 172501 [arXiv:nucl-th/0112060 [nucl-th]].

- [2] A. Gal, E.V. Hungerford and D.J. Millener, Rev. Mod. Phys. 88 (2016), 035004 [arXiv:1605.00557 [nucl-th]].

- [3] J. Haidenbauer, U.-G. Meißner and A. Nogga, Few Body Syst. 62 (2021), 105 [arXiv:2107.01134 [nucl-th]].

Primary author: Dr NOGGA, Andreas (Forschungszentrum Juelich GmbH)

Co-authors: Dr HAIDENBAUER, Johann (Forschungszentrum Juelich GmbH); Prof. MEISSNER, Ulf-G. (Universitaet Bonn and Forschungszentrum Juelich)

Presenter: Dr NOGGA, Andreas (Forschungszentrum Juelich GmbH)

Contribution ID: 53

Type: **Parallel-Goldstone Boson**

Measurement of the neutral pion transition form factor at low Q^2 at Jefferson Lab

The neutral pion transition form factor (TFF) plays an important role in tests of low energy QCD, and in the determination of the hadronic-light-by-light (HLbL) scattering contribution to the muon anomalous magnetic moment, $(g-2)$. Several measurements of this form factor exist in the large space-like Q^2 region, but the low Q^2 space-like region remains largely unexplored. This talk will present the details and impact of the proposed precision measurement of the TFF to be carried out in Hall B of Jefferson Lab. This experiment will use a 10.5 GeV electron beam, silicon target, and a low background beamline in Hall B to measure the Primakoff cross-section for neutral pion electro-production. These data will be used to extract the pion TFF in the Q^2 range of 0.001-0.1 GeV^2 , and constrain calculations for the HLbL contribution to a muon $(g-2)$ at low Q^2 . The author acknowledges support from D.O.E. grant DE-FG02-88ER40415

Primary author: Dr LARIN, Ilya (UMASS Amherst)

Presenter: Dr LARIN, Ilya (UMASS Amherst)

Contribution ID: 54

Type: **Parallel-Hadron Structure**

Resonance dynamics in U(3) chiral theory and its relevance in the determination of light-quark mass

I will first review our study of the resonance dynamics in the U(3) chiral theory. Then I will focus on the determination of the light-quark mass by taking the spectral functions from the U(3) chiral theory within the framework of QCD sum rule. It turns out that once the scalar dynamics in the spectral function is properly included, the resulting values of the light-quark mass in the isoscalar scalar QCD sum rules become compatible with those from other approaches.

Primary author: Prof. GUO, Zhi-Hui (Southeast University)

Presenter: Prof. GUO, Zhi-Hui (Southeast University)

Contribution ID: 55

Type: **Parallel-Hadron Structure**

Hadron physics results at KLOE-2 experiment

KLOE and KLOE-2 data (almost 8 fb^{-1}) constitute the largest sample ever collected at an electron-positron collider operating at the ϕ peak resonance.

In total it corresponds to the production of about 24 billion of ϕ mesons whose decays include about 8 billion pairs of neutral K mesons and about 300 million η mesons.

A wide hadron physics program, investigating rare meson decays, $\gamma - \gamma$ interaction, dark forces and hadronic cross section, is thus carried out by the KLOE-2 Collaboration.

The $\eta \rightarrow \pi^0 \gamma \gamma$ decay is a test bench for various models and effective theories like VMD (Vector Meson Dominance) or ChPT (Chiral

Perturbation Theory) which predict BR far from experimental value. KLOE-2, with its highly pure η sample produced

in $\phi \rightarrow \eta \gamma$ process, has a new preliminary measurement of this branching ratio.

Following the many contributions KLOE and KLOE-2 has done to Dark Matter searches, an alternative model, where the Dark Force mediator is an hypothetical leptophobic B boson, is investigated in the $\phi \rightarrow \eta B \rightarrow \eta \pi^0 \gamma, \eta \rightarrow \gamma \gamma$ channel.

A KLOE-2 distinctive feature is also the the possibility to investigate π^0 production from $\gamma \gamma$ scattering by

tagging final-state leptons from $e^+ e^- \rightarrow \gamma^* \gamma^* e^+ e^- \rightarrow \pi^0 e^+ e^-$ in coincidence with the π^0 in the barrel calorimeter. KLOE-2 aims to use this process to precisely measure the π^0 decay width into $\gamma \gamma$ to test low-energy QCD dynamics. Progresses made on the $\gamma^* \gamma^* \rightarrow \pi^0$ event counting will be reported.

A search for the P and CP violating decay $\eta \rightarrow \pi^+ \pi^-$ has been recently published by KLOE-2 using 1.6 fb^{-1} of KLOE data.

No signal is observed in the $\pi^+ \pi^-$ invariant mass spectrum, and the

upper limit on the branching fraction at 90% confidence level is $B(\eta \rightarrow \pi^+ \pi^-) < 4.9 \times 10^{-6}$, which is

three times lower than previous KLOE result. A combination of the two KLOE limits will be presented.

Moreover, the search for the double suppressed $\phi \rightarrow \eta \pi^+ \pi^-$ and the conversion

$\phi \rightarrow \eta \mu^+ \mu^-$ decays are performed at KLOE-2 with both $\eta \rightarrow \gamma \gamma$ and $\eta \rightarrow 3\pi^0$ final states. Clear signals are seen for the first time.

Finally, preliminary and promising results on the ω cross section measurement in the $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0 \gamma_{\text{ISR}}$ channel using the Initial State Radiation (ISR) method will be also presented.

Primary author: MANDAGLIO, Giuseppe (University of Messina)

Presenter: MANDAGLIO, Giuseppe (University of Messina)

Contribution ID: 56

Type: **Parallel-Few-Body Physics**

$A = 4 - 7$ Ξ hypernuclei based on interactions from chiral effective field theory

Experimental information on hypernuclei with double strangeness, in particular Ξ systems, is very limited because of low intensity of kaon beams and the extremely short life times of hyperons. In this contribution, I report on an investigation of the possible existence of bound Ξ states in systems with $A = 4 - 7$ baryons using the Jacobi NCSM [1] approach in combination with chiral NN [2] and Ξ N [3] interactions. Three shallow bound states for the NNN Ξ system (with $(J^\pi, T) = (1^+, 0)$, $(0^+, 1)$ and $(1^+, 1)$) with quite similar binding energies are found. The ${}^5_{\Xi}\text{H}(\frac{1}{2}^+, \frac{1}{2})$ and ${}^7_{\Xi}\text{H}(\frac{1}{2}^+, \frac{3}{2})$ hypernuclei are also clearly bound with respect to the thresholds ${}^4\text{He} + \Xi$ and ${}^6\text{He} + \Xi$, respectively. A perturbative estimate suggests that the decay widths of these states could be sufficiently small enough for them to be observed in experiment. Such an observation would provide important experimental constraints on Ξ N interactions.

References

- [1] S. Liebig, U.-G. Meißner, and A. Nogga, Eur. Phys. J. A, 52(4):103, 2016
- [2] E. Epelbaum, H. Hammer, and U.-G. Meißner, Rev. Mod. Phys., 81:1773-1825, 2009
- [3] J. Haidenbauer, U.-G. Meißner, Eur. Phys. J., A55(2):23, 2019

Primary author: Dr LE, Hoai (Forschungszentrum Juelich)

Co-authors: Dr NOGGA, Andreas (Forschungszentrum Juelich); Dr HAIDENBAUER, Johann (Forschungszentrum Juelich); Prof. MEISSNER, Ulf-G (HISKP and BCTP, Universität Bonn, Germany)

Presenter: Dr LE, Hoai (Forschungszentrum Juelich)

Contribution ID: 57

Type: **Parallel-Hadron Structure**

Results on spin sum rules and polarizabilities at low Q^2 .

We will report on recently published experimental results on spin sum rules, and particularly on the generalized spin polarizabilities $\gamma_0(Q^2)$ (for both the proton and neutron) and $\delta_{LT}(Q^2)$ (for the neutron).

The data were taken at Jefferson Lab in Hall A (neutron) and B (proton and deuteron) by experiments E97-110 and EG4, respectively. They covered the very low Q^2 domain, down to $Q^2 \sim 0.02 \text{ GeV}^2$, where Chiral Effective Field Theory (χ EFT) predictions should be valid. While some observables agree with the state-of-the-art χ EFT theoretical predictions, others are in tensions, including $\delta_{LT}^n(Q^2)$ for which χ EFT prediction was expected to be robust. This suggests that χ EFT does not yet consistently describe nucleon spin observables, even in the very low Q^2 domain covered by the experiments.

Primary author: DEUR, Alexandre (J)

Presenter: DEUR, Alexandre (J)

Contribution ID: 58

Type: **Plenary**

The REDTOP experiment: a low energy meson factory to explore dark matter and physics beyond the Standard model

The η and η' mesons are almost unique in the particle universe since they are Goldstone boson and the dynamics of their decay are strongly constrained. The integrated eta meson samples collected in earlier experiments have been about $\sim 10^9$ events, dominated by the WASA at Cosy experiment, limiting considerably the search for such rare decays. A new experiment, REDTOP, is being proposed, with the intent

of collecting more than 10^{13} eta/yr (10^{11} eta'/yr) for studying of rare η decays.

Such statistics are sufficient for investigating several symmetry violations, and for searches of new particles beyond the Standard Model.

With tagged-eta experiment the fully constrained kinematic of the process allows for searches of light dark matter with a “Missing 4-momentum technique” which, at present, cannot be exploited by any other existing or proposed experiment.

The physics program and the detector for REDTOP will be discussed during the presentation.

Primary author: Dr GATTO, Corrado (INFN & NIU)

Co-author: ON BEHALF OF: REDTOP, Collaboration (NIU)

Presenter: Dr GATTO, Corrado (INFN & NIU)

Contribution ID: 59

Type: **Plenary**

The Muon Proton Scattering Experiment (MUSE) at PSI

In 2010, high-precision studies of muonic hydrogen found notably smaller values for the charge radius than earlier results that have been extracted from elastic electron-scattering data and through the spectroscopy of atomic hydrogen. The MUon Scattering Experiment (MUSE) at the Paul Scherrer Institute (PSI) has been developed to address this so-called proton-radius puzzle. The experiment will measure elastic electron-proton and muon-proton scattering data with positively and negatively charged beams in a four-momentum-transfer range from 0.002 to 0.08 GeV². Each of the four sets of data will allow the extraction of the proton charge radius. In combination, the data test possible differences between the electron and muon interactions and two-photon exchange effects. The status of the experiment will be discussed.

- This material is based upon work supported by the National Science Foundation under NSF grant PHY-1913653. The MUSE experiment is supported by the Department of Energy, NSF, PSI, and the US-Israel Binational Science Foundation.

Primary author: STRAUCH, Steffen (University of South Carolina)

Presenter: STRAUCH, Steffen (University of South Carolina)

Contribution ID: 60

Type: **Parallel-Hadron Structure**

Overview of GPD Studies

Generalized Parton distributions (GPDs) are a new theoretical tool that was developed in the late 90s. GPDs not only link the well-known form factors and parton distribution functions but also provide much richer correlation information between the transverse location and the longitudinal momentum of partons. More importantly, they can access the contribution of the orbital angular momentum of quarks (and gluons) to the nucleon spin via Ji's spin sum-rule. Several exclusive physics processes can probe GPDs, including Deeply Virtual Compton Scattering (DVCS), Deep Virtual Meson Production (DVMP), Time-Like Compton Scattering (TCS), as well as the Double DVCS. A brief introduction of the GPDs will be given followed by a review of past, ongoing and future experimental programs at Jefferson Lab (SoLID, CLAS12, etc), Electron-Ion Collider in China and the U.S., and other places.

Primary author: YE, Zhihong (Tsinghua University)

Presenter: YE, Zhihong (Tsinghua University)

Contribution ID: 61

Type: **Parallel-Goldstone Boson**

HVP contribution to $g-2$

Hadronic vacuum polarization currently yields the dominant uncertainty in the Standard-Model prediction for the anomalous magnetic moment of the muon. While the phenomenological approach is only as accurate as the hadronic cross sections used as input, there are several aspects related to chiral dynamics that can be used as cross checks, including $\pi\pi$ dynamics and the chiral anomaly. In the talk I will give an overview over such aspects, including recent work to extrapolate the isovector HVP contribution to unphysical quark masses.

Primary author: HOFERICHTER, Martin (University of Bern)

Presenter: HOFERICHTER, Martin (University of Bern)

Contribution ID: 62

Type: **Parallel-Hadron Structure**

VCS and generalized polarizabilities from Mainz and JLab: Overview and new results

The Generalized Polarizabilities (GPs) are fundamental properties of the nucleon. They characterize the nucleon's response to an applied electromagnetic field, giving access to the polarization densities inside the nucleon. As such the GPs represent a central path towards a complete understanding of the nucleon dynamics. Previous measurements of the proton electric GP at intermediate four-momentum transfer squared have challenged the predictions of theoretical calculations, raising questions in regard to the underlying reasons responsible for a local enhancement of the electric GP. The measurement of the magnetic GP on the other hand promises to quantify the interplay of the paramagnetism and diamagnetism contributions inside the proton. An overview on this topic, new results from JLab and future prospects will be discussed in this talk.

Primary author: Prof. SPARVERIS, Nikos (Temple University)

Presenter: Prof. SPARVERIS, Nikos (Temple University)

Contribution ID: 63

Type: **not specified**

A theory of dark pions

We present a complete model of a dark QCD sector with light dark pions, broadly motivated by hidden naturalness arguments. The dark quarks couple to the Standard Model via irrelevant Z - and Higgs-portal operators, which encode the low-energy effects of TeV-scale fermions interacting through Yukawa couplings with the Higgs field. The dark pions, depending on their CP properties, behave as either composite axion-like particles (ALPs) mixing with the Z or scalars mixing with the Higgs. The dark pion lifetimes fall naturally in the most interesting region for present and proposed searches for long-lived particles, at the LHC and beyond. This is demonstrated by studying in detail three benchmark scenarios for the symmetries and structure of the theory. Within a coherent framework, we analyze and compare the GeV-scale signatures of flavor-changing meson decays to dark pions, the weak-scale decays of Z and Higgs bosons to hidden hadrons, and the TeV-scale signals of the ultraviolet theory. New constraints are derived from B decays at CMS and from Z -initiated dark showers at LHCb, focusing on the displaced dimuon signature. We also emphasize the strong potential sensitivity of ATLAS and CMS to dark shower signals with large multiplicities and long lifetimes of the dark pions. As a key part of our phenomenological study, we perform a new data-driven calculation of the decays of a light ALP to exclusive hadronic Standard Model final states. The results are provided in a general form, applicable to any model with arbitrary flavor-diagonal couplings of the ALP to fermions.

Primary authors: LI, Lingfeng (Brown University); LI, Lingfeng (HKUST)

Presenter: LI, Lingfeng (Brown University)

Contribution ID: 64

Type: **Parallel-Hadron Structure**

A Low Q^2 Measurement of the Proton Spin Structure Function g_2^p

We have extracted the spin structure functions g_1 and g_2 of the proton and their moments at $Q^2 < 0.13 \text{ GeV}^2$. This data was obtained with the Jefferson Lab polarized electron beam, a polarized solid NH_3 target, and the Jefferson Lab Hall A High Resolution Spectrometers. The structure functions were measured by calculating the asymmetry for both transverse and longitudinal kinematics and using it to form polarized cross section differences $\Delta\sigma_{\parallel}(\nu, Q^2)$ and $\Delta\sigma_{\perp}(\nu, Q^2)$. These structure functions were used to form several moments, many of which can be directly compared to predictions of Chiral Perturbation Theory (χ PT): \overline{F}_2 , δ_{LT} , \overline{d}_2 , γ_0 , and I_{LT} . These results represent the first experimental determination of δ_{LT} for the proton at low Q^2 . Current χ PT calculations differ in the measured region, and our data shows a strong preference for one of these calculations. Data published in 2004 showed a strong disagreement between neutron δ_{LT} data and chiral theory at low Q^2 , this “ δ_{LT} Puzzle” has since been well studied, but recently published neutron data shows a new discrepancy, making it very important to study the behavior of this moment for the proton. The proton results shown in this talk agree well with the χ PT calculation using a delta power counting scheme and less well with the calculation which uses an epsilon power counting scheme, with the difference in power counting being one known difference between the two.

Summary

We present proton spin structure function results g_1^p and g_2^p as well as their moments, comparing to leading calculations of chiral perturbation theory. These results show good agreement with one of these calculations for several 0th and 2nd order SSF moments.

Primary author: RUTH, David (U)

Co-authors: Dr CHEN, Jian-ping (Jefferson Lab); SLIFER, Karl (U)

Presenter: RUTH, David (U)

Contribution ID: 65

Type: **Parallel-Goldstone Boson**

Data-driven dispersive analysis of the $\pi\pi$ and πK scattering

We present a data-driven analysis of the resonant S-wave $\pi\pi \rightarrow \pi\pi$ and $\pi K \rightarrow \pi K$ reactions using the partial-wave dispersion relation. The contributions from the left-hand cuts are accounted for using the Taylor expansion in a suitably constructed conformal variable. The fits are performed to experimental and lattice data as well as Roy analyses. For the $\pi\pi$ scattering we present both a single- and coupled-channel analysis by including additionally the $K\bar{K}$ channel. For the latter the central result is the Omnès matrix, which is consistent with the most recent Roy and Roy-Steiner results on $\pi\pi \rightarrow \pi\pi$ and $\pi\pi \rightarrow K\bar{K}$, respectively. By the analytic continuation to the complex plane, we found poles associated with the lightest scalar resonances $\sigma/f_0(500)$, $f_0(980)$, and $\kappa/K_0^*(700)$ for the physical pion mass value and in the case of $\sigma/f_0(500)$, $\kappa/K_0^*(700)$ also for unphysical pion mass values.

Primary author: Ms DEINEKA, Oleksandra (Johannes Gutenberg University of Mainz)

Co-authors: Dr DANILKIN, Igor (Johannes Gutenberg University of Mainz); Prof. VANDER-HAEGHEN, Marc (Johannes Gutenberg University of Mainz)

Presenter: Ms DEINEKA, Oleksandra (Johannes Gutenberg University of Mainz)

Contribution ID: 66

Type: **not specified**

Probing few-body nuclear dynamics via ^3H and ^3He $(e,e'p)$ cross-section measurements at Jefferson Lab

While great progress has been made toward understanding the strong nucleon-nucleon (NN) interaction, there is still substantial uncertainty about the short-distance behavior of the nuclear force. Specifically, Quantum Monte Carlo (QMC) calculations using different state-of-the-art NN potential models can still make vastly different predictions at distance scales below 1 fm (small- r). Nucleon pairs at this separation distance are called Short-Range Correlations (SRCs). As a result of their repulsive small- r interaction, nucleons in an SRC pair fly apart from one another with high momenta (high- k). The study of SRCs is a powerful tool to constrain the NN interaction at small- r and high- k , and has significant implications not only nuclear structure, but to other fields as well, such as the astrophysics of neutron stars and the behavior of cold atomic gasses. In this talk, I will present results from the first measurement of the ^3He and $^3\text{H}(e,e'p)$ reactions in Hall A of the Thomas Jefferson National Accelerator Facility in kinematics in which the measured cross sections are expected to be sensitive to the underlying nucleon momentum distributions in the range 40 to 500 MeV/c. The resulting absolute cross sections were compared to precise cross-section calculations. These results defy the expectation of a $^3\text{He}/^3\text{H}$ ratio approaching unity for protons at high- k . This forces us to re-evaluate our understanding the reaction mechanism, of the $A=3$ wave function, and even of the NN interaction. The extracted absolute cross sections can be used to understand the shortcomings of current ^3He and ^3H cross section calculations and as a benchmark for future ones. Including the effects of outgoing nucleon rescattering improves agreement with the data at high missing momentum and suggests contributions from charge-exchange (SCX) rescattering. The isoscalar sum of ^3He plus ^3H is described by calculations to within the accuracy of the data over the entire p_{miss} range. This validates current models of the ground state of the three-nucleon system up to very high initial nucleon momenta.

Primary author: CRUZ-TORRES, Reynier (i)

Presenter: CRUZ-TORRES, Reynier (i)

Contribution ID: 67

Type: **Parallel-Few-Body Physics**

First covariant high-precision chiral nucleon-nucleon interaction up to next-to-next-to-leading order

We construct the next-to-leading and next-to-next-to-leading chiral nucleon-nucleon interaction in covariant baryon chiral perturbation theory. We show that a rather good description of the np phase shifts up to $E_{\text{lab}} = 200$ MeV can be achieved with a $\chi^2/\text{d.o.f.}$ less than 1. The resulting potential can be employed in ab initio studies of nuclear structure and reactions in a covariant framework.

Primary author: Dr GENG, Lisheng (Beihang University)

Presenter: Dr GENG, Lisheng (Beihang University)

Contribution ID: 68

Type: **not specified**

Recent global fits of the polarized and » unpolarized PDFs from JAM collaboration

The JAM collaboration has carried out several recent works in global QCD analysis, and I will summarize some of them in this talk.

The JAM-20-SIDIS, which was published in 2020, features simultaneous fit of the PDFs and FFs including unidentified charged hadrons and confirms the suppression of the strange quark PDF observed in JAM19.

More recently in October 2021, an analysis using the latest DIS data on helium and tritium from the MARATHON collaboration was published, which reveals an isovector component to the EMC effect at high x using a simultaneous study of nucleon PDFs and nuclear effects.

Later in this year, a simultaneous extraction of both helicity basis PDFs will be available on arXiv, in which we also critically studies the impact of theory assumptions on the spin-dependent PDFs. Lastly, I will talk about the sea asymmetries ($\bar{u} - \bar{d}$ or $\Delta\bar{u} - \Delta\bar{d}$) extracted with the latest data from the SeaQuest and STAR collaborations.

Primary author: 周, 一雨 (å)

Presenter: 周, 一雨 (å)

Contribution ID: 69

Type: **Parallel-Few-Body Physics**

Three pion and kaon scattering from lattice QCD

Hadron scattering information can be indirectly accessed through lattice QCD calculations of the finite-volume spectrum of multi-hadron states. The quantization condition connects the finite-volume energies to a description of scattering in the infinite-volume and has been very successful in the two-hadron sector. We will present an extension of the quantization condition to three-particle systems(FVU) and discuss its application to lattice results for three pion and kaon systems.

Primary author: CULVER, Christopher (University of Liverpool)

Presenter: CULVER, Christopher (University of Liverpool)

Contribution ID: 70

Type: **Parallel-Hadron Structure**

Possible bound states of $J/\psi J/\psi$

In a recent measurement LHCb reported pronounced structures in the $J/\psi J/\psi$ spectrum. One of the various possible explanations of those is that they emerge from non-perturbative interactions of vector charmonia. It is thus important to understand whether it is possible to form a bound state of two charmonia interacting through the exchange of gluons, which hadronise into two pions at the longest distance. In this paper, we demonstrate that, given our current understanding of hadron-hadron interactions, the exchange of correlated light mesons (pions and kaons) is able to provide sizeable attraction to the di- J/ψ system, and it is possible for two J/ψ mesons to form a bound state.

Primary author: 董, 相坤 (理论所)**Presenter:** 董, 相坤 (理论所)

Contribution ID: 71

Type: **Parallel-Hadron Structure**

Detecting the pure triangle singularity effect through the $\psi(2S)$ decay

In this talk, the triangle singularity mechanism is investigated in the two reactions of ψ decay, $\psi(2S) \rightarrow p\bar{p}\eta$ and $\psi(2S) \rightarrow \pi^+\pi^-K^+K^-$. They would generate a very narrow peak in the invariant mass spectrum of final states. In these processes, all the involved vertices are constrained by the experimental data. Thus, we can make a precise prediction here. We expect these effects can be observed by the Beijing Spectrometer and Super Tau-Charm Facility in the future.

Primary author: 吴, 佳俊 (University of Chinese Academy of Sciences)

Presenter: 吴, 佳俊 (University of Chinese Academy of Sciences)

Contribution ID: 72

Type: **Parallel-Few-Body Physics**

Baryon masses and currents in SU(3) BChPT x 1/Nc

Baryon Chiral Perturbation Theory (BChPT) and the $1/N_c$ expansion provide systematic frameworks for the strong interactions at low energy. A combined framework of both expansions has been developed and applied for baryons with three light-quark-flavors. The small scale expansion of the combined approach is identified as the ξ -expansion, in which the power counting of the expansions is linked according to $O(p) = O(1/N_c) = O(\xi)$. Results obtained from applying the combined framework to baryon masses, sigma terms and vector & axial-vector currents in SU(3) will be discussed in this talk.

Primary authors: FERNANDO, Ishara (University of Virginia); Prof. GOITY, Jose (Hampton University & Jefferson Lab)

Presenter: FERNANDO, Ishara (University of Virginia)

Contribution ID: 73

Type: **Parallel-Few-Body Physics**

Towards the Study of Short Range Correlations in Radioactive Nuclei: The transparent nucleus in inverse kinematics

Understanding the structure of a strongly-interacting quantum mechanical system such as atomic nuclei is a formidable challenge in physics. Nucleon knockout reactions with high energy probes are widely used to reveal the inner structure of nuclei, however they cannot be applied to study unstable nuclei. We recently demonstrated the feasibility to access Short-Range Correlation (SRC) properties in nuclei with hadronic probes in inverse kinematics, opening the pathway for such studies in short-lived nuclei at upcoming accelerator facilities. The experiment was carried out at the JINR (Russia), a ^{12}C beam at 48 GeV/c impinged on a liquid hydrogen target and the reaction products were measured kinematically complete with the BM@N detector setup. We show that by selecting the fragment in the $^{12}\text{C}(p,2p)^{11}\text{B}$ reaction limitations posed by final-state interactions are overcome and single nucleon properties are probed in a single-step knockout reaction. The extracted ground-state distributions are in agreement with theoretical calculations. We probe SRCs in the same way by the break up of SRC pairs in $^{12}\text{C}(p,2pN)^{10}\text{B}/^{10}\text{Be}$ reactions. We not only identify SRCs in such kinematical conditions for the first time but also deduce factorization and other pair properties from direct measurements. I will discuss the recent results and the future experimental program towards the study of SRCs in radioactive nuclei.

Primary author: KAHLBOW, Julian (MIT)

Presenter: KAHLBOW, Julian (MIT)

Contribution ID: 74

Type: **not specified**

opening speech

Monday, 15 November 2021 19:00 (10 minutes)

Presenter: Prof. WANG, Yifang (IHEP)

Session Classification: Plenary session (Chair: Xiaoyan Shen)

Contribution ID: 75

Type: **not specified**

Chiral Dynamics: Theory and Experiment - A Tribute to Aron Bernstein

Monday, 15 November 2021 19:10 (40 minutes)

Presenter: Prof. MEISSNER, Ulf-G. (Universitaet Bonn and Forschungszentrum Juelich)

Session Classification: Plenary session (Chair: Xiaoyan Shen)

Contribution ID: 76

Type: **not specified**

Highlights of BESIII Physics

Monday, 15 November 2021 19:50 (40 minutes)

Presenter: Dr MA, Hailong (IHEP)

Session Classification: Plenary session (Chair: Xiaoyan Shen)

Contribution ID: 77

Type: **not specified**

Chiral effective theory in the Higgs sector

Monday, 15 November 2021 20:30 (40 minutes)

Presenter: JUAN JOSE SANZ-CILLERO (Madrid Univ.)

Session Classification: Plenary session (Chair: Xiaoyan Shen)

Contribution ID: 78

Type: **not specified**

Theoretical aspects of virtual states, bound states, and resonances in hadron physics from Friedrichs model

Monday, 15 November 2021 21:30 (40 minutes)

Presenter: 肖, 志广 (Sichuan University)

Session Classification: Plenary session (Chair : Ulf-G. Meissner)

Contribution ID: 79

Type: **not specified**

Experimental Status on the Proton Charge Radius

Monday, 15 November 2021 22:10 (40 minutes)

Presenter: Prof. GAO, Haiyan (Duke Univ.)

Session Classification: Plenary session (Chair : Ulf-G. Meissner)

Contribution ID: 80

Type: **not specified**

Statistically rigorous analyses of light nuclei with chiral interactions

Monday, 15 November 2021 22:50 (40 minutes)

Presenter: FURNSTAHL, Dick (Ohio State University)

Session Classification: Plenary session (Chair : Ulf-G. Meissner)

Contribution ID: **81**

Type: **not specified**

Review of theoretical calculations on π^0 radiative decay width

Tuesday, 16 November 2021 19:30 (20 minutes)

Presenter: KAMPF, Karol (Charles University)

Session Classification: Goldstone Boson Dynamics (Chair: Bastian Kubis)

Contribution ID: 82

Type: **not specified**

Status of π^0 , η and η' experimental program in KLOE-II

Presenter: TBD, TBD

Contribution ID: 83

Type: **not specified**

Light Meson decays at BESIII

Tuesday, 16 November 2021 18:50 (20 minutes)

Presenter: KANG, Xiaolin (CUG)

Session Classification: Goldstone Boson Dynamics (Chair: Bastian Kubis)

Contribution ID: 84

Type: **not specified**

Dispersive approach to strong three-body decays of eta and eta' mesons

Tuesday, 16 November 2021 19:50 (20 minutes)

Presenter: ISKEN, Tobias (GSI)

Session Classification: Goldstone Boson Dynamics (Chair: Bastian Kubis)

Contribution ID: 85

Type: **not specified**

π^0 - η - η' mixing from $V \rightarrow P\gamma$ and $P \rightarrow V\gamma$ decays

Tuesday, 16 November 2021 21:30 (20 minutes)

Presenter: ESCRIBANO, Rafel (Universitat Autònoma de Barcelona)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: 86

Type: **not specified**

A theoretical analysis of the semileptonic decays η/η' $\rightarrow \pi^0 l^+ l^-$ and η' $\rightarrow \eta l^+ l^-$

Tuesday, 16 November 2021 20:10 (20 minutes)

Presenter: ROYO, Emilio (Universitat Autònoma de Barcelona (UAB) and Institut de Física d'Altes Energies (IFAE))

Session Classification: Goldstone Boson Dynamics (Chair: Bastian Kubis)

Contribution ID: 87

Type: **not specified**

Patterns of C- and CP-violation in hadronic eta and eta' three-body decays

Tuesday, 16 November 2021 20:30 (20 minutes)

Presenter: AKDAG, Hakan (Bonn University)

Session Classification: Goldstone Boson Dynamics (Chair: Bastian Kubis)

Contribution ID: 88

Type: **not specified**

Review of light quark mass ratio determination via $\eta \rightarrow 3\pi$

Tuesday, 16 November 2021 21:10 (20 minutes)

Presenter: PASSEMAR, Emilie (Indiana University/JLab)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: 89

Type: **not specified**

Theoretical analysis of the doubly radiative decays η/η' \rightarrow $\pi^0 \gamma \gamma$ and $\eta' \rightarrow \eta \gamma \gamma$

Tuesday, 16 November 2021 21:50 (20 minutes)

Presenter: GONZÁLEZ-SOLÍS, Sergi (Indiana University)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: **90**

Type: **not specified**

The JLab Eta Factory (JEF) experiment

Tuesday, 16 November 2021 22:10 (20 minutes)

Presenter: SOMOV, Alexander (Jefferson Lab)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: 91

Type: **not specified**

The radiative decay width measurement of the η -meson at GlueX

Tuesday, 16 November 2021 22:30 (20 minutes)

Presenter: JAEGLE, Igal (Jefferson Lab)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: 92

Type: **not specified**

Measurement of the π^0 Transition Form Factor via the Primakoff effect

Tuesday, 16 November 2021 22:50 (20 minutes)

Presenter: LARIN, Ilya (UMASS Amherst)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: 93

Type: **not specified**

Pion electronic decay and lepton universality

Tuesday, 16 November 2021 23:10 (20 minutes)

Presenter: POČANIĆ, Dinko (University of Virginia)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: 94

Type: **not specified**

Measurements of charged and neutral pion electromagnetic polarizabilities at GlueX

Tuesday, 16 November 2021 23:30 (20 minutes)

Presenter: MISKIMEN, Rory (University of Massachusetts)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: 95

Type: **not specified**

Talk to be presented at the 2nd session on Nov.17th

Tuesday, 16 November 2021 19:10 (20 minutes)

Presenter: ANTONELLI, Antonella (INFN)

Session Classification: Goldstone Boson Dynamics (Chair: Bastian Kubis)

Contribution ID: 96

Type: **not specified**

Short-distance constraints in hadronic-light-by-light for the muon $g-2$

Wednesday, 17 November 2021 18:30 (20 minutes)

Presenter: BIJNENS, Johan (Lund University)

Session Classification: Goldstone Boson Dynamics (Chair: Andrzej Kupsc)

Contribution ID: 97

Type: **not specified**

HVP contribution to $g-2$

Wednesday, 17 November 2021 18:50 (20 minutes)

Presenter: HOFERICHTER, Martin (University of Bern)

Session Classification: Goldstone Boson Dynamics (Chair: Andrzej Kupsc)

Contribution ID: 98

Type: **not specified**

Overview of the EFT epsilon' /epsilon

Wednesday, 17 November 2021 19:10 (20 minutes)

Presenter: GISBERT, Hector (TU Dortmund)

Session Classification: Goldstone Boson Dynamics (Chair: Andrzej Kupsc)

Contribution ID: 99

Type: **not specified**

OPE of Green functions and its phenomenological applications

Wednesday, 17 November 2021 19:30 (20 minutes)

Presenter: KADAVY, Tomas (Charles University)

Session Classification: Goldstone Boson Dynamics (Chair: Andrzej Kupsc)

Contribution ID: **100**

Type: **not specified**

Dilaton chiral perturbation theory and applications

Wednesday, 17 November 2021 19:50 (20 minutes)

Presenter: SHAMIR, Yigal (Tel Aviv university)

Session Classification: Goldstone Boson Dynamics (Chair: Andrzej Kupsc)

Contribution ID: **101**

Type: **not specified**

NNLO Positivity Bounds on Chiral Perturbation Theory for a General Number of Flavours

Wednesday, 17 November 2021 20:10 (20 minutes)

Presenter: Sjö, Mattias (Lund University)

Session Classification: Goldstone Boson Dynamics (Chair: Andrzej Kupsc)

Contribution ID: **102**

Type: **not specified**

Three-pion scattering in the Chiral Perturbation Theory

Wednesday, 17 November 2021 20:30 (20 minutes)

Presenter: HUSEK, Tomáš (IFIC (UV-CSIC))

Session Classification: Goldstone Boson Dynamics (Chair: Andrzej Kupsc)

Contribution ID: **103**

Type: **not specified**

The REDTOP experiment: a low energy meson factory to explore dark matter and physics beyond the Standard model

Wednesday, 17 November 2021 21:30 (20 minutes)

Presenter: Dr GATTO, Corrado (INFN & NIU)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gasparian)

Contribution ID: **104**

Type: **not specified**

ChPT and BSM dark mesons

Wednesday, 17 November 2021 21:50 (20 minutes)

Presenter: LI, Lingfeng (Brown University)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gaspasrian)

Contribution ID: **105**

Type: **not specified**

Review of Lattice QCD Calculations for the kaon decays

Wednesday, 17 November 2021 21:10 (20 minutes)

Presenter: KELLY, Christopher (Colmbia University)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gasparian)

Contribution ID: **106**

Type: **not specified**

A dispersive analysis of low energy pion photo- and electroproduction

Wednesday, 17 November 2021 22:10 (20 minutes)

Presenter: CAO, Xiong-Hui (Peking University)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gasparian)

Contribution ID: **107**

Type: **not specified**

The dispersive analysis of pip and piK scattering

Wednesday, 17 November 2021 22:30 (20 minutes)

Presenter: DEINEKA, Olekandra (Johannes Gutenberg University Mainz)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gasparian)

Contribution ID: **108**

Type: **not specified**

Dispersive determination of low energy pi K interactions

Wednesday, 17 November 2021 23:30 (20 minutes)

Presenter: Dr RODAS, ARKAITZ (College of William and Mary)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gasparian)

Contribution ID: **109**

Type: **not specified**

Dispersive analysis of the Primakoff reaction **$\gamma K \rightarrow K\pi$**

Presenter: STAMEN, Dominik (Bonn University)

Contribution ID: **110**

Type: **not specified**

Pion-mass dependence of $\gamma(^*) \pi \rightarrow \pi \pi$ and $\pi \rightarrow \pi \pi$

Wednesday, 17 November 2021 22:50 (20 minutes)

Presenter: NIEHUS, Malwin (HISKP)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gasparian)

Contribution ID: 111

Type: **not specified**

Nuclear forces in a manifestly Lorentz-invariant formulation of chiral EFT

Thursday, 18 November 2021 19:00 (40 minutes)

Presenter: REN, Xiulei (Johannes Gutenberg-Universität Mainz)

Session Classification: Plenary session(Chair: Evgeny Epelbaum)

Contribution ID: 112

Type: **not specified**

Precision tests of fundamental physics with eta and eta' mesons

Thursday, 18 November 2021 19:40 (40 minutes)

Presenter: KUBIS, Bastian (Bonn University)

Session Classification: Plenary session(Chair: Evgeny Epelbaum)

Contribution ID: 113

Type: **not specified**

An overview of the recent and ongoing activities by the LENPIC

Thursday, 18 November 2021 20:20 (40 minutes)

Presenter: KREBS, Hermann (Bochum Univ.)

Session Classification: Plenary session(Chair: Evgeny Epelbaum)

Contribution ID: **114**

Type: **not specified**

Spin sum rules and polarizabiities at long distance

Thursday, 18 November 2021 21:20 (40 minutes)

Presenter: SLIFER, Karl (Univ. of New Hampshire)

Session Classification: Plenary session (Chair: Mohammad W. Ahmed)

Contribution ID: 115

Type: **not specified**

An Overview of Recent and Ongoing Developments in Lattice Simulations of Nuclei

Thursday, 18 November 2021 22:00 (40 minutes)

Presenter: LEE, Dean (Michigan State Univ.)

Session Classification: Plenary session (Chair: Mohammad W. Ahmed)

Contribution ID: **116**

Type: **not specified**

Overview of experimental study on nucleon structure

Thursday, 18 November 2021 22:40 (40 minutes)

Presenter: MEZIANI, Zein-Eddine (Argonne National Laboratory)

Session Classification: Plenary session (Chair: Mohammad W. Ahmed)

Contribution ID: 117

Type: **not specified**

Weak transitions in light nuclei using chEFT interactions and currents

Thursday, 18 November 2021 23:20 (39 minutes)

Presenter: PASTORE, Saori (Washington Univ.)

Session Classification: Plenary session (Chair: Mohammad W. Ahmed)

Contribution ID: **118**

Type: **not specified**

An overview on muonic spectroscopy with light nuclei at CREMA

Friday, 19 November 2021 19:00 (40 minutes)

Presenter: POHL, Randolph (Johannes Gutenberg-Universität)

Session Classification: Plenary session (Chair: Johan Bijmens)

Contribution ID: **119**

Type: **not specified**

Precision calculations of charge radii of light nuclei

Friday, 19 November 2021 19:40 (40 minutes)

Presenter: FILIN, Arseniy (Bochum University)

Session Classification: Plenary session (Chair: Johan Bijnens)

Contribution ID: **120**

Type: **not specified**

Three-nucleon interaction in pionless and pionfull effective field theory

Friday, 19 November 2021 20:20 (40 minutes)

Presenter: GIRLANDA, Luca (INFN)

Session Classification: Plenary session (Chair: Johan Bijnens)

Contribution ID: **121**

Type: **not specified**

Nuclear Equation of State with EFT uncertainties rigorously quantified

Friday, 19 November 2021 21:20 (40 minutes)

Presenter: DRISCHLER, Christian (Michigan State University)

Session Classification: Plenary session (Chair: Hanqing Zheng)

Contribution ID: **122**

Type: **not specified**

The PrimEx result on the π^0 lifetime

Friday, 19 November 2021 22:00 (40 minutes)

Presenter: GASPARIAN, Ashot (North Carolina A& T State Univ.)

Session Classification: Plenary session (Chair: Hanqing Zheng)

Contribution ID: **123**

Type: **not specified**

Hadronic Light-by-light contribution to muon (g-2)

Friday, 19 November 2021 22:40 (40 minutes)

Presenters: JIN, Luchang; JIN, Luchang (University of Connecticut)

Session Classification: Plenary session (Chair: Hanqing Zheng)

Contribution ID: **124**

Type: **not specified**

Closing speech

Friday, 19 November 2021 23:30 (10 minutes)

Presenter: FANG, Shuangshi (IHEP)

Session Classification: Plenary session (Chair: Hanqing Zheng)

Contribution ID: **125**

Type: **not specified**

Opening Speech

Contribution ID: 126

Type: **not specified**

Analysis of Tcc including chiral dynamics and three-body cut

Tuesday, 16 November 2021 18:30 (20 minutes)

Presenter: Dr DU, Meng-Lin (IFIC, Valencia U.)

Session Classification: Hadron Structure (Chair: Fengkun Guo)

Contribution ID: 127

Type: **not specified**

Detecting the pure triangle singularity effect through the ψ (2S) decay

Tuesday, 16 November 2021 18:50 (20 minutes)

Presenter: WU, jiajun (UCAS)

Session Classification: Hadron Structure (Chair: Fengkun Guo)

Contribution ID: **128**

Type: **not specified**

Possible bound states of $J/\psi J/\psi$ from dispersion relation

Tuesday, 16 November 2021 19:10 (20 minutes)

Presenter: DONG, Xiangkun (ITP)

Session Classification: Hadron Structure (Chair: Fengkun Guo)

Contribution ID: 129

Type: **not specified**

Resonance dynamics in U(3) chiral theory and its relevance in the determination of light-quark mass

Tuesday, 16 November 2021 19:30 (20 minutes)

Presenter: GUO, Zhihui (Southeast University)

Session Classification: Hadron Structure (Chair: Fengkun Guo)

Contribution ID: **130**

Type: **not specified**

On QCD contribution to vacuum energy

Presenter: GEGELIA, Jambul (R)

Contribution ID: **131**

Type: **not specified**

The Pole Counting Rule and X, Y, Z States

Tuesday, 16 November 2021 20:10 (20 minutes)

Presenter: CHEN, hao (Peking Univ.)

Session Classification: Hadron Structure (Chair: Fengkun Guo)

Contribution ID: 132

Type: **not specified**

Two-particle scattering from finite-volume quantization conditions using the plane wave basis

Presenter: MENG, Lu (Ruhr University Bochum)

Contribution ID: 133

Type: **not specified**

Low-energy Kbar N interaction from high-energy nuclear collision

Tuesday, 16 November 2021 19:50 (20 minutes)

Presenter: KAMIYA, Yuki (Helmholtz-Institut für Strahlen- und Kernphysik)

Session Classification: Hadron Structure (Chair: Fengkun Guo)

Contribution ID: 134

Type: **not specified**

Results on spin sum rules and polarizabilities at low Q^2

Tuesday, 16 November 2021 21:10 (20 minutes)

Presenter: DEUR, Alexandre (Thomas Jefferson National Accelerator Facility)

Session Classification: Hadron Structure (Chair: Jianping Chen)

Contribution ID: 135

Type: **not specified**

Chiral EFT of nucleon polarizabilities and two-photon-exchange corrections in muonic hydrogen

Tuesday, 16 November 2021 21:30 (20 minutes)

Presenter: PASCALUTSA, Vladimir (Mainz Univ.)

Session Classification: Hadron Structure (Chair:Jianping Chen)

Contribution ID: **136**

Type: **not specified**

Compton Scattering and nucleon polarizabilities

Tuesday, 16 November 2021 21:50 (20 minutes)

Presenter: GRIESSHAMMER, Harald (George Washington Universit)

Session Classification: Hadron Structure (Chair:Jianping Chen)

Contribution ID: 137

Type: **not specified**

The nucleon sigma term from Lattice QCD

Tuesday, 16 November 2021 22:10 (20 minutes)

Presenter: GUPTA, Rajan (Los Alamos National Laboratory)

Session Classification: Hadron Structure (Chair:Jianping Chen)

Contribution ID: **138**

Type: **not specified**

Compton scattering from HIGS

Tuesday, 16 November 2021 22:30 (20 minutes)

Presenter: AHMED, Mohammad (North Carolina Central University)

Session Classification: Hadron Structure (Chair:Jianping Chen)

Contribution ID: **139**

Type: **not specified**

Recent global fit from JAM collaboration

Tuesday, 16 November 2021 22:50 (20 minutes)

Presenter: ZHOU, Yiyu (South China Normal University)

Session Classification: Hadron Structure (Chair:Jianping Chen)

Contribution ID: 140

Type: **not specified**

A Low Q^2 Measurement of the Proton Spin Structure Function g_2^p

Tuesday, 16 November 2021 23:10 (20 minutes)

Presenter: RUTH, David (University of New Hampshire)

Session Classification: Hadron Structure (Chair:Jianping Chen)

Contribution ID: **141**

Type: **not specified**

A high-precision covariant chiral nucleon-nucleon interaction

Tuesday, 16 November 2021 18:30 (20 minutes)

Presenter: GENG, Lisheng (Beijing Univ)

Session Classification: Few-Body Dynamics (Chair: John Bulava)

Contribution ID: **142**

Type: **not specified**

Neutron-antineutron oscillation and the deuteron lifetime

Tuesday, 16 November 2021 18:50 (20 minutes)

Presenter: Dr LONG, Bingwei (Sichuan University)

Session Classification: Few-Body Dynamics (Chair: John Bulava)

Contribution ID: 143

Type: **not specified**

Explicit renormalization of the nucleon-nucleon interaction in chiral EFT and non-perturbative effects

Tuesday, 16 November 2021 19:10 (20 minutes)

Presenter: GASPARYAN, Ashot (Ruhr University of Bochum)

Session Classification: Few-Body Dynamics (Chair: John Bulava)

Contribution ID: 144

Type: **not specified**

Constraints on the Lambda-neutron interaction from charge symmetry breaking of $A=4$ hypernuclei

Tuesday, 16 November 2021 19:30 (20 minutes)

Presenter: NOGGA, Andreas (Forschungszentrum Juelich GmbH)

Session Classification: Few-Body Dynamics (Chair: John Bulava)

Contribution ID: 145

Type: **not specified**

Nuclear structure corrections in muonic atoms from chiral effective field theory

Tuesday, 16 November 2021 19:50 (20 minutes)

Presenter: LI MULI, Simone (Johannes Gutenberg University Mainz)

Session Classification: Few-Body Dynamics (Chair: John Bulava)

Contribution ID: **146**

Type: **not specified**

(speaker not available)

Tuesday, 16 November 2021 20:10 (20 minutes)

Session Classification: Few-Body Dynamics (Chair: John Bulava)

Contribution ID: 147

Type: **not specified**

Fast & rigorous constraints on chiral three-nucleon forces from few-body observables

Tuesday, 16 November 2021 20:30 (20 minutes)

Presenter: PHILLIPS, Daniel (Ohio University)

Session Classification: Few-Body Dynamics (Chair: John Bulava)

Contribution ID: 148

Type: **not specified**

Towards the Study of Short Range Correlations in Radioactive Nuclei: The transparent nucleus in inverse kinematics

Tuesday, 16 November 2021 21:10 (20 minutes)

Presenter: KAHLBOW, Julian (MIT)

Session Classification: Few-Body Dynamics (Chair: Lisheng Geng)

Contribution ID: **149**

Type: **not specified**

Fluctuations and phases in baryonic matter

Tuesday, 16 November 2021 21:30 (20 minutes)

Presenter: BRANDES, Len (Technical University of Munich)

Session Classification: Few-Body Dynamics (Chair: Lisheng Geng)

Contribution ID: 150

Type: **not specified**

A=4-7 Xi hypernuclei based on interactions from chiral effective field theory

Tuesday, 16 November 2021 22:10 (20 minutes)

Presenter: LE, Hoai (Forschungszentrum Juelich)

Session Classification: Few-Body Dynamics (Chair: Lisheng Geng)

Contribution ID: 151

Type: **not specified**

Probing few-body nuclear dynamics via ^3H and ^3He $(e,e' p)$ cross-section measurements at Jefferson Lab

Tuesday, 16 November 2021 22:30 (20 minutes)

Presenter: CRUZ-TORRES, Reynier (LBNL, Berkeley)

Session Classification: Few-Body Dynamics (Chair: Lisheng Geng)

Contribution ID: **152**

Type: **not specified**

Two-nucleon with various methods

Presenter: WALKER-LOUD, Andre (LBNL)

Contribution ID: **153**

Type: **not specified**

Extracting few-body matrix elements from lattice QCD

Tuesday, 16 November 2021 22:50 (20 minutes)

Presenter: ILLA, Marc (University of Washington)

Session Classification: Few-Body Dynamics (Chair: Lisheng Geng)

Contribution ID: 154

Type: **not specified**

Strangeness $S=-3$ and -4 baryon-baryon interactions in chiral effective field theory

Tuesday, 16 November 2021 21:50 (20 minutes)

Presenter: Dr HAIDENBAUER, Johann (Forschungszentrum Juelich GmbH)

Session Classification: Few-Body Dynamics (Chair: Lisheng Geng)

Contribution ID: **155**

Type: **not specified**

Talk 15

Contribution ID: **156**

Type: **not specified**

Overview of TMD studies

Wednesday, 17 November 2021 18:30 (20 minutes)

Presenter: LIU, Tianbo (Shandong University)

Session Classification: Hadron Structure (Chair: Hanqing Zheng)

Contribution ID: **157**

Type: **not specified**

Overview of GPD Studies

Wednesday, 17 November 2021 18:50 (20 minutes)

Presenter: YE, Zhihong (Tsinghua University)

Session Classification: Hadron Structure (Chair: Hanqing Zheng)

Contribution ID: **158**

Type: **not specified**

Structure-dependent electromagnetic finite-size effects

Wednesday, 17 November 2021 19:10 (20 minutes)

Presenter: HERMANSSON-TRUEDSSON, Nils (University of Bern)

Session Classification: Hadron Structure (Chair: Hanqing Zheng)

Contribution ID: **159**

Type: **not specified**

The N/D study on the singularity structure of πN scattering amplitudes

Wednesday, 17 November 2021 19:30 (20 minutes)

Presenter: LI, Quzhi (Peking Univ.)

Session Classification: Hadron Structure (Chair: Hanqing Zheng)

Contribution ID: **160**

Type: **not specified**

Hadron physics results at KLOE-2 experiment

Presenter: MANDAGLIO, Giuseppe (University of Messina)

Contribution ID: **161**

Type: **not specified**

Chiral EFT of nucleons and pions in the presence of external gravitational field

Wednesday, 17 November 2021 19:50 (20 minutes)

Presenter: ALHARAZIN, Herzallah (Institut für Theoretische Physik II)

Session Classification: Hadron Structure (Chair: Hanqing Zheng)

Contribution ID: **162**

Type: **not specified**

New insights into the nucleon form factor and proton radius from dispersive analysis

Wednesday, 17 November 2021 20:10 (20 minutes)

Presenter: LIN, Yonghui (Bonn University)

Session Classification: Hadron Structure (Chair: Hanqing Zheng)

Contribution ID: **163**

Type: **not specified**

Proton radius with MUSE experiment

Wednesday, 17 November 2021 20:30 (20 minutes)

Presenter: STRAUCH, Steffen (University of South Carolina)

Session Classification: Hadron Structure (Chair: Hanqing Zheng)

Contribution ID: **164**

Type: **not specified**

d/u at high-x from JLab MARATHON experiment

Presenter: NYCZ, Michael (Virginia Univ.)

Contribution ID: **165**

Type: **not specified**

Pion photoproduction in CHPT

Presenter: BLIN, Astrid Hiller (Jlab)

Contribution ID: **166**

Type: **not specified**

BSM search and hadron structure from parity violating electron scattering.

Wednesday, 17 November 2021 21:10 (20 minutes)

Presenter: Dr ZHENG, Xiaochao (University of Virginia)

Session Classification: Hadron Structure (Chair: Harald Griesshammer)

Contribution ID: **167**

Type: **not specified**

Isospin breaking effects in radiative corrections to the axial charge in neutron beta decay

Wednesday, 17 November 2021 21:30 (20 minutes)

Presenter: HAYEN, Leendert (NCSU)

Session Classification: Hadron Structure (Chair: Harald Griesshammer)

Contribution ID: **168**

Type: **not specified**

Baryon masses and currents in SU(3) BChPT x 1/Nc

Wednesday, 17 November 2021 21:50 (20 minutes)

Presenter: FERNANDO, Ishara (University of Virginia)

Session Classification: Hadron Structure (Chair: Harald Griesshammer)

Contribution ID: **169**

Type: **not specified**

Two-loop calculation of the nucleon self-energy

Wednesday, 17 November 2021 22:10 (20 minutes)

Presenter: CONRAD, Nils Dirk (Ruhr-Universität Bochum)

Session Classification: Hadron Structure (Chair: Harald Griesshammer)

Contribution ID: 170

Type: **not specified**

The nucleon sigma term from Lattice QCD

Presenter: GUPTA, Rajan (LANL)

Contribution ID: 171

Type: **not specified**

Review of the experimental activity at RIKEN to explore the three nucleon interactions

Wednesday, 17 November 2021 18:30 (20 minutes)

Presenter: SEKIGUCHI, Kimiko (Tohoku University)

Session Classification: Few-Body Dynamics (Chair: Xu Feng)

Contribution ID: 172

Type: **not specified**

Three-nucleon force from lattice QCD

Wednesday, 17 November 2021 18:50 (20 minutes)

Presenter: DOI, Takumi (RIKEN))

Session Classification: Few-Body Dynamics (Chair: Xu Feng)

Contribution ID: 173

Type: **not specified**

Pion-nucleon scattering from lattice QCD

Wednesday, 17 November 2021 19:10 (20 minutes)

Presenter: BULAVA, John (DESY)

Session Classification: Few-Body Dynamics (Chair: Xu Feng)

Contribution ID: 174

Type: **not specified**

Most charming dibaryon from lattice QCD

Wednesday, 17 November 2021 19:30 (20 minutes)

Presenter: LYU, Yan (school of physics Peking University)

Session Classification: Few-Body Dynamics (Chair: Xu Feng)

Contribution ID: 175

Type: **not specified**

Three pion and kaon scattering from lattice QCD

Wednesday, 17 November 2021 19:50 (20 minutes)

Presenter: CULVER, Christopher (University of Liverpool)

Session Classification: Few-Body Dynamics (Chair: Xu Feng)

Contribution ID: 176

Type: **not specified**

A1(1260) from lattice QCD

Wednesday, 17 November 2021 20:10 (20 minutes)

Presenter: Dr MAI, Maxim (GWU)

Session Classification: Few-Body Dynamics (Chair: Xu Feng)

Contribution ID: 177

Type: **not specified**

Two-particle scattering from finite-volume quantization conditions using the plane wave basis

Wednesday, 17 November 2021 20:30 (20 minutes)

Presenter: MENG, Lu (Ruhr University Bochum)

Session Classification: Few-Body Dynamics (Chair: Xu Feng)

Contribution ID: 178

Type: **not specified**

Deuteron VVCS and nuclear structure effects in muonic deuterium at N3LO in pionless EFT

Wednesday, 17 November 2021 21:10 (20 minutes)

Presenter: LENSKY, Vadim (JGU, Mainz)

Session Classification: Few-Body Dynamics (Chair: Emile Passemar)

Contribution ID: 179

Type: **not specified**

The Generalized GDH Sum Rule: Measuring the ^3He Spin Structure at Low Q^2

Wednesday, 17 November 2021 21:30 (20 minutes)

Presenter: PENG, Chao (Argonne National Laboratory)

Session Classification: Few-Body Dynamics (Chair: Emile Passemar)

Contribution ID: **180**

Type: **not specified**

Renormalization of CP-violating nuclear forces

Wednesday, 17 November 2021 21:50 (20 minutes)

Presenter: SHAIN PORUVELIL, Sachin (Massachusetts U)

Session Classification: Few-Body Dynamics (Chair: Emile Passemar)

Contribution ID: **181**

Type: **not specified**

Large-Nc constraints for Beyond the Standard Model few-nucleon currents in effective field theory

Wednesday, 17 November 2021 22:10 (20 minutes)

Presenter: RICHARDSON, Thomas (Duke Univ)

Session Classification: Few-Body Dynamics (Chair: Emile Passemar)

Contribution ID: **182**

Type: **not specified**

Nucleon-level Effective Theory of $\mu \rightarrow e$ Conversion

Wednesday, 17 November 2021 22:30 (20 minutes)

Presenter: RULE, Evan (UC, Berkeley)

Session Classification: Few-Body Dynamics (Chair: Emile Passemar)

Contribution ID: **183**

Type: **not specified**

Chiral EFT for neutrinoless double beta decay

Wednesday, 17 November 2021 22:50 (20 minutes)

Presenter: DEKENS, Wouter (UC San Diego)

Session Classification: Few-Body Dynamics (Chair: Emile Passemar)

Contribution ID: **184**

Type: **not specified**

Talk 14

Contribution ID: **185**

Type: **not specified**

Talk 15

Contribution ID: 186

Type: **not specified**

The Generalized GDH Sum Rule: Measuring the ^3He Spin Structure at Low Q^2

The Gerasimov-Drell-Hearn (GDH) sum rule, as a fundamental relation between real photon absorption and the anomalous magnetic moment to the spin structure of the target. The generalized form of GDH sum rule extends this relation to finite four-momentum transfer squared (Q^2). Jefferson Lab experiment E97-110 was carried out with longitudinally polarized electron beam scattering off the polarized ^3He target in Hall A. The experiment measured the spin dependent structure functions g_1 and g_2 for ^3He at small scattering angles of 6° and 9° with a beam energy from 1.1 GeV to 4.4 GeV. The generalized GDH sum and moments of the spin dependent structure functions were extracted by integrating the measured g_1 and g_2 over the quasi-elastic and resonance region, and beyond at low Q^2

from 0.02 to 0.3 GeV^2 . The data in this low Q^2 region benchmarks predictions regarding the neutron spin structure by Chiral Perturbation Effective Field Theories. Recovery of the real-photon point ($Q^2 \rightarrow 0$) for the generalized GDH sum rule is also tested with the low Q^2 results. In this talk, we will present the experimental results of various moments for ^3He .

Primary author: PENG, Chao (Argonne National Laboratory)

Presenter: PENG, Chao (Argonne National Laboratory)

Contribution ID: **187**

Type: **not specified**

Study of deuteron-dark matter scattering within the χ EFT framework

Presenter: FILANDRI, Elena (University of Trento)

Contribution ID: **188**

Type: **not specified**

The Muon Proton Scattering Experiment (MUSE) at PSI

Contribution ID: **189**

Type: **not specified**

The Generalized GDH Sum Rule: Measuring the ^3He Spin Structure at Low Q^2

Presenter: PENG, Chao (A)

Contribution ID: 190

Type: **Parallel-Hadron Structure**

Nucleon Polarizabilities from Few-Nucleon Systems

An Update Low-energy Compton scattering probes the nucleon's two-photon response to electric and magnetic fields. It tests the symmetries and strengths of the interactions between constituents, and with photons. For convenience, this energy-dependent information is often compressed into the two scalar dipole polarizabilities α_{E1} and β_{M1} at zero photon energy. In addition, spin polarizabilities are particularly interesting since they parametrize the stiffness of the spin in external electro-magnetic fields (nucleonic Faraday effect) and probe the spin-dependent pion-nucleon interaction. Combined with emerging lattice QCD computations, polarizabilities provide stringent tests of chiral symmetry and hadron structure. Compton scattering in light nuclei also tests the chiral symmetry of the charged pion-exchange contribution to nuclear binding. %This talk focuses on the first-ever description of 4He. It uses the transition-density formalism, an efficient and general method for calculating interactions of external probes with light nuclei. One- and two-body transition densities that encode the nuclear structure of the target are evaluated once and stored. They are then convoluted with an interaction kernel to produce observables. The same densities can be used with different kernels for any reaction in which a probe interacts perturbatively with the target. The method exploits factorization between nuclear structure and interaction kernel in Chiral EFT at energies $\sim m_\pi$. It takes full advantage of the numerical power of modern few-nucleon methods, is markedly more computationally efficient and applicable to a wide array of nuclei and reactions. %The 4He Compton results converge well order-by-order and are in good agreement with the data between 50 and 90 MeV, with the expected residual dependence on the NN and 3N potential. However, data appear noisier than the reported experimental uncertainties allow for, so an extraction of the polarizabilities is not attempted.

Primary author: W. GRIESSHAMMER, Harald (George Washington University)

Presenter: W. GRIESSHAMMER, Harald (George Washington University)

Contribution ID: 191

Type: **Parallel-Goldstone Boson**

Pion electronic decay and lepton universality

Involving only light elementary particles, charged pion decays are characterized by simple dynamics, few available decay channels, and extremely well controlled radiative and loop corrections. The comparative simplicity of pion (and muon) decays allows them to be described with unprecedented precision within the Standard Model, (SM), typically with relative uncertainties of $\sim 10^{-4}$ or lower, reflecting the strong influence of the applicable SM symmetries. Given the highly precise SM theoretical description, pion decays are used as selective and sensitive probes of SM parameters, and of possible SM extensions. In particular, $V-A$ helicity suppression of the $\pi^+ \rightarrow e^+ \nu_e(\gamma)$, or $\pi e2(\gamma)$ decay amplifies the sensitivity to pseudoscalar terms by a factor of ~ 8000 , enabling indirect searches for non-SM pseudoscalar terms, as well as scalar and tensor terms, through loop effects, with good sensitivity to interesting regions of the beyond-SM parameter space, such as supersymmetric extensions.

Experimental data on the allowed rare electronic decay of charged pions: $\pi e2(\gamma)$, the muon radiative decay, $\mu^+ \rightarrow e^+ \nu_e \nu_\mu \gamma$ as well as the semileptonic, $\pi^+ \rightarrow \pi^0 e^+ \nu_e(\gamma)$, or $\pi e3(\gamma)$ decay, present an internally consistent picture, so far in excellent agreement with SM predictions. However, experimental accuracy is lagging behind that of the theoretical description for all above processes. The PEN experiment at PSI is studying the $\pi e2(\gamma)$ decay with the primary goal to reach a relative precision of 5×10^{-4} in $R_{\pi e/\mu}$, its branching ratio. We review the current status of the field, the PEN research program, the present status of the PEN data analysis, and the expected uncertainty limits.

Primary author: POČANIĆ, Dinko (University of Virginia)

Presenter: POČANIĆ, Dinko (University of Virginia)

Contribution ID: 192

Type: **Parallel-Few-Body Physics**

Two-nucleons with various methods

Lattice QCD calculations of two-nucleon systems have suffered from a significant systematic uncertainty for more than a decade. Two different methods of computing the interaction energy have yielded qualitatively different solutions: the use of local creation operators yields deeply bound di-nucleon systems, while calculations that use momentum space creation operators or the HAL QCD potential observe both di-nucleon systems to be unbound. These observations have been made with heavy pions masses where the stochastic signal-to-noise is exponentially larger than at light pion masses. This is not an academic exercise, but a challenge that must be overcome in order for the lattice QCD and broader physics communities to have confidence in lattice QCD calculations of two-nucleon (two-baryon) interactions and two-nucleon electroweak matrix elements. I will present progress that is being made in computing the two-nucleon system with all methods in use, the local creation operators, momentum space creation operators and the HAL QCD potential, for the first time on the same set of gauge configurations. This will enable us to isolate the discrepancy in the spectrum to the method used, and hopefully, resolve this long-standing discrepancy.

Primary author: WALKER-LOUD, Andre (LBNL)

Presenter: WALKER-LOUD, Andre (LBNL)

Contribution ID: **193**

Type: **not specified**

Improved Standard-Model prediction for the dilepton decay of the neutral pion (20')

Presenter: HOID, Bai-Long (Universität Bern)

Contribution ID: **194**

Type: **not specified**

Determining the nucleon mass and sigma term from lattice QCD

Wednesday, 17 November 2021 22:30 (20 minutes)

Presenter: MILLER, Nolan (University of North Carolina, Chapel Hill)

Session Classification: Hadron Structure (Chair: Harald Griesshammer)

Contribution ID: **195**

Type: **not specified**

Hadron physics results at KLOE-2 experiment

Presenter: P. D. RIO, Elena (INFN)

Contribution ID: **196**

Type: **not specified**

VCS and generalized polarizabilities from Mainz and JLab: Overview and new results

Wednesday, 17 November 2021 22:50 (20 minutes)

Presenter: SPARVERIS, Nikos (Temple University)

Session Classification: Hadron Structure (Chair: Harald Griesshammer)

Contribution ID: 197

Type: **Parallel-Few-Body Physics**

Large- N_c constraints for Beyond the Standard Model few-nucleon currents in effective field theory

Low energy experiments that search for Beyond the Standard Model (BSM) physics often rely on nuclear targets.

Therefore, it is imperative that we obtain a clear theoretical picture of the nuclear physics involved. Effective field theory (EFT) provides a model-independent framework to capture the nuclear physics in terms of few-nucleon currents.

However, every operator in an EFT is accompanied by an undetermined low energy coefficient that must be determined from data or a nonperturbative quantum chromodynamics (QCD) calculation such as a lattice calculation.

For many processes, these determinations are not yet possible; thus, other theoretical constraints are necessary in order to guide the interpretation of experimental bounds.

Here, we review recent constraints obtained from the large- N_c limit of QCD, where N_c is the number of colors, for BSM few-nucleon currents relevant for neutrinoless double beta decay and dark matter direct detection.

Primary author: R. RICHARDSON, Thomas (Duke University)

Presenter: R. RICHARDSON, Thomas (Duke University)

Contribution ID: **198**

Type: **not specified**

Status of π^0 , η and η' experimental program in KLOE-II

Tuesday, 16 November 2021 18:30 (20 minutes)

Presenter: P. D. RIO, Elena (INFN)

Session Classification: Goldstone Boson Dynamics (Chair: Bastian Kubis)

Contribution ID: **199**

Type: **not specified**

Improved Standard-Model prediction for the dilepton decay of the neutral pion(20')

Tuesday, 16 November 2021 23:50 (9 minutes)

Presenter: HOID, Bai-Long (Universität Bern)

Session Classification: Goldstone Boson Dynamics (Chair: Liping Gan)

Contribution ID: **200**

Type: **not specified**

On QCD contribution to vacuum energy

Tuesday, 16 November 2021 20:30 (20 minutes)

Presenter: GEGELIA, Jambul (Ruhr-University-Bochum)

Session Classification: Hadron Structure (Chair: Fengkun Guo)

Contribution ID: **201**

Type: **not specified**

Bochum: Venue of the CD2024

Friday, 19 November 2021 23:20 (10 minutes)

Presenter: EPELBAUM, Evgeny (Ruhr-Universitaet Bochum)

Session Classification: Plenary session (Chair: Hanqing Zheng)

Contribution ID: 202

Type: **Parallel-Hadron Structure**

Isospin breaking effects in radiative corrections to the axial charge in neutron beta decay

Precision searches in neutron beta decay have been at the inception and continued testing of the Standard Model, fueled in part by progress in electroweak radiative corrections. Recently, a substantial shift in the evaluation of the so-called γW box have caused a significant shift in V_{ud} , the up-down CKM matrix element, inspiring substantial additional research including radiative corrections to the weak axial charge. The latter is additionally a clean channel when looking for exotic right-handed currents by comparing to precision lattice QCD results. As additional electroweak radiative corrections can mimic New Physics, we have performed heavy-baryon effective field theory calculations and quantify the effects of explicit isospin symmetry breaking. We will present the current state of the art and report on progress and future paths.

Primary author: HAYEN, Leendert (NCSU)

Presenter: HAYEN, Leendert (NCSU)

Contribution ID: **203**

Type: **not specified**

Two-nucleon with various methods

Wednesday, 17 November 2021 23:10 (20 minutes)

Presenter: WALKER-LOUD, Andre (LBNL)

Session Classification: Few-Body Dynamics (Chair: Emile Passemar)

Contribution ID: 204

Type: **not specified**

Dispersive analysis of the Primakoff reaction $\gamma K \rightarrow K\pi$

Wednesday, 17 November 2021 23:10 (20 minutes)

Presenter: STAMEN, Dominik (Bonn University)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gasparian)

Contribution ID: **205**

Type: **not specified**

Recent results from NA62 (20')

Wednesday, 17 November 2021 23:50 (9 minutes)

Presenter: ANTONELLA, antonelli (INFN)

Session Classification: Goldstone Boson Dynamics (Chair: Ashot Gasparian)