

Analysis of Little Higgs Model with T-parity at ILC

'10 3/28 Y. Takubo (Tohoku U.)

E. Kato(Tohoku U.), E. Asakawa(Meiji-gakuin U.),
K. Fujii(KEK), S. Matsumoto(Toyama U.),
H. Yamamoto (Tohoku U.)

Little hierarchy problem


There are two predictions of the energy scale for new physics (Λ).

- $\Lambda < 1 \text{ TeV}$: <10% fine tuning of Higgs mass.
- $\Lambda > 10 \text{ TeV}$: EW precision measurements
 - The global fit of the EW parameters.
(Γ_Z , M_W/M_Z , $\sin^2\theta_W$, ...)

$$m_h^2 = m_0^2 + \Lambda^2$$

$100^2 \Leftrightarrow 1000^2$

→ There is a discrepancy between two predictions.




Some physics models are proposed to solve little hierarchy problem.

→ **Little Higgs model (with T-parity)**


Little Higgs mechanism

- Higgs is a pseudo NG boson of a global symmetry of SU(5) .
- The symmetry breaks to SO(5) at $\Lambda \sim 10$ TeV.
 - VEV: $f \sim 1\text{TeV}$
 - $[\text{SU}(2)_L \times \text{U}(1)_Y]$ is a subgroup of SO(5).
- The little Higgs partners contribute to cancel quadratic divergent term of M_h at 1-loop level.
 - The new physics at 1 TeV is not necessary.

$$\text{SU}(5): [\text{SU}(2) \times \text{U}(1)]^2$$



$$\text{SO}(5): \text{SU}(2)_L \times \text{U}(1)_Y$$



$$\text{U}(1)_Y$$


→ Little hierarchy problem can be solved.

The diagram shows two Feynman diagrams for the Higgs mass correction. The first diagram is a top quark loop with external Higgs lines, labeled Λ_{SM} . The second diagram is a top quark loop with a top partner T loop, labeled Λ_{NP} . The sum of these two diagrams is zero, indicating the cancellation of quadratic divergences.


Importance of heavy gauge bosons

Heavy gauge bosons

- The heavy gauge bosons appears as the little Higgs partners of SM gauge bosons.
 - $\gamma, Z, W \leftrightarrow A_H, Z_H, W_H$
 - The masses have information of $VEV(f)$.
 - A_H becomes stable, requiring T-parity.
 - A_H is a dark matter candidate.
- **VEV and abundance of the dark matter can be evaluated by measurement of heavy gauge bosons.**



Sensitivity of ILC to the heavy gauge bosons was studied.



Parameter choice for simulation study

Constraint by WMAP result


- The dark matter density was determined by WMAP.
 - $\Omega h^2 = 1.119 \pm 0.009$
- Annihilation xsec of $A_H \sim F(M_h, f)$
 - M_h and f are restricted by Ωh^2 .

Parameters for this study


- f : 580 GeV
- M_h : 134 GeV
- M_{A_H} : 81.9 GeV
- M_{W_H} : 368 GeV
- M_{Z_H} : 369 GeV

A_H , W_H , and Z_H can be observed at ILC.

Annihilation processes of A_H



f v.s. m_h with WMAP constraint






Analysis modes

According to the beam energy at ILC, three analysis modes were selected.

- $A_H + Z_H$ @ $E_{CM} = 500$ GeV
 - xsec: 1.91 fb
 - $M_{A_H} + M_{Z_H} = 450.9$ GeV
- $W_H^+ + W_H^-$ @ $E_{CM} = 1$ TeV
 - xsec: 277 fb
 - $M_{W_H} + M_{W_H} = 736$ GeV

- $Z_H + Z_H$ @ $E_{CM} = 1$ TeV
 - xsec: 277 fb
 - $M_{Z_H} + M_{Z_H} = 736$ GeV



Analysis result for $Z_H Z_H$ is shown.


Simulation study

Simulation condition


- Signal: $Z_H Z_H \rightarrow H H A_H A_H$ (98.0fb)
- BG:
 - WWZ (5.9fb)
 - $\nu\nu WW$ (6.7fb)
 - WW (3,932fb)
 - tt (193fb)

Simulation tools

- Event generator: PhysSim
 - ISR, FSR, beamstrahlung, and beam energy spread are considered.
- The fast-simulator for GLD was used.



Event display of a $W_H W_H$ event





Event reconstruction

- All the events are reconstructed as 4-jet events.
- Two Higgs masses are reconstructed to minimize χ^2 value:


$$\chi^2 = \frac{(M_{H1} - M_H)^2}{\sigma_{MH}^2} + \frac{(M_{H2} - M_H)^2}{\sigma_{MH}^2}$$

- Many BG events contaminate in the signal region.
 → The selection cut was applied.






χ^2 cut

- The χ^2 cut was applied to select well reconstructed events.
 - $\chi^2 < 80$ was selected.
- The main part of WW background can be rejected.





of b-tagged jets

- The b-tagging is applied to select $H \rightarrow bb$.
 - b-tag requirement: 2 tracks with 3 sigma displacement from IP.
- Selection: $N_b \geq 1$
- ➔ tt events become the main background.



Acoplanarity cut

- Acoplanarity distribution is investigated.
 - Acoplanarity = $\phi - \pi$
- The signal has larger acoplanarity because A_H is missing.
 - $t\bar{t}$ events have a sharp peak around 0 degree.
- Acoplanarity > 25 degree is selected.



Cut summary

A number of events at each selection cut are summarized.


- The backgrounds are rejected efficiently.

	$Z_H Z_H$	WWZ	$\nu\nu WW$	WW	tt
Xsec (fb)	98.0	5.9	6.9	3,932	192.9
No cut	49,730	2,961	3,340	894,500	96,470
$\chi^2 < 80$	41,550	871	1,607	144,900	66,680
$N_b \geq 1$	27,980	115	537	13,210	64,120
Acop > 25 deg.	21,360	96	443	8,463	8,354

The masses of A_H and Z_H are extracted.

Extraction of M_{A_H} and M_{Z_H}

- The edge of the Higgs energy distribution has information of masses of A_H and Z_H .
- The masses are extracted by fitting the Higgs energy distribution.
 - M_{A_H} : 81.9 ± 5.1 GeV (True value: 81.9 GeV) → **6.2%**
 - M_{Z_H} : 368.0 ± 5.2 GeV (True value: 369.0 GeV) → **1.4%**



Mass resolution of heavy gauge bosons

The measurement accuracy of the heavy gauge bosons is summarized.

$E_{\text{CM}} = 500\text{GeV}$

- $A_{\text{H}} + Z_{\text{H}}$
 - $M_{A_{\text{H}}} : 83.2 \pm 13.3 \text{ GeV (16.2\%)}$
 - $M_{Z_{\text{H}}} : 366.0 \pm 16.0 \text{ GeV (4.3\%)}$

$E_{\text{CM}} = 1\text{TeV}$

- $W_{\text{H}}^{+} + W_{\text{H}}^{-}$
 - $M_{A_{\text{H}}} : 82.46 \pm 1.18 \text{ GeV (1.4\%)}$
 - $M_{W_{\text{H}}} : 367.8 \pm 0.83 \text{ GeV (0.2\%)}$

} Previous study
(Phys. Rev. D79, 075013 (2009))

- $Z_{\text{H}} + Z_{\text{H}}$
 - $M_{A_{\text{H}}} : 81.9 \pm 5.2 \text{ GeV (6.2\%)}$
 - $M_{Z_{\text{H}}} : 368.0 \pm 5.2 \text{ GeV (1.4\%)}$

Precision measurement of heavy gauge bosons is possible at ILC.

Summary

- Little Higgs model with T-parity is one of the candidate of the new physics.
- Measurement of the heavy gauge bosons is important to confirm Little Higgs model.
- $Z_H Z_H$ is analyzed to extract masses of A_H and Z_H at $E_{CM} = 1$ TeV.
 - Masses of A_H and Z_H can be measured with 6.2% and 1.4% accuracy, respectively, at $E_{CM} = 1$ TeV.
- The simultaneous mass fitting will be performed with $W_H W_H$ and $Z_H Z_H$ analyses.