

Long-Lived Particle Search with Lepton Colliders


Yulei Zhang^[1], Xiang Chen^[1], Jifeng Hu^[2], Liang Li^[1]

1 Shanghai Jiao Tong University

2 South China Normal University



$$e^+e^- \rightarrow Zh \rightarrow \nu\bar{\nu} + SS1 + SS2 \rightarrow \nu\bar{\nu}q\bar{q}q\bar{q}$$



Beyond the Standard Model


- # Long-Lived Particle



The lifetime frontier ...

- Large majority of current collider experiment searches and analysis strategies assume the new particle decays **promptly**.
- Particle lifetimes span a very wide range and **long lifetimes** can generically appear in the **BSM theories**.
- Dedicated searches for long-lived BSM particles are necessary.

Sixth workshop of the LHC LLP Community




What is a long-lived particle?


Object (neutral or charged) decaying a **macroscopic** and **reconstructible** distance from IP

Signal signature of a long-lived particle:



Neutral LLP decays are a spectacular signature, and the **burst of energy** appearing out of nowhere sets it apart from the collision point.



Phys. Rev. Lett. **122**, 131801 – 2019.04.03




LLP Searches at Lepton Colliders



- Energy: 240 GeV
- Mass of SS1: 0.1-50 GeV
- Mass of SS2: 0.1-50 GeV
- SS1, SS2's lifetime $\tau = R_{\text{out}}/c = 6m/c = 20 \text{ ns}$

Basic Setup



- Muon Detector
 - $R_{in} \approx 4m$
 - $R_{out} \approx 6m$
- $\Delta t = t_{Hit} - r_{Hit}/c$
- Dominant Background
 - $e^+e^- \rightarrow ZH$
 - $e^+e^- \rightarrow qq$
- **Full simulation** with CEPC official software



Signal range: LLPs decay within 0~6 meters

ML based Analysis




Deep Residual Network, ResNet

- Firstly, appeared in the ILSVRC 2015 classification challenges (ImageNet Large Scale Visual Recognition Challenge)
- ResNet18, ResNet50, ResNet101...



Revolution of Depth





Configuration


- Mapping the raw detector information to a 2D image
- Input Format: image with resolution of $(R, \phi) = 200 \times 200$ and 1 to 3 channel(s)
 - $R_i = i \times \Delta R_i$, R starts from 0 m to 8 m.
 - ϕ starts from $-\pi$ to π
 - Energy is the sum of both Tracker hits and Calorimeter hits.
 - Time is the maximum ΔT ($E > 0.1 \text{ GeV}$) within (R, ϕ) pixel
- Model: ResNet18 (Classification), ResNet50 (Vertex Finding)
- **Binary Cross Entropy Loss:** $loss(x_i, y_i) = -\omega_i[y_i \log(x_i) + (1 - y_i)\log(1 - x_i)]$



Signal: $e^+e^- \rightarrow Zh \rightarrow q\bar{q} + (SS1 + SS2) \rightarrow q\bar{q}(q\bar{q}v\bar{v})$



Hits from LLP Signal






Hits from Z-boson

Energy




Max. ΔT

2-channel



Background: $e^+e^- \rightarrow q\bar{q}$





Preselection



	Signal	ZH	eeqq
Raw Entries	0.5×10^6	0.93×10^6	0.99×10^7
$50 < m_{qq} < 180 \ \&\& \ 35 < m_{recoil} < 175 \ \&\& \ 0.25 \leq y_{12} \leq 0.72$	3.8×10^5	182,844	848,529
Efficiency	72.41%	19.66%	8.57%



ResNet18 (2 channel: $E, \Delta T$)



	$n_1 = 1 \ \& \ n_2 = 1$	ϵ	Weight
Signal	377742	99.99%	$\times 1$
ZH	0	0.00%	$\times 1$
qq	0	0.00%	$\times 25$

ML-based and Cut-based comparison ($Z \rightarrow \bar{\nu}\nu$)

	Signal: $Z \rightarrow \bar{\nu}\nu$	$e^+e^- \rightarrow q\bar{q}$	$e^+e^- \rightarrow Zh$	Total
# of Events in 5.6 ab^{-1}		2.5×10^8	1.0×10^6	2.01×10^8
# of Events simulated	$\sim 1.0 \times 10^6$	$(\sim 0.99 \times 10^7)^*$	$\sim 1.37 \times 10^6$	$\sim 2.87 \times 10^6$
$E_{\text{missing}} > 190 \text{ GeV}$ && $n_{\text{rec}} < 8$	88,077	290	3,361	3,651
$ML \text{ score} > 0.95$	87,050	0	0	0
Efficiency (ML-based)	98.83%	—	—	—
$E_{2j} \geq 30 \text{ GeV}$ && $\min(\Delta T_{j_1}, \Delta T_{j_2}) > 3 \text{ ns}$	66,325	0	0	0
Efficiency (Cut-based)	73.89%			


* Due to the limited computing power, $\sim 10^7$ events were simulated so far

Expected Limits



	Signal	Total Background	Expected Limits
$e^+e^- \rightarrow Zh \rightarrow (Z: \bar{q}q)\bar{q}q\bar{\nu}\nu$	373308	0.02 (CR)	2.4×10^{-5}
$e^+e^- \rightarrow Zh \rightarrow (Z: \bar{\nu}\nu)\bar{q}q\bar{\nu}\nu$	87,050	0.02 (CR)	9.8×10^{-5}
Combined limit: 1.9×10^{-5}			

- Limits are the minimal branching ratio of Higgs decaying to LLPs (the smaller the better).
- Cosmic Ray(CR) veto efficiency is calculated by the filter that the time difference of two clusters on the outermost cell must be less than 2.4 meters. (signal inefficiency~ 2.1%)
- Signal Yield: $n_s = \mathcal{L} \times \sigma(e^+e^- \rightarrow Zh) \times \sigma(Z \rightarrow qq, \bar{\nu}\nu) \times \epsilon_{sig} \times \epsilon_{CR}$


Sensitivity





- Best branching ratio exclusion limit at decay length around a few meters: $BR(h \rightarrow XX) > \sim 10^{-5}$ for most LLP masses
- Good sensitivity for low LLP mass (as low as 0.1 GeV)



Preliminary Study: Vertex Finding With ML






SmoothL1 Loss: Vertex Finding



Smooth L1 Loss: $loss(x, y) = \frac{1}{n} \sum_{i=1}^n \begin{cases} 0.5 \times (y_i - f(x_i))^2, & \text{if } |y_i - f(x_i)| < 1 \\ |y_i - f(x_i)| - 0.5, & \text{otherwise} \end{cases}$

Preliminary Study: Vertex Finding Results



$$\Delta R = R_{\text{pred}} - R_{\text{truth}} \quad \Delta\phi = \phi_{\text{pred}} - \phi_{\text{truth}}$$

- pixel size: 7.1cm
- Initial result looks very promising

Summary

- Long-Lived Particle ($h \rightarrow q\bar{q}\nu\bar{\nu}$) study done with future lepton collider
 - current results based on CEPC_v4 geometry setup
- For background-free channel (both for ML-based and cut-based), $ZH \rightarrow \bar{\nu}\nu + SS_1 + SS_2$, ML can increase signal efficiency from 73.9% to 98.8%.
- Working on exploring other possibilities with ML: e.g., vertex finding
- First attempt to apply AI image recognition techniques to raw detector hits
 - Very good sensitivity reached ($\sim 10^{-5}$) with (expected) 10^6 Higgs statistics compared to current LHC limits ($\sim 10^{-4}$).



Thanks

Yulei Zhang

✉ zc_1994@163.com