



A ν scalar in the early universe and $(g - 2)_\mu$

Jia Liu (刘佳)

Peking University

With Navin McGinnis, Carlos E.M. Wagner and Xiao-Ping Wang (王小平 北航)

1810.11028 [JHEP 1903 (2019) 008]

2001.06522 [JHEP 2004 (2020) 197]

2110.14665



The 2021 international workshop on the high energy Circular Electron-Positron Collider

11/11/2021

Outline

- The light dark sector and $(g - 2)_{e/\mu}$
- The cosmological triangle
- The ν scalar in the early universe and $(g - 2)_\mu$
- Summary

The heavy and the light



- Dark sector particles
 - New light weakly coupled particles
 - Do not interact with the known strong, weak, or electromagnetic forces
- Today we focus on the light dark sector particles

The news from muon g-2

First results from Fermilab's Muon g-2 experiment strengthen evidence of new physics

April 7, 2021



Media contact

- Tracy Marc, Fermilab, media@fnal.gov, 224-290-7803


The long-awaited [first results](#) from the Muon g-2 experiment at the U.S. Department of Energy's Fermi National Accelerator Laboratory show fundamental particles called muons behaving in a way that is not predicted by scientists' best theory, the Standard Model of particle physics. This [landmark result](#), made with unprecedented precision, confirms a discrepancy that has been gnawing at researchers for decades.

The strong evidence that muons deviate from the Standard Model calculation might hint at exciting new physics. Muons act as a window into the subatomic world and could be interacting with yet undiscovered particles or forces.

"Today is an extraordinary day, long awaited not only by us but by the whole international physics community," said Graziano Venanzoni, co-spokesperson of the Muon g-2 experiment and physicist at the Italian National Institute for Nuclear Physics. "A large amount of credit goes to our young researchers who, with their talent, ideas and enthusiasm, have allowed us to achieve this incredible result."


- $\Delta a_\mu = (2.51 \pm 0.59) \times 10^{-9}$
- The Brookhaven + Fermilab results
 - **4.2 σ tension to the SM**

- The big news from the muons
- The muonic window to New Physics



The status of electron g-2

Parker et al., Science 360, 191–195 (2018)




$$2018 \text{ Cs : } \Delta a_e = a_e^{\text{exp}} - a_e^{\text{th}} = (-88 \pm 36) \times 10^{-14}$$

- **Negative value and a (- 2.4 σ) discrepancy**

- The two experiments are in tension at $\sim 4\sigma$, waiting for future experiments to see if there is something interesting in electron sector.

Morel et al., Nature 588, 61–65 (2020)



$$2020 \text{ Rb : } \Delta a_e = a_e^{\text{exp}} - a_e^{\text{th}} = (48 \pm 36) \times 10^{-14}$$

- Positive value and a (+ 1.6 σ) discrepancy
- **A 2.4 σ discrepancy with its own result in 2011!**


The new physics models for muon $g-2$

- The heavy solutions: SUSY, leptoquark, vector-like heavy leptons etc...
- The light solutions:
 - Mostly bosonic neutral mediators
 - Charged mediators should be quite heavy due to strong constraints from low energy experiments
 - Vectors: dark photon solution
 - Scalars: dark Higgs, axion-like particles, ...

Light dark photon and muon g-2

- Flavor universal : kinetic mixing dark photon

Bauer et al: 1803.05466 (JHEP)



$$\mathcal{L} \supset \epsilon F'_{\mu\nu} B^{\mu\nu}$$

$$\Rightarrow \mathcal{L} \supset \epsilon e A'_\mu J_{em}^\mu$$

$\text{Pi}0 \rightarrow A' + \text{gamma}, A' \rightarrow e^+ e^-$


$e^+ e^- \rightarrow \text{gamma } A'$

Visible dark photon can not explain muon g-2!

Light dark photon and muon g-2


- Flavor universal : kinetic mixing dark photon
- Further experimental updates

Visible A' : $A' \rightarrow \ell^+ \ell^-$



LHCb: 1910.06926 (PRL)

Invisible A' : $A' \rightarrow$ Missing Energy




BESIII: 1907.07046 (PRD)

Light dark photon and muon g-2

- Flavor universal : kinetic mixing dark photon
- Recent developments:
 - Incorporating Inelastic Dark Matter
 - $A' \rightarrow \chi_1 \chi_2, \chi_2 \rightarrow \chi_1 + f\bar{f}$
 - A' decays to MET + soft objects
 - Muon g-2 and dark matter simultaneously satisfied for large mass splitting

$$\Delta \equiv \frac{m_{\chi_2} - m_{\chi_1}}{m_{\chi_1}} = 0.4$$





(a) iDM: $\Delta = 0.4, \alpha_D = 0.1$. With muon $g - 2$ and DM regimes.

Tsai et al: 1908.07525 (PRL)

Light dark photon and muon g-2

- Flavor specific : $U(1)_{L_\mu - L_\tau}$ dark photon
- $m_{A'} \in [10, 100]$ MeV still viable for muon g-2


$$\mathcal{L} = g' A'_\alpha \times \left(\bar{L}_\mu \gamma^\alpha L_\mu - \bar{L}_\tau \gamma^\alpha L_\tau + \bar{\ell}_\mu \gamma^\alpha \ell_\mu - \bar{\ell}_\tau \gamma^\alpha \ell_\tau \right)$$



Light dark photon and muon g-2

- Flavor specific : $U(1)_{L_\mu-L_\tau}$ dark photon
- People are still proposing new proposals: e.g. MUonE

$$\mathcal{L} = g' A'_\alpha \times \left(\bar{L}_\mu \gamma^\alpha L_\mu - \bar{L}_\tau \gamma^\alpha L_\tau + \bar{\ell}_\mu \gamma^\alpha \ell_\mu - \bar{\ell}_\tau \gamma^\alpha \ell_\tau \right)$$



- Similar to NA64 μ : 10^{12} muons on target from CERN SPS upgraded muon beam
- MUonE: 150 GeV muon + e at rest
- $\mu e \rightarrow \mu e$ measuring hadronic vacuum polarization (HVP) contribution for g-2
- $\mu e \rightarrow \mu e Z' \rightarrow \mu e + \text{MET}$

Light dark scalar and muon g-2


- Effective Lagrangian:

$$\mathcal{L}_{\text{eff}} \supset \sum_q \epsilon_q \frac{m_q}{v} \phi \bar{q} q + \sum_\ell \epsilon_\ell \frac{m_\ell}{v} \phi \bar{\ell} \ell + \epsilon_W \frac{2m_W^2}{v} \phi W_\mu^+ W^{\mu-}$$

- **Universal:** $\epsilon \equiv \epsilon_q = \epsilon_W = \epsilon_\ell$ (Higgs portal + singlet)
- **Lepton specific:** $\epsilon_q \approx \epsilon_W \neq \epsilon_\ell$ (from type-X 2HDM + singlet)
- **Muonic specific:** $\epsilon_\mu \neq 0$, others = 0

- **Muon g-2:**

$$\Delta a_\mu = \frac{m_\mu^2}{8\pi^2 v^2} \epsilon_\ell^2 \int_0^1 dx \frac{(1-x)^2(1+x)}{(1-x)^2 + x(m_\phi/m_\mu)^2}$$



JL, Carlos E.M. Wagner, Xiao-ping Wang: 1810.11028 [JHEP]


Scalars are less constrained comparing with A' , due to smaller coupling to e

Light dark scalar and muon g-2

- Effective Lagrangian:

$$\mathcal{L}_{\text{eff}} \supset \sum_q \epsilon_q \frac{m_q}{v} \phi \bar{q}q + \sum_\ell \epsilon_\ell \frac{m_\ell}{v} \phi \bar{\ell}\ell + \epsilon_W \frac{2m_W^2}{v} \phi W_\mu^+ W^{\mu-}$$

- **Universal:** $\epsilon \equiv \epsilon_q = \epsilon_W = \epsilon_\ell$ (Higgs portal + singlet)
- **Lepton specific:** $\epsilon_q \approx \epsilon_W \neq \epsilon_\ell$ (from type-X 2HDM + singlet)
- **Muonic specific:** $\epsilon_\mu \neq 0$, others = 0
- **New experiment updates for leptonphilic/universal dark scalar**



Utilizes the large tau coupling: $e^+e^- \rightarrow \tau^+\tau^-\phi$

Outline

- The light dark sector and $(g - 2)_{e/\mu}$
- The cosmological triangle
- The ν scalar in the early universe and $(g - 2)_{\mu}$
- Summary


The cosmological triangle

- Referring to Axion-Like Particle searches with photon couplings, works for scalar as well

$$\mathcal{L}_{\text{ALP}} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m_\phi^2 \phi^2 - \frac{g_{\phi\gamma}}{4} \phi F_{\mu\nu} \tilde{F}^{\mu\nu}$$

$$\tau_{\phi\gamma} = \Gamma_{\phi\gamma}^{-1} = \frac{64\pi}{m_\phi^3 g_{\phi\gamma}^2}$$

- Blue solid line from BBN constraints: Y_p , D/H
- Blue dot dashed line: allow varying neutrino chemical potential and ΔN_{eff}
- Blue dashed line: N_{eff} constraints from CMB measurements



Depta et al: 2002.08370 (JCAP)

The cosmological triangle

- Referring to Axion-Like Particle searches with photon couplings, works for scalar as well

$$\mathcal{L}_{\text{ALP}} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m_\phi^2 \phi^2 - \frac{g_{\phi\gamma}}{4} \phi F_{\mu\nu} \tilde{F}^{\mu\nu}$$


$$\tau_{\phi\gamma} = \Gamma_{\phi\gamma}^{-1} = \frac{64\pi}{m_\phi^3 g_{\phi\gamma}^2}$$

- N_{eff} from CMB constraints constraints:

- SM value: $\left(T_\nu^0 / T_\gamma^0 \right) = \left(\frac{4}{11} \right)^{1/3}$ due to $e^+ e^-$ annihilation

- $\phi \rightarrow \gamma\gamma$ injects entropy to photons after neutrino decoupling ($T_D^0 \sim 2.3$ MeV)

- Further lowering the ratio T_ν / T_γ



Depta et al: 2002.08370 (JCAP)


The cosmological triangle

- Referring to Axion-Like Particle searches with photon couplings, works for scalar as well

$$\mathcal{L}_{\text{ALP}} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m_\phi^2 \phi^2 - \frac{g_{\phi\gamma}}{4} \phi F_{\mu\nu} \tilde{F}^{\mu\nu}$$

$$\tau_{\phi\gamma} = \Gamma_{\phi\gamma}^{-1} = \frac{64\pi}{m_\phi^3 g_{\phi\gamma}^2}$$

- If one allows extra cosmological setup: ΔN_{eff} (e.g. dark radiation), to compensate the low T_ν
- A triangle area is still allowed for MeV ALP, similar for dark scalar




Depta et al: 2002.08370 (JCAP)

Outline

- The light dark sector and $(g - 2)_{e/\mu}$
- The cosmological triangle
- The ν scalar in the early universe and $(g - 2)_{\mu}$
- Summary

The ν scalar in the early universe and $(g - 2)_\mu$

- Our motivation:
 - Light dark scalar ($\lesssim 30$ MeV) with muon coupling same as SM Higgs ($\sim m_\mu/v_h$) can solve $(g - 2)_\mu$



JL, Carlos E.M. Wagner, Xiao-ping Wang: 1810.11028 [JHEP]


The ν scalar in the early universe and $(g - 2)_\mu$

- Our motivation:

- Light dark scalar ($\lesssim 30$ MeV) with muon coupling same as SM Higgs ($\sim m_\mu/v_h$) can solve $(g - 2)_\mu$
- Such coupling naturally induce photon couplings at 1-loop

- $\mathcal{L}_{\text{eff}} \supset -\frac{g_{\gamma\gamma}}{4}\phi F_{\mu\nu}F^{\mu\nu}$

- Right in the cosmological triangle
- We are not satisfactory with adding hand-waiving ΔN_{eff}
- We do it dynamically by coupling scalar to ν (neutrinos) and solve the ν mass problem at the same time



JL, Navin McGinnis, Carlos E.M. Wagner, Xiao-ping Wang
2110.14665

The model setup

- Low energy model: $\mathcal{L}_{\text{eff}} \supset -g_\mu \phi \bar{\mu} \mu - \left(\frac{g_{\nu_a}}{2} \phi \nu_a \cdot \nu_a + h.c. \right)$
- 1-loop induced photon coupling: $\mathcal{L}_{\text{eff}} \supset -\frac{g_{\gamma\gamma}}{4} \phi F_{\mu\nu} F^{\mu\nu}$
- $(g-2)_\mu$ and cosmological triangle fixes:
 - $m_\phi \sim 1 \text{ MeV}$, $g_\mu \sim m_\mu/v_h$ and $g_\gamma \sim -\frac{2\alpha g_\mu}{3\pi m_\mu}$
 - Only neutrino coupling g_ν is free parameter
 - Problem: $\phi \rightarrow \gamma\gamma$ injects entropy to photon plasma, leads to lower T_ν



The solution: delayed neutrino decoupling

- Solution: ϕ coupling to ν , delayed neutrino decoupling T_D
 - It forces $T_\nu = T_\gamma$ ($T_\gamma < T_D$), therefore entropy injection from e^+e^- shares in ν/γ sectors
 - Therefore, it effectively raises T_ν and compensate the $\phi \rightarrow \gamma\gamma$ entropy injection
 - The new decoupling T_D is determined by s-channel resonant interaction $\nu\nu \leftrightarrow \phi^* \leftrightarrow \gamma\gamma$

$$\frac{dn_\nu}{dt} + 3Hn_\nu = \langle \sigma v \rangle_{\text{res}} (n_{\nu,\text{eq}}^2 - n_\nu^2)$$

$$R \equiv n_\nu^{\text{eq}} \langle \sigma v \rangle \approx \frac{8\sqrt{2\pi}}{3\xi(3)} \Gamma_\phi \text{BR}_{\gamma\gamma} \text{BR}_{\nu\nu} x^{-3/2} e^{-x} \quad \text{V.S. Hubble rate}$$

The delayed neutrino decoupling



- Solution: ϕ coupling to ν , delayed neutrino decoupling T_D
 - A small $BR(\phi \rightarrow \nu\nu)$ solve the cosmological triangle problem
 - We check if it also solves the neutrino mass problem

UV model

Field	$SU(2)_L$	$U(1)_Y$	Z_2
μ_R	1	-1	-1
ϕ	1	0	-1
H	2	$\frac{1}{2}$	+1
H'	2	$\frac{1}{2}$	-1
N	1	0	+1
N'	1	0	-1

$$\begin{aligned}
 \mathcal{L}_{UV} = & y_\mu \bar{L}_\mu H' \mu_R + y_{N,i} (L_i \cdot H) N + y'_{N,j} (L_j \cdot H') N' \\
 & + \lambda_N N \cdot N' \phi + \mu_\phi H'^\dagger H \phi \\
 & + \frac{1}{2} m_N N \cdot N + \frac{1}{2} m_{N'} N' \cdot N' + h.c.
 \end{aligned} \tag{25}$$

- Muon specific coupling $g_\mu = y_\mu \frac{v \mu_\phi}{\sqrt{2} m_{H'}}$
- Two sterile neutrino setup:
 - One active neutrino is massless
 - Two massive active neutrino can be obtained
 - ϕ couples to heaviest active neutrino dominantly
 - Fits to $\text{BR}(\phi \rightarrow \nu\nu) \sim \mathcal{O}(0.1\%)$

$$m_1 = 0, \quad m_2 \simeq 8.7 \times 10^{-3} \text{eV}, \quad m_3 \simeq 0.059 \text{eV}$$

$$g_{\nu_3} \simeq 2.5 \times 10^{-10}$$

Summary

- Light dark sector can solve $(g - 2)_\mu$ problem, but is under severe constraints
- Dark scalar solution is less constrained than dark photon
- However, needs extra ΔN_{eff} to save the cosmological triangle
- We build a dynamic model to save the cosmological triangle
 - Solve $(g - 2)_\mu$ and neutrino mass simultaneously

Backup slides

Some twist for e/muon g-2 combined

UV model details

Field	$SU(2)_L$	$U(1)_Y$	Z_2
μ_R	1	-1	-1
ϕ	1	0	-1
H	2	$\frac{1}{2}$	+1
H'	2	$\frac{1}{2}$	-1
N	1	0	+1
N'	1	0	-1

$$\begin{aligned}
 \mathcal{L}_{UV} = & y_\mu \bar{L}_\mu H' \mu_R + y_{N,i} (L_i \cdot H) N + y'_{N,j} (L_j \cdot H') N' \\
 & + \lambda_N N \cdot N' \phi + \mu_\phi H'^\dagger H \phi \\
 & + \frac{1}{2} m_N N \cdot N + \frac{1}{2} m_{N'} N' \cdot N' + h.c.
 \end{aligned} \tag{25}$$

$$g_\mu = y_\mu \frac{v \mu_\phi}{\sqrt{2} m_{H'}^2}$$

$$\begin{pmatrix} \vec{\nu} \\ N \\ N' \end{pmatrix} \approx \begin{pmatrix} \mathbb{I}_{3 \times 3} & \frac{\vec{y}_N v + \vec{y}_{N'} v'}{\sqrt{2} \lambda_N v_\phi} & \frac{\vec{y}_N v - \vec{y}_{N'} v'}{\sqrt{2} \lambda_N v_\phi} \\ -\frac{\vec{y}_{N'}^T v'}{\lambda_N v_\phi} & \frac{1}{\sqrt{2}} + z & -\frac{1}{\sqrt{2}} + z \\ -\frac{\vec{y}_N^T v}{\lambda_N v_\phi} & \frac{1}{\sqrt{2}} + z & \frac{1}{\sqrt{2}} + z \end{pmatrix} \begin{pmatrix} \vec{\nu}' \\ \tilde{N} \\ \tilde{N}' \end{pmatrix}$$

$$M_{ij} = (y_{N,i} y_{N',j} + y_{N,j} y_{N',i}) \frac{\sqrt{2} v v'}{2 \lambda_N v_\phi},$$

$$\tilde{m}_{\nu_1} = 0,$$

$$\tilde{m}_{\nu_2} = \frac{v v'}{\sqrt{2} \lambda_N v_\phi} (|\vec{y}_N| |\vec{y}_{N'}| - \vec{y}_N \cdot \vec{y}_{N'}) \sim 0.1 \text{eV},$$

$$\tilde{m}_{\nu_3} = \frac{v v'}{\sqrt{2} \lambda_N v_\phi} (|\vec{y}_N| |\vec{y}_{N'}| + \vec{y}_N \cdot \vec{y}_{N'}) \sim 0.1 \text{eV},$$

$$\tilde{m}_N \approx \frac{\lambda_N v_\phi}{\sqrt{2}} + \frac{m_N + m_{N'}}{2} + \mathcal{O}\left(\frac{1}{\lambda_N v_\phi}\right),$$

$$\tilde{m}_{N'} \approx \frac{\lambda_N v_\phi}{\sqrt{2}} - \frac{m_N + m_{N'}}{2} \mathcal{O}\left(\frac{1}{\lambda_N v_\phi}\right),$$

$$g_{\nu_a} = \frac{\tilde{m}_{\nu_a}}{v_\phi}.$$

$$m_1 = 0, \quad m_2 \simeq 8.7 \times 10^{-3} \text{eV}, \quad m_3 \simeq 0.059 \text{eV}$$

$$g_{\nu_3} \simeq 2.5 \times 10^{-10}$$