Precision Calculations for Heavy Quark Decays

Yu-Ming Wang

Nankai University

Workshop on Hadron Structure at High-Energy, High-Luminosity Facilities October 26, 2021, Nanjing

Yu-Ming Wang (Nankai)

QCD and Heavy Quark Decays

October 26, 2021 1/30

Why precision calculations?

- Understanding the general properties of power expansion in EFTs (HQET, SCET, NRQCD).
- Interesting to understand the strong interaction dynamics of heavy quark decays.
 - Factorization properties of the subleading-power amplitudes.
 - Renormalization and asymptotic properties of the higher-twist B-meson DAs.
 - Interplay of different QCD techniques.
- Precision determinations of the CKM matrix elements $|V_{ub}|$ and $|V_{cb}|$. Power corrections, QED corrections, BSM physics.
- Crucial to understand the CP violation in *B*-meson decays. Strong phase of $\mathscr{A}(B \to M_1 M_2) @ m_b$ scale in the leading power.
- Indispensable for understanding the flavour puzzles (continuously updated).
 - P'_5 and $R_{K^{(*)}}$ anomalies in $B \to K^{(*)} \ell^+ \ell^-$.
 - $R_{D^{(*)}}$ anomalies in $B \to D^{(*)} \ell \bar{\nu}_{\ell}$.
 - Color suppressed hadronic *B*-meson decays.
 - ▶ Polarization fractions of penguin dominated $B_{(s)} \rightarrow VV$ decays.

Theory tools for precision flavor physics

New Physics:
$$\mathscr{L}_{NP}$$

 \downarrow
EW scale (m_W) : $\mathscr{L}_{SM} + \mathscr{L}_{D>4}$
 \downarrow
Heavy-quark scale (m_b) : $\mathscr{L}_{eff} = -\frac{G_F}{\sqrt{2}} \sum_i C_i Q_i + \mathscr{L}_{eff,D>6}$
 \downarrow
QCD scale (Λ_{QCD})

• Aim: $\langle f | Q_i | \bar{B} \rangle = ?$

- QCD factorization [Diagrammatic approach].
- SCET factorization [Operator formalism].
- TMD factorization.
- (Light-cone) QCD sum rules.
- Lattice QCD.

• Key concepts: Factorization, Resummation, Evolution.

Yu-Ming Wang (Nankai)

• The light-ray HQET matrix element [Grozin, Neubert, 1997]:

$$\langle 0|\bar{q}_{\beta}(z)[z,0]h_{\nu\alpha}(0)|\bar{B}(\nu)\rangle = -\frac{i\tilde{f}_{B}m_{B}}{4} \left[\frac{1+\psi}{2}\left\{2\,\tilde{\phi}_{B}^{+}(t,\mu) + \frac{\tilde{\phi}_{B}^{-}(t,\mu) - \tilde{\phi}_{B}^{+}(t,\mu)}{t}\,\not\xi\right\}\gamma_{5}\right]_{\alpha\beta}$$

Important EOM constraints in HQET [Kawamura, Kodaira, Qiao, Tanaka, 2001, 2003].

• Evolution equation at one loop [Lange, Neubert, 2003]:

$$\frac{d\phi_B^+(\omega,\mu)}{d\ln\mu} = -\left[\Gamma_{\rm cusp}(\alpha_s)\ln\frac{\mu}{\omega} + \gamma_+(\alpha_s)\right]\phi_B^+(\omega,\mu) - \omega\int_0^\infty d\eta\,\Gamma_+(\omega,\eta,\alpha_s)\,\phi_B^+(\eta,\mu)\,.$$

This is an integro-differential equation!

• (Relatively) complicated solution [Lee, Neubert, 2005]:

$$\begin{split} \phi^+_B(\omega,\mu) &= e^{V-2\gamma_{Eg}} \frac{\Gamma(2-g)}{\Gamma(g)} \int_0^\infty \frac{d\eta}{\eta} \phi^+_B(\eta,\mu_0) \left(\frac{\max(\omega,\eta)}{\mu_0}\right)^g \\ &\times \frac{\min(\omega,\eta)}{\max(\omega,\eta)} {}_2F_1 \left(1-g,2-g,2,\frac{\min(\omega,\eta)}{\max(\omega,\eta)}\right), \\ g(\mu,\mu_0) &= \int_{\alpha_s(\mu_0)}^{\alpha_s(\mu)} d\alpha \frac{\Gamma_{\mathrm{cusp}}(\alpha)}{\beta(\alpha)} \approx \frac{2C_F}{\beta_0} \ln \frac{\alpha_s(\mu_0)}{\alpha_s(\mu)}. \end{split}$$

Making the QCD resummation for enhanced logarithms complicated!

Yu-Ming Wang (Nankai)

• Better understanding from the RGE in coordinate space [Braun, Ivanov, Korchemsky, 2004]:

$$\frac{d\tilde{\phi}_B^+(t,\mu)}{d\ln\mu} = -\Gamma_{\rm cusp}(\alpha_s) \left\{ \left[\ln(i\,\tilde{\mu}\,t) - \frac{1}{4} \right] \tilde{\phi}_B^+(t,\mu) + \int_0^1 du \,\frac{\bar{u}}{u} \left[\tilde{\phi}_B^+(t,\mu) - \tilde{\phi}_B^+(\bar{u}\,t,\mu) \right] \right\}.$$

- Absence of the local operator product expansion due to the non-analytical term!
- $\tilde{\phi}_B^+(t,\mu)$ only mixes into $\tilde{\phi}_B^+(\bar{u}t,\mu)$ with $0 \le \bar{u} \le 1$ under renormalization.

• Renormalization of $[\bar{q}_s(t\bar{n})\Gamma b_v(0)]$ does not commute with the short-distance expansion.

$$[(\bar{q}_{s}Y_{s})(t\bar{n})\vec{\eta}\,\Gamma(Y_{s}^{\dagger}b_{v})(0)]_{R}\neq\sum_{p=0}\frac{t^{p}}{p!}\left[\bar{q}_{s}(0)\,(n\cdot\overleftarrow{D})^{p}\,\vec{\eta}\,\Gamma b_{v}(0)\right]_{R}.$$

- Many other examples in QCD physics (e.g., Light-cone projection and renormalization)!
- Non-negative moments of the B-meson distribution amplitude ill defined.
- ▶ Non-trivial generalization of the QCD equations of motion beyond the tree level.

Yu-Ming Wang (Nankai)

• Fourier/Mellin transformation:

$$\varphi_B^+(\theta,\mu) = \int_0^\infty \frac{d\omega}{\omega} \left(\frac{\omega}{\mu}\right)^{-i\theta} \phi_B^+(\omega,\mu) \quad \Leftrightarrow \quad \phi_B^+(\omega,\mu) = \int_{-\infty}^\infty \frac{d\theta}{2\pi} \left(\frac{\omega}{\mu}\right)^{i\theta} \varphi_B^+(\theta,\mu).$$

• Solution to the RGE in Mellin space:

$$\varphi_B^+(\theta,\mu) = e^{V-2\gamma_{ES}} \left(\frac{\mu}{\mu_0}\right)^{i\theta} \frac{\Gamma(1-i\theta)}{\Gamma(1+i\theta)} \cdot \frac{\Gamma(1+i(\theta+ig))}{\Gamma(1-i(\theta+ig))} \varphi_B^+(\theta+ig,\mu_0).$$

Already very symmetric solution in Mellin space.

• Yet simper solution to the integral-differential equation exists?

⇔ Eigenfunctions of the Lange-Neubert kernel [Bell, Feldmann, YMW and Yip, 2013].

$$\begin{split} \phi_B^+(\theta,\mu) &:= \frac{\Gamma(1-i\theta)}{\Gamma(1+i\theta)} \bar{\rho}_B^+(\theta,\mu) \\ &= \frac{\Gamma(1-i\theta)}{\Gamma(1+i\theta)} \int_0^\infty \frac{d\omega'}{\omega'} \rho_B^+(\omega',\mu) \left(\frac{\omega'}{\mu}\right)^{-i\theta}. \end{split}$$

Yu-Ming Wang (Nankai)

• Linear differential equation [Bell, Feldmann, YMW and Yip, 2013]:

$$\frac{d\rho_B^+(\omega',\mu)}{d\ln\mu} = -\left[\Gamma_{\rm cusp}(\alpha_s)\ln\frac{\mu}{\dot{\omega}'} + \gamma_+(\alpha_s)\right]\rho_B^+(\omega',\mu)\,.$$

Local evolution in the dual space!

• Integral transformation:

$$\begin{split} \phi^+_B(\omega,\mu) &= \int_0^\infty \frac{d\omega'}{\omega'} \sqrt{\frac{\omega}{\omega'}} J_1\left(2\sqrt{\frac{\omega}{\omega'}}\right) \rho^+_B(\omega',\mu), \\ \rho^+_B(\omega',\mu) &= \int_0^\infty \frac{d\omega}{\omega} \sqrt{\frac{\omega}{\omega'}} J_1\left(2\sqrt{\frac{\omega}{\omega'}}\right) \phi^+_B(\omega,\mu). \end{split}$$

Eigenfunction of the Lange-Neubert kernel at one-loop is the Bessel function!

• Interesting transformation from the coordinate space to the dual space:

$$\rho_B^+(\omega',\mu) = \int \frac{dt}{2\pi} \left(1 - \exp\left[-\frac{i}{\omega' t}\right] \right) \tilde{\phi}_B^+(t,\mu) \,.$$

 $\rho_B^+(\omega',\mu)$ cannot be constructed from the local OPE of $\tilde{\phi}_B^+(t,\mu)$.

Yu-Ming Wang (Nankai)

• Solution to the RGE in dual space [Bell, Feldmann, Wang, Yip, 2013]:

$$ho_B^+(\omega',\mu) = e^V \left(rac{\mu_0}{\omega'}
ight)^{-g}
ho_B^+(\omega',\mu_0).$$

Very compact expression in a full analytical form!

• Solution to the RGE in momentum space:

$$\phi_B^+(\omega,\mu) = e^V \int_0^\infty \frac{d\omega'}{\omega'} \sqrt{\frac{\omega}{\omega'}} J_1\left(2\sqrt{\frac{\omega}{\omega'}}\right) \left(\frac{\mu_0}{\hat{\omega}'}\right)^{-g} \rho_B^+(\omega',\mu_0) d\omega'$$

Still a beautiful expression!

• The logarithmic inverse moments of LCDA and spectral function:

$$\int_0^\infty \frac{d\omega}{\omega} \ln^n\left(\frac{\omega}{\mu}\right) \phi_B^+(\omega,\mu) \stackrel{n=0,1,2}{=} \int_0^\infty \frac{d\omega'}{\omega'} \ln^n\left(\frac{\hat{\omega}'}{\mu}\right) \rho_B^+(\omega',\mu) \equiv L_n(\mu).$$

Non-trivial mixing of $L_n(\mu)$ in dual space under renormalization.

$$\frac{dL_n(\mu)}{d\ln\mu} = \Gamma_{\rm cusp}(\alpha_s) L_{n+1}(\mu) - \gamma_+(\alpha_s) L_n(\mu) - nL_{n-1}(\mu).$$

More complicated RGE for the logarithmic inverse moments in momentum space.

Yu-Ming Wang (Nankai)

• Collinear conformal symmetry for the Lange-Neubert kernel [Braun, Manashov, 2014]:

$$\left(\frac{d}{d\ln\mu}+\mathscr{H}_{\rm LN}\right)O_+(z,\mu)=0.$$

HLN is almost determined by the commutation relations completely [Knoedlseder, Offen, 2011]

$$[S_+, \mathscr{H}_{\mathrm{LN}}] = 0, \qquad [S_0, \mathscr{H}_{\mathrm{LN}}] = 1$$

The beautiful solution in terms of S_+ :

$$\mathscr{H}_{\rm LN} = \ln(i\,\mu\,S^+) - \psi(1) - \frac{5}{4}\,.$$

• Generators of the collinear conformal group:

$$S_+ = z^2 \partial_z + 2jz$$
, $S_0 = z \partial_z + j$, $S_- = -\partial_z$.

Eigenfunctions of S_+ [Braun, Manashov, 2014]:

$$iS_+ Q_s(z) = sQ_s(z), \qquad Q_s(z) = -\frac{1}{z^2} e^{is/z}$$
$$\langle e^{-i\omega z} | Q_s(z) \rangle = \frac{1}{\sqrt{\omega s}} J_1(2\sqrt{\omega s}).$$

Wide applications of the conformal symmetry in high energy physics!

Yu-Ming Wang (Nankai)

• RG evolution of $\phi_B^+(\omega,\mu)$ at two loops [Braun, Ji, Manashov, 2019; Liu, Neubert, 2020]:

$$\begin{aligned} \frac{d\phi_B^+(\omega,\mu)}{d\ln\mu} &= \left[\Gamma_{\rm cusp}(\alpha_s)\ln\frac{\omega}{\mu} - \gamma_\eta(\alpha_s)\right]\phi_B^+(\omega,\mu) + \Gamma_{\rm cusp}(\alpha_s)\int_0^\infty dx\Gamma(1,x)\phi_B^+(\omega/x,\mu) \\ &+ \left(\frac{\alpha_s}{2\pi}\right)^2 C_F \int_0^1 \frac{dx}{1-x}h(x)\phi_B^+(\omega/x,\mu)\,. \end{aligned}$$

The last missing element for the NLL predictions of exclusive B-meson decay observables!

- The two-loop eigenfunctions depend on the strong coupling α_s [Braun, Ji, Manashov, 2019].
- Applying the Laplace transformation of the LCDA [Galda, Neubert, 2020]

$$\tilde{\phi}^+_B(\eta,\mu) = \int_0^\infty \frac{d\omega}{\omega} \, \phi^+_B(\omega,\mu) \, \left(\frac{\omega}{\bar{\omega}}\right)^{-\eta} \, ,$$

 \Rightarrow the general solution to the two-loop RGE of $\phi_B^+(\omega,\mu)$

$$\begin{split} \tilde{\phi}_{B}^{+}(\eta,\mu) &= \exp\left[S(\mu_{0},\mu) + a_{\gamma}(\mu_{0},\mu) + 2\gamma_{E}a_{\Gamma}(\mu_{0},\mu)\right] \left(\frac{\bar{\omega}}{\mu_{0}}\right)^{-a_{\Gamma}(\mu_{0},\mu)} \\ &\times \frac{\Gamma(1+\eta+a_{\Gamma}(\mu_{0},\mu))\Gamma(1-\eta)}{\Gamma(1-\eta-a_{\Gamma}(\mu_{0},\mu))\Gamma(1+\eta)} \exp\left[\int_{\alpha_{s}(\mu_{0})}^{\alpha_{s}(\mu)} \frac{d\alpha}{\beta(\alpha)}\mathscr{G}(\eta+a_{\Gamma}(\mu_{\alpha},\mu),\alpha)\right] \\ &\times \tilde{\phi}_{B}^{+}(\eta+a_{\Gamma}(\mu_{0},\mu),\mu_{0}). \end{split}$$

Yu-Ming Wang (Nankai)

Double Radiative B_q -Meson Decays

• Leading-order contributions at $\mathcal{O}(\alpha_s^0)$:

$$b \longrightarrow \gamma(p) \qquad b \longrightarrow \gamma(p) \qquad b \longrightarrow \gamma(p) \qquad b \longrightarrow \gamma(p) \qquad c \rightarrow \gamma(p) \qquad c \rightarrow \gamma(p) \qquad c \rightarrow \gamma(q) \qquad c \rightarrow \gamma(q)$$

Kinematics:

$$p_{\mu} = \frac{n \cdot p}{2} \, \bar{n}_{\mu} \equiv \frac{m_{B_q}}{2} \, \bar{n}_{\mu} \,, \qquad q_{\mu} = \frac{\bar{n} \cdot q}{2} \, n_{\mu} \equiv \frac{m_{B_q}}{2} \, n_{\mu} \,.$$

Interplay of the soft and collinear QCD dynamics!

Decay amplitude:

$$\bar{\mathscr{A}}(\bar{B}_q \to \gamma \gamma) = -\frac{4G_F}{\sqrt{2}} \,\frac{\alpha_{\rm em}}{4\pi} \,\varepsilon_1^{*\alpha}(p) \,\varepsilon_2^{*\beta}(q) \sum_{p=u,c} V_{pb} V_{pq}^* \sum_{i=1}^8 C_i T_{i,\alpha\beta}^{(p)} \,.$$

Hadronic tensors:

$$\begin{split} T_{7,\alpha\beta} &= 2\,\overline{m}_b(\mathbf{v})\int d^4x\,e^{iq\cdot x}\,\langle 0|\mathrm{T}\left\{j^{\mathrm{em}}_{\beta}(x),\bar{q}_L(0)\sigma_{\mu\alpha}\,p^{\mu}b_R(0)\right\}|\bar{B}_q(p+q)\rangle \\ &+\left[p\leftrightarrow q,\alpha\leftrightarrow\beta\right],\\ T^{(p)}_{i,\alpha\beta} &= -(4\,\pi)^2\int d^4x\int d^4y\,e^{ip\cdot x}\,e^{iq\cdot y}\,\langle 0|\mathrm{T}\left\{j^{\mathrm{em}}_{\alpha}(x),j^{\mathrm{em}}_{\beta}(y),P^{(p)}_i(0)\right\}|\bar{B}_q(p+q)\rangle,\\ &(i=1,\ldots,6,8). \end{split}$$

Main task: Construct the SCET factorization formulae beyond the leading power.

Yu-Ming Wang (Nankai)

General aspects of $B_q \rightarrow \gamma \gamma$

• Parametrization:

$$T^{(p)}_{i,\alpha\beta} = i m^3_{B_q} \left[\left(g^{\perp}_{\alpha\beta} - i \varepsilon^{\perp}_{\alpha\beta} \right) F^{(p)}_{i,L} - \left(g^{\perp}_{\alpha\beta} + i \varepsilon^{\perp}_{\alpha\beta} \right) F^{(p)}_{i,R} \right] \,.$$

- Only two helicity form factors due to the Ward identities and the transversity conditions.
- Similar decomposition for the radiative leptonic *B*-meson decay amplitude.
- Hierarchy structure due to the chiral weak interactions and helicity conservation:

$$F_{i,L}^{(p)}:F_{i,R}^{(p)}=1:\left(rac{\Lambda_{ ext{QCD}}}{m_b}
ight).$$

In analogy to the two-body nonleptonic $B \rightarrow VV$ decays [Beneke, Rohrer, Yang, 2007].

• Transversity form factors:

$$F_{i,\parallel}^{(p)} = F_{i,L}^{(p)} - F_{i,R}^{(p)} \,, \qquad F_{i,\perp}^{(p)} = F_{i,L}^{(p)} + F_{i,R}^{(p)} \,.$$

The two-photon final states as the CP eigenstates with the eigenvalues +1 and -1.

Yu-Ming Wang (Nankai)

Current status of $B_q \rightarrow \gamma \gamma$

- QCD factorization at leading power in Λ/m_b and at NLO in α_s [Descotes-Genon, Sachrajda, 2003].
 - No collinear strong interaction dynamics at LP.
 - ► The two-loop $b \rightarrow q \gamma$ matrix elements of QCD penguin operators NOT included.
 - \Rightarrow A complete factorization-scale independence at NLO is absent!
- Subleading power contributions from the weak annihilation diagrams [Bosch, Buchalla, 2002].
 - Complex perturbative hard functions evaluated at one loop.
 - Diagrammatic factorization established at two loops.
- The new (technical)-ingredients from [Shen, YMW, Wei, 2020]:
 - A complete NLL calculation for the LP contribution \Rightarrow 2-loop evolution of ϕ_B^+ .
 - The NLP factorization for the energetic photon radiation off the light quark. The so-called "soft form factor" defined in [Beneke, Rohrwild, 2011] is factorizable!
 - The NLP factorization for the light-quark mass effect.
 - The NLP factorization for the SCET current $J^{(A2)} \supset (\bar{\xi}_{\overline{hc}} W_{\overline{hc}}) \gamma_{\alpha}^{\perp} P_L \left(\frac{i \vec{p}_{\top}}{2m_h} \right) h_{\nu}.$
 - ▶ The NLP factorization for the subleading twist *B*-meson LCDAs.
 - The resolved photon contribution with the dispersion technique.
- Referee Report: "The authors have endeavored to study the decay $B_{d,s} \rightarrow \gamma \gamma$ quite comprehensively, and achieved much progress far beyond the seminal paper by Descotes-Genon and Sachrajda on that subject".

Yu-Ming Wang (Nankai)

SCET factorization at leading power

 $\ \ \ O CD \rightarrow SCET_I \ matching \ at \ LP:$

$$\sum_{i=1}^{8} C_{i} T_{i,\alpha\beta}^{(p)} = \sum_{i=1}^{8} C_{i} H_{i}^{(p)} \left\{ \int d^{4}x e^{iq \cdot x} \langle 0| T \left\{ f_{\beta,\text{SCET}_{I}}^{\text{em}}(x), \left[(\bar{\xi}_{\text{hc}} W_{\overline{\text{hc}}}) \gamma_{\alpha}^{\perp} P_{L} h_{\nu} \right] (0) \right\} | \bar{B}_{q} \rangle + [p \leftrightarrow q, \alpha \leftrightarrow \beta] \right\}.$$

Universal SCET_I correlation function independent of the weak-operator index!

• Perturbative matching coefficients at NLO:

$$\begin{split} \sum_{i=1}^8 C_i H_i^{(p)} &= (-2\,i)\,\overline{m}_b(\nu)\,m_{B_q}\,V_{7,\mathrm{eff}}^{(p)}\,,\\ V_{7,\mathrm{eff}}^{(p)} &= C_7^{\mathrm{eff}}\,C_{T_1}^{(\mathrm{A0})} + \sum_{i=1,\ldots,6,8} \frac{\alpha_s(\mu)}{4\pi}\,C_i^{\mathrm{eff}}\,F_{i,7}^{(p)}\,. \end{split}$$

- The hard function $C_{T_1}^{(A0)}$ from matching the heavy-light tensor current onto SCET_I.
- The hard functions $F_{i,7}^{(p)}$ (i = 1, ..6, 8) from perturbative matching of the $b \rightarrow q\gamma$ matrix elements [Buras et al, 2002].
- $V_{7,\text{eff}}^{(p)}$ depends on both the electro-weak scale and the heavy-quark mass scale.

Yu-Ming Wang (Nankai)

SCET factorization at leading power

• $SCET_I \rightarrow SCET_{II}$ matching in coordinate space:

$$\begin{aligned} \mathscr{T}_{\alpha\beta} &= \int d^4 x e^{iq \cdot x} \left\langle 0 | \mathbf{T} \left\{ j^{\text{em}}_{\beta,\text{SCET}_{\mathrm{I}}}(x), \left[\left(\bar{\xi}_{\overline{hc}} W_{\overline{hc}} \right) \gamma^{\perp}_{\alpha} P_L h_v \right] (0) \right\} | \bar{B}_q \right\rangle \\ &= \int dt \, \mathscr{I} \left(\frac{\bar{n} \cdot q}{\mu^2 v \cdot x} \right) \left\langle 0 | (\bar{q}_s Y_s)(tn) \gamma^{\perp}_{\beta} \not n \gamma^{\perp}_{\alpha} P_L (Y^{\dagger}_s h_v)(0) | \bar{B}_q \right\rangle. \end{aligned}$$

• Only need the one-loop jet function (2-loop result by Liu and Neubert, 2020).

$$J = 1 + \frac{\alpha_s(\mu) C_F}{4\pi} \left[\ln^2 \left(\frac{\mu^2}{m_b \omega} \right) - \frac{\pi^2}{6} - 1 \right] + \mathcal{O}(\alpha_s^2).$$

- ▶ The soft dynamics encoded in the twist-two HQET *B*-meson LCDA.
- The resulting LP factorization formula:

$$egin{aligned} egin{aligned} &oldsymbol{\mathscr{A}_{\mathrm{LP}}}(ar{B}_q o \gamma\gamma) = i \, rac{4\,G_F}{\sqrt{2}} \, rac{lpha_{\mathrm{em}}}{4\pi} \, arepsilon_1^{stlpha}(p) \, arepsilon_2^{steta}(q) \, \left[g_{lpha\beta}^{\perp} - iarepsilon_{lpha\beta}^{\perp}
ight] \, e_q f_{B_q} \, m_{B_q}^2 \, K^{-1}(m_b,\mu) \ &\left[\sum_{p=u,c} V_{pb} V_{pq}^{st} \, \overline{m}_b(\mathbf{v}) \, V_{7,\mathrm{eff}}^{(p)}(m_b,\mu,\mathbf{v})
ight] \, \int_0^\infty rac{d\omega}{\omega} \, \phi_B^+(\omega,\mu) \, J(m_b,\omega,\mu) \, . \end{aligned}$$

Factorization of the hard, hard-collinear and soft dynamics.

Yu-Ming Wang (Nankai)

SCET factorization at leading power

- No common choice of the factorization scale to avoid the parametrically large logarithms.
 ⇒ QCD rsummation for the enhanced logarithms with the standard RG formalism.
- RG evolution functions for the hard functions [Beneke, Rohrwild, 2011]:

$$egin{array}{rl} V^{(p)}_{7,\mathrm{eff}}(m_b,\mu,
u) &=& \hat{U}_1(m_b,\mu_h,\mu) \, V^{(p)}_{7,\mathrm{eff}}(m_b,\mu_h,
u) \,, \ K^{-1}(m_b,\mu) &=& \hat{U}_2(m_b,\mu_h,\mu) \, K^{-1}(m_b,\mu_h) \,. \end{array}$$

- Taking the factorization scale of order $\sqrt{m_b \Lambda_{QCD}} \Rightarrow$ No resummation for the jet function.
- Implementing the 2-loop evolution of φ⁺_B(ω, μ) as discussed before.

 → The NLL resummation improved expression for the LP decay amplitude:

$$\sum_{i=1}^{8} C_i F_{i,L}^{(p),\text{LP}} = -\frac{e_q f_{B_q}}{m_{B_q}} \hat{U}_1(m_b, \mu_h, \mu) \hat{U}_2(m_b, \mu_h, \mu) K^{-1}(m_b, \mu_h)$$
$$\overline{m}_b(\mathbf{v}) V_{7,\text{eff}}^{(p)}(m_b, \mu_h, \mathbf{v}) \mathscr{R}(m_b, \mu_0, \mu).$$

 \hookrightarrow Can be achieved alternatively in dual space [Shen, Wei, Zhao, Zhou, 2020].

Yu-Ming Wang (Nankai)

• The NLP factorization from the hard-collinear propagator:



• The NLP correlation function in coordinate space:

$$T_{7,\alpha\beta}^{\mathrm{hc,NLP}} = \left[-\frac{e_q \,\overline{m}_b(\mathbf{v}) m_{B_q}}{4 \, \pi^2} \right] \int d^4 x \, \frac{e^{i q \cdot x}}{x^2} \left(2 \, v_\mu - n_\mu \right) \\ \times \frac{\partial}{\partial x_\mu} \left\langle 0 | \bar{q}(x) \, \gamma_\beta^{\perp} \not n \, \vec{n} \, \gamma_\alpha^{\perp} \, P_R \, h_\nu(0) | \bar{B}_q \right\rangle.$$

• The classical equation of motion for the non-local operator [Kawamura et al, 2001]:

$$v_{\mu} \frac{\partial}{\partial x_{\mu}} \left[\bar{q}(x) \Gamma h_{\nu}(0) \right] = i \int_{0}^{1} du \, \bar{u} \, \bar{q}(x) \, g_{s} \, G_{\alpha\beta}(ux) \, x^{\alpha} \, v^{\beta} \, \Gamma h_{\nu}(0) + (v \cdot \partial) \left[\bar{q}(x) \, \Gamma h_{\nu}(0) \right].$$

 \Rightarrow The yielding NLP factorization formula

$$T_{7,\alpha\beta}^{\mathrm{hc,NLP}} = \left[-2ie_{q}\overline{m}_{b}(\mathbf{v})f_{B_{q}}m_{B_{q}}\right]\left[g_{\alpha\beta}^{\perp}-i\varepsilon_{\alpha\beta}^{\perp}\right]\left\{\frac{1}{2}-\left(\frac{\Lambda}{\lambda_{B_{q}}}\right)\right.\\ \left.+\int_{0}^{\infty}d\omega_{1}\int_{0}^{\infty}d\omega_{2}\frac{1}{\omega_{2}}\left[\frac{1}{\omega_{2}}\ln\frac{\omega_{1}}{\omega_{1}+\omega_{2}}+\frac{1}{\omega_{1}}\right]\Psi_{4}(\omega_{1},\omega_{2},\mu)\right\}.$$

Yu-Ming Wang (Nankai)

QCD and Heavy Quark Decays

2 7 3

• The NLP correction form the non-vanishing quark mass:

$$T_{7,\alpha\beta}^{m_q,\,\mathrm{NLP}} = \left[-ie_q \,\overline{m}_b(\mathbf{v}) \, m_q f_{B_q} \, m_{B_q}\right] \left[g_{\alpha\beta}^{\perp} - i\varepsilon_{\alpha\beta}^{\perp}\right] \int_0^{\infty} d\omega \, \frac{\phi_B^-(\omega,\mu)}{\omega} \, .$$

- Rapidity divergence implies the breakdown of the naive soft-collinear factorization.
- Nonperturbative parametrization of the convolution integral [Beneke, Neubert, 2003]:

- Alternative estimate from the LCSR method also possible.
- The complete SCET factorization in demand but extremely challenging.

Yu-Ming Wang (Nankai)

• The NLP contribution from the subleading SCET current [Beneke, Feldmann, 2002]

$$J^{(A2)} \supset (\bar{\xi}_{\overline{hc}} W_{\overline{hc}}) \gamma_{\alpha}^{\perp} P_L \left(\frac{i \overrightarrow{D}_{\top}}{2 m_b} \right) h_{\nu} + \dots, \qquad D_{\top}^{\mu} \equiv D^{\mu} - (\nu \cdot D) \nu^{\mu}.$$

Arise from the HQET representation of the QCD b-quark field.

• The resulting non-local hadronic matrix element

$$T_{7,\alpha\beta}^{A2,\text{NLP}} = \left[-\frac{ie_q m_{B_q}^2}{2} \right] \int d^4x \int \frac{d^4\ell}{(2\pi)^4} \exp\left[i(q-\ell) \cdot x\right] \frac{1}{\ell^2 + i0} \\ \times \langle 0|\bar{q}(x) \gamma_{\beta}^{\perp} \not n \not n \gamma_{\alpha}^{\perp} \not D P_L h_v(0)|\bar{B}_q \rangle \,.$$

 \Rightarrow The soft-collinear factorization formula with the aid of the HQET equations of motion

$$T_{7,\alpha\beta}^{42,\,\mathrm{NLP}} = \left[i e_q f_{B_q} m_{B_q}^2\right] \left[g_{\alpha\beta}^{\perp} - i \varepsilon_{\alpha\beta}^{\perp}\right] \left\{\frac{1}{2} \left(\frac{\bar{\Lambda}}{\lambda_{B_q}}\right) - 1 + \int_0^\infty d\omega_1 \int_0^\infty d\omega_2 \frac{1}{\omega_1(\omega_1 + \omega_2)} \Phi_3(\omega_1,\omega_2,\mu)\right\}.$$

- The convolution integral free of the end-point divergence.
- ► In agreement with the counterpart contribution to $B \rightarrow \gamma \ell \nu$ [Beneke, Braun, Ji, Wei, 2018].

Yu-Ming Wang (Nankai)

• The two-particle higher-twist *B*-meson LCDAs up to the $\mathcal{O}(x^2)$ accuracy:

$$\begin{split} \langle 0 | (\bar{q}_s Y_s)_\beta(x) \ (Y_s^{\dagger} h_v)_\alpha(0) | \bar{B}_q \rangle \\ &= -\frac{i \tilde{f}_{B_q}(\mu) m_{B_q}}{4} \int_0^\infty d\omega \, e^{-i\omega v \cdot x} \left[\frac{1+\psi}{2} \left\{ 2 \left[\phi_B^+(\omega,\mu) + x^2 \, g_B^+(\omega,\mu) \right] \right. \\ &\left. - \frac{1}{v \cdot x} \left[(\phi_B^+(\omega,\mu) - \phi_B^-(\omega,\mu)) + x^2 \, (g_B^+(\omega,\mu) - g_B^-(\omega,\mu)) \right] \right\} \gamma_s \right]_{\alpha\beta} \end{split}$$

- The three-particle higher-twist B-meson LCDAs up to twist-6 [Braun, Ji, Manashov, 2017].
 - The EOM relation between the two-particle and three-particle LCDAs.

$$-2\frac{d^2}{d\omega^2}g_B^-(\omega,\mu) = \left[\frac{3}{2} + (\omega-\bar{\Lambda})\frac{d}{d\omega}\right]\phi_B^-(\omega,\mu) - \frac{1}{2}\phi_B^+(\omega,\mu)$$
$$+\int_0^\infty \frac{d\omega_2}{\omega_2}\left[\frac{d}{d\omega} - \frac{1}{\omega_2}\right]\Psi_5(\omega,\omega_2,\mu) + \int_0^\omega \frac{d\omega_2}{\omega_2^2}\Psi_5(\omega-\omega_2,\omega_2,\mu).$$

- The systematic parametrization requires 8 independent LCDAs.
- The higher-twist factorization formula at tree level:

$$\begin{split} T_{7,\alpha\beta}^{\mathrm{HT,NLP}} \simeq \left[-ie_{q}\,\overline{m}_{b}(\mathbf{v})f_{B_{q}}\,m_{B_{q}}\right] \left[g_{\alpha\beta}^{\perp}-i\varepsilon_{\alpha\beta}^{\perp}\right] \left\{-1+2\int_{0}^{\infty}d\omega\ln\omega\Delta\phi_{B}^{-}(\omega,\mu)\right.\\ \left.-2\int_{0}^{\infty}d\omega_{2}\,\frac{1}{\omega_{2}}\,\Phi_{4}(\omega_{1}=0,\omega_{2},\mu)\right\}. \end{split}$$

Yu-Ming Wang (Nankai)

• The NLP contribution from the weak annihilation diagram:

The resulting helicity form factors:



- ► The weak-annihilation effect will spoil the large-recoil symmetry.
- Massive quark loops generate the non-trivial strong phase. \Leftrightarrow Dual to the final-state rescattering $\bar{B}_q \rightarrow H_c H_{\bar{c}} \rightarrow \gamma \gamma$ at hadronic level.
- Tree-operator contributions consistent with [Bosch, Buchalla, 2002].
- The NLP contribution from the (anti)-collinear photon radiation off the bottom-quark:

$$T_{7,\alpha\beta}^{e_b,\mathrm{NLP}} = \left[-ie_q f_{B_q} m_{B_q}^2\right] \left[g_{\alpha\beta}^{\perp} - i\varepsilon_{\alpha\beta}^{\perp}\right].$$

- Local correction preserves the large-recoil symmetry!
- ► In analogy to the P_7 contribution to $B_q \rightarrow \gamma \ell \bar{\ell}$ with the *B*-type insertion [Beneke, Bobeth, YMW, 2020].

Yu-Ming Wang (Nankai)

The resolved photon contribution

- The NLP contribution from the "hadronic" component of the on-shell photon [Ball, Braun, 2002].
- The dispersion technique [Khodjamirian, 1999; Braun, Khodjamirian, 2013; YMW, 2016]:

$$\begin{split} \tilde{T}_{7,\alpha\beta} &= 2\overline{m}_b(\mathbf{v}) \int d^4x e^{iq\cdot x} \left\langle 0 \right| \mathcal{T} \left\{ J^{\text{em}}_{\beta}(x), \bar{q}_L(0) \sigma_{\mu\alpha} p^{\mu} b_R(0) \right\} \left| \bar{B}_q(p+q) \right\rangle \Big|_{q^2 < 0} \\ &+ \left[p \leftrightarrow q, \alpha \leftrightarrow \beta \right] \\ &= -\frac{i e_q \overline{m}_b(\mathbf{v}) m^2_{B_q}}{2} \left\{ \left(g^{\perp}_{\alpha\beta} - i \varepsilon^{\perp}_{\alpha\beta} \right) \tilde{F}_{7,L}(\bar{n} \cdot q, n \cdot q) + \left[p \leftrightarrow q, \alpha \leftrightarrow \beta \right] \right\}. \end{split}$$

Power countering scheme:

$$\bar{n} \cdot q \sim \mathscr{O}(m_b), \qquad n \cdot q \sim \mathscr{O}(\Lambda).$$

• The hadronic dispersion relation:

$$\begin{split} \tilde{T}_{7,\alpha\beta} &= -i\overline{m}_{b}(\mathbf{v})m_{B_{q}}^{2}\left(g_{\alpha\beta}^{\perp}-i\varepsilon_{\alpha\beta}^{\perp}\right)\left\{\sum_{V}\frac{c_{V}f_{V}m_{V}T_{1}^{B_{q}\rightarrow V}\left(0\right)}{\bar{n}\cdot q\left(m_{V}^{2}/\bar{n}\cdot q-n\cdot q-i0\right)}\right.\\ &+ \int_{\omega_{s}}^{\infty}d\omega'\frac{\rho^{\mathrm{had}}(\bar{n}\cdot q,\omega')}{\omega'-n\cdot q-i0}\right\}.\end{split}$$

The constant c_V determined by the flavour factor and the electric charge of the QED quark-current.

• LCSR for the tensor $B \rightarrow V$ form factors:

$$\sum_{V} \frac{c_V f_V m_V}{\bar{n} \cdot q} \exp\left[-\frac{m_V^2}{\bar{n} \cdot q \,\omega_M}\right] T_1^{B_q \to V}(0) = \frac{e_q}{2} \frac{1}{\pi} \int_0^{\omega_s} d\omega' \, e^{-\omega'/\omega_M} \operatorname{Im}_{\omega'} \tilde{F}_{7,L}(\bar{n} \cdot q, \omega').$$

Yu-Ming Wang (Nankai)

The resolved photon contribution

• Improved dispersion relations (setting $n \cdot q = 0$) [Master formula]:

$$\begin{split} T_{7,\alpha\beta} &= -\frac{ie_q \,\overline{m}_b(\mathbf{v}) \, m_{B_q}^2}{2} \left(g_{\alpha\beta}^{\perp} - i \varepsilon_{\alpha\beta}^{\perp} \right) \left\{ \frac{1}{\pi} \, \int_0^\infty \frac{d\omega'}{\omega'} \, \mathrm{Im}_{\omega'} \, \tilde{F}_{7,L}(\bar{n} \cdot q, \omega') \right. \\ & + \underbrace{\frac{1}{\pi} \, \int_0^{\omega_{\mathrm{s}}} \, d\omega' \, \left[\frac{\bar{n} \cdot q}{m_V^2} \exp\left(\frac{m_V^2 - \bar{n} \cdot q \, \omega'}{\bar{n} \cdot q \, \omega_M} \right) - \frac{1}{\omega'} \right] \, \mathrm{Im}_{\omega'} \, \tilde{F}_{7,L}(\bar{n} \cdot q, \omega')} \right\} + \left[p \leftrightarrow q, \alpha \leftrightarrow \beta \right] \, . \end{split}$$

nonperturbative modification

• Power counting scheme for the sum-rule parameters:

$$\omega_{s} = \frac{s_{0}}{\bar{n} \cdot q} \sim \mathscr{O}\left(\frac{\Lambda_{\text{QCD}}^{2}}{m_{B_{q}}}\right), \qquad \omega_{M} = \frac{M^{2}}{\bar{n} \cdot q} \sim \mathscr{O}\left(\frac{\Lambda_{\text{QCD}}^{2}}{m_{B_{q}}}\right).$$

 \Rightarrow Nonperturbative modification yields the soft non-factorizable contribution.

• Spectral density at tree level:

$$\frac{1}{\pi} \operatorname{Im}_{\omega'} \tilde{F}_{7,L}(\bar{n} \cdot q, \omega') = f_{B_q} \underbrace{\phi_B^+(\omega', \mu)}_{O(1/\Lambda)} + \mathscr{O}(\alpha_s, \Lambda/m_b).$$
of $\mathscr{O}(1/\Lambda)[\mathscr{O}(1/m_b)]$ for $\omega' \sim \mathscr{O}(\Lambda)[\omega' \sim \mathscr{O}(\Lambda^2/m_b)]$

Power suppressed soft contribution!

• Alternative LCSR calculation with the subleading-twist photon LCDAs [Shen, YMW, 2018].

Yu-Ming Wang (Nankai)

Summary for the helicity amplitudes of $B_q \rightarrow \gamma \gamma$

• Final expressions for the factorized NLP corrections:

$$\begin{split} \sum_{i=1}^{8} C_{i} F_{i,L}^{(p), \text{fac}, \text{NLP}} &= C_{7}^{\text{eff}} \left[F_{7,L}^{\text{hc}, \text{NLP}} + F_{7,L}^{m_{q}, \text{NLP}} + F_{7,L}^{\text{A2}, \text{NLP}} + F_{7,L}^{\text{HT}, \text{NLP}} + F_{7,L}^{e_{b}, \text{NLP}} \right] \\ &+ \frac{f_{B_{q}}}{m_{B_{q}}} \left[\mathscr{F}_{V}^{(p), \text{WA}} - \mathscr{F}_{A}^{(p), \text{WA}} \right], \\ \sum_{i=1}^{8} C_{i} F_{i,R}^{(p), \text{fac}, \text{NLP}} &= \frac{f_{B_{q}}}{m_{B_{q}}} \left[\mathscr{F}_{V}^{(p), \text{WA}} + \mathscr{F}_{A}^{(p), \text{WA}} \right]. \end{split}$$

Large-recoil symmetry violation from the weak annihilation correction completely.

• Final expressions for the two helicity amplitudes:

$$\begin{split} \vec{\mathscr{A}_L} &= \sum_{p=u,c} V_{pb} \, V_{pq}^* \, \sum_{i=1}^8 \, C_i \, \left[F_{i,L}^{(p),\,\mathrm{LP}} + F_{i,L}^{(p),\,\mathrm{fac},\,\mathrm{NLP}} + F_{i,L}^{(p),\,\mathrm{soft},\,\mathrm{NLP}} \right] \,, \\ \vec{\mathscr{A}_R} &= \sum_{p=u,c} \, V_{pb} \, V_{pq}^* \, \sum_{i=1}^8 \, C_i \, \left[F_{i,R}^{(p),\,\mathrm{LP}} + F_{i,R}^{(p),\,\mathrm{fac},\,\mathrm{NLP}} + F_{i,R}^{(p),\,\mathrm{soft},\,\mathrm{NLP}} \right] \,. \end{split}$$

The fundamental nonperturbative functions: HQET *B*-meson LCDAs.
 ⇒ Key hadronic inputs for exclusive *B*-meson decay phenomenologies.

Yu-Ming Wang (Nankai)

B-meson distribution amplitudes in HQET

- The LP contribution depends on λ_{B_a} and the inverse-logarithmic moments.
- Applying the general ansatz of the 2- and 3- particle B-meson LCDAs

$$\begin{split} \phi_B^+(\boldsymbol{\omega},\boldsymbol{\mu}) &= \boldsymbol{\omega} f(\boldsymbol{\omega}), \ \Phi_3(\boldsymbol{\omega}_1,\boldsymbol{\omega}_2,\boldsymbol{\mu}_0) = -\frac{1}{2} \,\kappa(\boldsymbol{\mu}_0) \left[\lambda_E^2(\boldsymbol{\mu}_0) - \lambda_H^2(\boldsymbol{\mu}_0) \right] \,\boldsymbol{\omega}_1 \,\boldsymbol{\omega}_2^2 f'(\boldsymbol{\omega}_1 + \boldsymbol{\omega}_2), \\ \Phi_4(\boldsymbol{\omega}_1,\boldsymbol{\omega}_2,\boldsymbol{\mu}_0) &= \frac{1}{2} \,\kappa(\boldsymbol{\mu}_0) \left[\lambda_E^2(\boldsymbol{\mu}_0) + \lambda_H^2(\boldsymbol{\mu}_0) \right] \,\boldsymbol{\omega}_2^2 f(\boldsymbol{\omega}_1 + \boldsymbol{\omega}_2), \\ \Psi_4(\boldsymbol{\omega}_1,\boldsymbol{\omega}_2,\boldsymbol{\mu}_0) &= \kappa(\boldsymbol{\mu}_0) \,\lambda_E^2(\boldsymbol{\mu}_0) \,\boldsymbol{\omega}_1 \,\boldsymbol{\omega}_2 f(\boldsymbol{\omega}_1 + \boldsymbol{\omega}_2). \end{split}$$

The factorized NLP corrections can then be parameterized by the local HQET parameters.

• The NLP soft contribution sensitive to the precise shape of $\phi_B^+(\omega,\mu)$.

$$\phi_B^+(\omega,\mu_0) = \int_0^\infty ds \sqrt{ws} J_1(2\sqrt{ws}) \eta_+(s,\mu_0), \qquad \eta_+(s,\mu_0) = {}_1F_1(\alpha;\beta;-s\omega_0).$$

Such three-parameter ansatz [Beneke, Braun, Ji, Wei, 2018] is advantageous, since the resulting RG evolution can be done analytically in terms of $_2F_2$ functions.

Yu-Ming Wang (Nankai)

Theory predictions for the helicity amplitudes

• Breakdown of the various QCD mechanisms:



- NLL effects stabilize the factorizationscale dependence.
- Factorizable NLP effects around \$\mathcal{O}\$(30%).
- Destructive effects from the NLP soft corrections.
- Strong phase from the 2-loop matrix element of P^c₂ and the weak annihilation.

Yu-Ming Wang (Nankai)

Phenomenological observables for $B_q \rightarrow \gamma \gamma$

• Time-integrated branching fraction (for the flavour-tagged measurement):



The yielding theory predictions

$$\mathscr{BR}(B_d \to \gamma \gamma) = \left(1.44^{+1.35}_{-0.80}\right) \times 10^{-8}, \qquad \mathscr{BR}(B_s \to \gamma \gamma) = \left(3.17^{+1.96}_{-1.74}\right) \times 10^{-7}.$$

with the dominant uncertainties from λ_{B_q} , $\hat{\sigma}_{B_q}^{(1)}$, $\hat{\sigma}_{B_q}^{(2)}$ and the QCD renormalization scale v.

- Both the factorizable and soft NLP corrections are numerically important.
- The ratio of the two branching ratios for $B_{d,s} \rightarrow \gamma \gamma$

$$\frac{\mathscr{BR}(B_s \to \gamma\gamma)}{\mathscr{BR}(B_d \to \gamma\gamma)} = 33.80 \left(\frac{\lambda_{B_d}}{\lambda_{B_s}}\right)^2 + \mathcal{O}\left(\frac{\Lambda}{m_b}, \alpha_s\right)$$

The λ_{B_q} -scaling violation effect due to the NLP contributions approximately (10-20)%.

Yu-Ming Wang (Nankai)

Phenomenological observables for $B_q \rightarrow \gamma \gamma$

• Time-dependent CP asymmetries:

$$A_{\rm CP}^{\chi}(t) = \frac{\bar{\Gamma}^{\chi}(\bar{B}_q(t) \to \gamma\gamma) - \Gamma^{\chi}(B_q(t) \to \gamma\gamma)}{\bar{\Gamma}^{\chi}(\bar{B}_q(t) \to \gamma\gamma) + \Gamma^{\chi}(B_q(t) \to \gamma\gamma)} = -\frac{\mathscr{A}_{\rm CP}^{\rm dir,\,\chi}\cos(\Delta m_q t) + \mathscr{A}_{\rm CP}^{\rm mix,\,\chi}\sin(\Delta m_q t)}{\cosh(\Delta \Gamma_q t/2) + \mathscr{A}_{\Lambda\Gamma}^{\chi}\sinh(\Delta \Gamma_q t/2)}$$



Yu-Ming Wang (Nankai)

Current status of $B \rightarrow \gamma \ell \, \bar{\nu}_{\ell}$ (for Belle II)

- Factorization properties at leading power [Korchemsky, Pirjol, Yan, 2000; Descotes-Genon, Sachrajda, 2002; Lunghi, Pirjol, Wyler, 2003; Bosch, Hill, Lange, Neubert, 2003].
- Leading power contributions at NLL and (partial)-subleading power corrections at tree level [Beneke, Rohrwild, 2011].
- Subleading power corrections from the dispersion technique:
 - Soft two-particle correction at tree level [Braun, Khodjamirian, 2013].
 - Soft two-particle correction at one loop [YMW, 2016].
 - ▶ Three-particle *B*-meson DA's contribution at tree level [YMW, 2016; Beneke et al, 2018].
 - Subleading effective current and twist-5 and 6 corrections at tree level. [Beneke et al, 2018].
- Subleading power corrections from the direct QCD approach:
 - Hadronic photon corrections at tree level up to the twist-4 accuracy [Khodjamirian, Stoll, Wyler, 1995; Ali, Braun, 1995; Eilam, Halperin, Mendel, 1995].
 - Hadronic photon corrections of twist-two at one loop and of higher-twist at tree level [Ball, Kou, 2003; YMW, Shen, 2018].
- Leading power contributions at NNLL and the updated NLP corrections:
 - ► Two-loop RG evolution of $\phi_B^+(\omega, \mu)$ derived in [Braun, Ji, Manashov, 2019].
 - Two-loop jet function obtained in [Liu, Neubert, 2020].
 - Further improvement on $B \rightarrow \gamma \ell \bar{v}_{\ell}$ will appear soon [Cui, Shen, Wang, YMW, Wei, 2021].
 - Interesting generalization to $B_u^- \to \ell' \bar{\ell}' \ell \bar{\nu}_\ell$ in [Wang, YMW, Wei, 2021].

Theoretical wishlist

- Systematic understanding of the (higher-twist) *B*-meson distribution amplitudes.
 - Renormalization properties beyond the one-loop approximation [conformal symmetry].
 - Perturbative constraints at large ω_i [OPE technique].
 - Renormalon analysis and the renormalization-scheme dependence.
 - Precision determinations of the inverse moment λ_B .
- QCD factorization for the subleading power corrections.
 - SCET analysis for the pion-photon form factor as the first step [operator structures, symmetry constraints, etc].
 - General treatment of the rapidity divergences in the (naïve)-factorization formulae.
 - Rigorous factorization proof taking into account the Glauber gluons.
 - Novel resummation techniques for enhanced logarithms.
- Technical issues for future improvements.
 - Factorization techniques for the electromagnetic corrections.
 - ▶ NNLO QCD computations for $B \rightarrow V\gamma$ and $B \rightarrow V\ell\ell$.
 - QCD factorization for the radiative and electroweak penguin decays of the Λ_b -baryon.
 - Improved understanding of the parton-hadron duality violation.
- Very promising future for QCD aspects of heavy-quark physics!

Yu-Ming Wang (Nankai)