




Possible Bound States of $J/\psi J/\psi$

Xiang-Kun Dong


Feb. 17, 2022



- X.K. Dong, V. Baru, F.K. Guo, C. Hanhart, and A. Nefediev, [Phys.Rev.Lett. 126 \(2021\) 13, 132001](#)
- X.K. Dong, V. Baru, F.K. Guo, C. Hanhart, A. Nefediev and B.S. Zou, [Sci. Bull. 66 \(2021\) 24, 2462-2470](#)

- Coupled-channel interpretation of LHCb data
- Interactions between $J/\psi J/\psi$ from dispersive relation:
soft gluon exchange \rightarrow two pion exchange + ...
- Possible bound states of $J/\psi J/\psi$

$X(6900)$ in LHCb measurement



- ✓ A **narrow** resonance-like structure at 6.9 GeV, $X(6900)$
- ✓ A **broad** structure just above double- J/ψ threshold
- ✓ A **dip** near 6.7 GeV

➤ $X(6900)$ compact tetraquark?

Predictions dated back to 1970s

Y. Iwasaki, Prog.Theor.Phys. 1975;54:492

K.-T. Chao, Z.Phys.C7, 317(1981),

A.M. Badalian et al., PRD25,2370(1982), ...


Many theoretical investigations

...

Figure 3: Invariant mass spectra of weighted di- J/ψ candidates with $p_T^{\text{di-}J/\psi} > 5.2 \text{ GeV}/c$ and overlaid projections of the $p_T^{\text{di-}J/\psi}$ -threshold fit using (a) the NRSPS plus DPS model, (b) model I, and (c) model II.

LHCb, Sci.Bull.65,1983(2020)

$X(6900)$ in LHCb measurement



Discussions of general threshold behaviors, see X.-K. Dong, F.-K. Guo and B.-S. Zou PRL126, 152001(2021)

- Fully charmed tetraquark state?
 - 6.9 GeV **too high** for ground state
 - The gap 700 MeV **too large** for ground and 1st excited states
 - **No lighter** states (**easier**) observed
- Threshold effects. Near threshold,
 - **Breit-Wigner** fits **mislead** rather than **educate**
 - **Breit-Wigner** parameters (M and Γ) **hide nature** of states
 - Threshold effects sometimes play critical role
- Coupled-channel approach, Minimal models with X.-K. Dong et al, PRL126,132001(2021)
 - **most** relevant channels (2 models)
 - **minimal necessary** orders in interactions

Coupled-channel models

Two channel model

$J/\psi J/\psi$ & $\psi(2S) J/\psi$

$$V_{2\text{ch}}(E) = \begin{pmatrix} a_1 + b_1 k_1^2 & c \\ c & a_2 + b_2 k_2^2 \end{pmatrix}$$

Three channel model

$J/\psi J/\psi$, $\psi(2S) J/\psi$ & $\psi(3770) J/\psi$

$$V_{3\text{ch}}(E) = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{12} & a_{22} & a_{23} \\ a_{13} & a_{23} & a_{33} \end{pmatrix}$$


- η_c and h_c spin-0, HQSS
- $\chi_{cJ} \rightarrow \psi$ by ω exchange

Lippmann-Schwinger equation

$$T(E) = V(E) \cdot [1 - G(E)V(E)]^{-1}$$



Production amplitude in $J/\psi J/\psi$ channel (channel 1)

$$\mathcal{M}_1 = \alpha e^{-\beta E^2} \left[b + G_1(E)T_{11}(E) + \sum_{i=2,3} r_i G_i(E)T_{i1}(E) \right]$$





Hints of a $J/\psi J/\psi$ molecule

X.-K. Dong et al, PRL126,132001(2021)



	2-ch. fit	3-ch. fit 1	3-ch. fit 2
a_0 (fm)	≤ -0.49 or ≥ 0.48	$-0.61^{+0.29}_{-0.32}$	≤ -0.60 or ≥ 0.99
r_0 (fm)	$-2.18^{+0.66}_{-0.81}$	$-0.06^{+0.03}_{-0.04}$	$-0.09^{+0.08}_{-0.05}$
\bar{X}_A	$0.39^{+0.58}_{-0.12}$	$0.91^{+0.04}_{-0.07}$	$0.95^{+0.04}_{-0.06}$

X(6900) is uncertain
X(6200) is robust



➤ Compositeness of $X(6200)$



$$\bar{X}_A = (1 + 2|r_0/a_0|)^{-1/2}$$

$\bar{X}_A = 1$ for molecule and
 0 for compact state

I. Matusche et al, EPJA57(2021)3,101

➤ Large molecular component
in $X(6200)$ in 3-channel fit.

Hints of a $J/\psi J/\psi$ molecule




➤ Support of existence of $X(6200)$ from independent analysis

Z.-R. Liang et al, Phys.Rev.D
104 (2021) 3, 034034

Further Tests

- ✓ Data in the $\psi(2S)J/\psi$ channel \Rightarrow distinguish between the models
- ✓ Data in the $\eta_c\eta_c$ channel \Rightarrow verify predictive power of the models
- ✓ Data on $Y\Upsilon$ production \Rightarrow check in complementary sector
- ✓ Lattice simulation of double- J/ψ (η_c) scattering \Rightarrow independent test

X.-K. Dong et al, PRL126,132001(2021)



Binding mechanism??? \Rightarrow (dis)prove X(6200) nature!!!

- Van der Waals interaction between color dipoles
H. Fujii and D. Kharzeev, Phys.Rev.D 60, 114039 (1999).
- Long-range potential from two pion exchange between 2 S-wave bottomonia
N. Brambilla et al, Phys.Rev.D 93, 054002 (2016)

Evidence of an $\Omega_{ccc}\Omega_{ccc}$ bound state from HAL QCD method

Yan Lyu et al,
Phys.Rev.Lett. 127 (2021) 7, 072003

At long-range, soft gluon exchange \rightarrow two pion exchange + heavier...

OPE highly suppressed by isospin

Evidence of an $\Omega_{ccc}\Omega_{ccc}$ bound state

Yan Lyu et al, *Phys.Rev.Lett.* 127 (2021) 7, 072003

Evidence of an $\Omega_{ccc}\Omega_{ccc}$ bound state from HAL QCD method




FIG. 1. The $\Omega_{ccc}\Omega_{ccc}$ potential $V(r)$ in the 1S_0 channel as a function of separation r at Euclidean time $t/a = 25$ (red square), 26 (blue diamond), and 27 (green circle).





FIG. 4. The dimensionless ratio of the effective range r_{eff} and the scattering length a_0 as a function of r_{eff} . The red up(down)-pointing triangle and the blue right(left)-pointing triangle correspond to $\Omega_{ccc}\Omega_{ccc}$ system and $\Omega\Omega$ system in the 1S_0 channel with (without) the Coulomb repulsion, respectively. The black circle represents NN system in the 3S_1 - 3D_1 channel. The green square (nn) and diamond (pp) correspond to NN system in the 1S_0 channel. The error bars for $\Omega_{ccc}\Omega_{ccc}$ are the quadrature of the statistical and systematic errors in Eqs. (4) and (8).

Two pion/kaon exchange potential



➤ J/ψ / ψ $\pi\pi$ coupling $\mathcal{M}_\pi(t) =$

➤ $\psi\psi\pi\pi$ couplings related to the chromopolarisabilities, α_{AB} , satisfying [A. Sibirtsev et al, PRD71, 076005\(2005\)](#)

$$\alpha_{J/\psi J/\psi} \alpha_{\psi(2S)\psi(2S)} \geq \alpha_{\psi(2S)J/\psi}^2$$

➤ Chiral Lagrangian (& HQSS)

$$\mathcal{L}_{\psi_\alpha \psi_\beta \Phi \Phi} = \frac{c_1^{(\alpha\beta)}}{2} \langle J_\beta^\dagger J_\alpha \rangle \text{Tr}[u_\mu u^\mu] + \frac{c_2^{(\alpha\beta)}}{2} \langle J_\beta^\dagger J_\alpha \rangle \text{Tr}[u_\mu u_\nu v^\mu v^\nu] + \text{H.c.},$$



LECs \rightarrow $J/\psi, \psi(2S), \dots$ \rightarrow π, K, \dots

➤ Non-perturbative interaction of scalar-isoscalar $\pi\pi$ and $K\bar{K}$ channel is necessary.

✓ A polynomial times Omnès function

[Omnès, Nuovo Cim. 8 \(1958\) 316](#)

Unitarity



Two pion/kaon exchange potential

- $c_i^{(12)}$ from fits to BESII data on $\psi(2S) \rightarrow J/\psi\pi\pi$ decay.

[BES Collaboration], PLB645, 19 (2007)

- Difference between c_i^{11} and c_i^{12}

$$\xi \equiv \frac{\alpha_{J/\psi J/\psi}}{\alpha_{\psi(2S) J/\psi}} \approx 1 \sim 3$$

estimated by

- overlap of quark model wavefunctions

$$\alpha_{AB} \propto \int d^3r \psi_A^*(\vec{r}) e^{-i\vec{q}_c \cdot \vec{r}/2} \psi_B(\vec{r}) \equiv I_{AB}(q_c)$$



- S -Wave $J/\psi\pi$ scattering length

$$a_0 = \frac{(c_1^{(11)} + c_2^{(11)})m_\pi^2}{2\pi F_\pi^2(M_{J/\psi} + m_\pi)} \approx 0.0036 \xi \text{ fm}$$

$$|a_0^{\text{lat}}| \sim 0.01 \text{ fm}$$

K. Yokokawa et al. PRD74, 034504 (2006)
 L. Liu et al. PoS LATTICE2008, 112 (2008)
 X.-H. Liu et al. EPJC73, 2284(2013)


P -Wave $J/\psi\pi$
 Scattering
 Lattice QCD



$$\mathcal{M}_1[J/\psi\pi \rightarrow J/\psi\pi] \approx 8\pi (M_{J/\psi} + m_\pi) k^2 a_1$$

$$a_1 = -\frac{M_{J/\psi} c_1 - m_\pi c_2}{6\pi F_\pi^2 M_{J/\psi} (M_{J/\psi} + m_\pi)} \approx -\frac{c_1}{6\pi F_\pi^2 M_{J/\psi}}$$

Potentials from dispersive relation

➤ **Dispersive relation** $\mathcal{M}_\pi(t) =$  $= \frac{1}{\pi} \int_{4m_\pi^2}^{\infty} dt' \frac{\text{Im}\mathcal{M}_\pi(t')}{t' - t - i\epsilon}$


➤ **Two pion (kaon) exchange potential**

$$\mathcal{M}_{J/\psi J/\psi}(t) = \mathcal{M}_\pi(t) + \mathcal{M}_K(t)$$

$$V_{\text{exch}}(q, \Lambda) = \frac{-1}{4\pi M_{J/\psi}^2}$$

$$\times \int_{4m_\pi^2}^{\infty} d\mu^2 \frac{\text{Im}\mathcal{M}_{J/\psi J/\psi}(\mu^2)}{\mu^2 + q^2} e^{-\frac{q^2 + \mu^2}{\Lambda^2}}$$

Form factor to regularize the UV divergence in both q and μ and keep the long-range potential unchanged.



$J/\psi J/\psi$ molecular state

- Renormalization with a **contact term** $V_{\text{CT}}(q, \Lambda) = C e^{-\frac{q^2}{\Lambda^2}}$

$$V_{\text{tot}}(q, \Lambda) = V_{\text{CT}}(q, \Lambda) + V_{\text{exch}}(q, \Lambda)$$

- Lippmann Schwinger equation


$$T(E; k', k) = V_{\text{tot}}^S(k', k, \Lambda) + \int \frac{d^3l}{(2\pi)^3} \frac{V_{\text{tot}}^S(k', l, \Lambda) T(E; l, k)}{E - l^2/M_{J/\psi} + i\epsilon}$$

- Parameters:

$\Lambda(1 \sim 3 \text{ GeV})$, $\xi(1 \sim 3)$ and V_{ct} in GeV^{-2}

- Poles for $\xi = 1$.

Solid for bound states and dashed for virtual states





$J/\psi J/\psi$ molecular state

Strategy:

- ✓ Near threshold poles (bound or virtual), $E_B = 1$ or 5 MeV fixed (hints from LHCb data)
- ✓ $\xi (=1,2,3)$ fixed
- ✓ Contribution of V_{exch} to the binding, R

$$R \equiv \frac{V_{\text{exch}}^S(k'=0, k=0, \Lambda)}{V_{\text{tot}}^S(k'=0, k=0, \Lambda)}$$




$e^{-(q^2 + \mu^2)^n / \Lambda^{2n}}$ for meson exchanges
 $e^{-q^{2n} / \Lambda^{2n}}$ for the contact term

Weak dependence on n

Plausible?

- Contact term + two pion/kaon exchange leads to a molecule of $J/\psi J/\psi$
- $J/\psi J/\psi$ interaction suppressed by OZI or $\Lambda_{\text{QCD}}^2/m_c^2$, **no reason** for $V_{\text{ct}} \gg V_{\text{exch}}$
- **Naturalness**: contact term is of the same order as two pion/kaon exchange
- Reasonable cutoff, **1 ~ 1.5 GeV**
- We take it **plausible** if **two pion/kaon** exchange has **sizeable** contributions to the binding of $J/\psi J/\psi$, characterized by the ratio

$$R \equiv \frac{V_{\text{exch}}^S(k'=0, k=0, \Lambda)}{V_{\text{tot}}^S(k'=0, k=0, \Lambda)} \gtrsim 1/2$$




Summary



- LHCb $J/\psi J/\psi$ data can be well described by the coupled-channel method.
- Hints of a state near $J/\psi J/\psi$ threshold, with large **molecular** component.
- **Soft gluon** exchange between $J/\psi J/\psi$ described by **two pion and kaon** exchange.
- Coupling constants of $\psi(2S)J/\psi\pi\pi$ coupling extracted from BESII data on $\psi(2S) \rightarrow J/\psi\pi\pi$ decay.
- $J/\psi J/\psi\pi\pi$ coupling is argued to be **larger** than $\psi(2S)J/\psi\pi\pi$ coupling.
- With reasonable cutoff Λ , two pion and kaon exchanges provide **sizeable** contribution to the $J/\psi J/\psi$ attraction.
- The **binding of $J/\psi J/\psi$** system is **plausible**, given our current understanding of hadron-hadron interaction.

- ✓ Data in the $\psi(2S)J/\psi$ channel \Rightarrow distinguish between the models
- ✓ Data in the $\eta_c\eta_c$ channel \Rightarrow verify predictive power of the models
- ✓ Data on $\Upsilon\Upsilon$ production \Rightarrow check in complementary sector
- ✓ Lattice simulation of **double- J/ψ (η_c) scattering** \Rightarrow independent test
- ✓ Take a look at $J/\psi e^+e^-$ or $J/\psi\mu^+\mu^-$ channels
- ✓ **Lattice simulations of S - and P -wave $J/\psi\pi$ scattering** $\Rightarrow c_1$ and c_2



THANK YOU . 