

New Physics at CEPC

Xuai Zhuang (IHEP)

zhuangxa@ihep.ac.cn

高能物理大会

Aug. 8-11, 2022



中国科学院高能物理研究所
Institute of High Energy Physics
Chinese Academy of Sciences

Introduction

- Besides as a higgs factory, CEPC has a good potential to search for the direct production of new physics states
- With a very clean collision background, CEPC has the discovery advantage in many scenarios which are challenging at hadron colliders (large Bgs, large pile-up, trigger constraints from high energy objects, and difficulties in obj. Rec and ID, ...).
 - ✓ Exotic Higgs
 - ✓ SUSY
 - ✓ Dark Matter or Dark sector
 - ✓ Long-lived particles
 - ✓ More exotics: Heavy neutrinos, Axion-like particles, EW phase transition, ...

Brief summary of BSM search @CEPC


- BSM working group formed @ 2021.4 Yangzhou WS
- Big updates presented
 - @ 2021.11 CEPC WS (13 talks)
 - @ 2022.5 CEPC WS (17 talks)
- BSM white paper is scheduled and going-on smoothly:
 - Preliminary organizers: Liantao Wang, Bruce Mellado, Xuai Zhuang, Jia Liu
 - ✓ **More to be invited, volunteers are very welcome!**
 - Timeline (TBD): collect inputs and a very brief white paper draft ready by end of 2022; First paper draft is ready by next Spring?
- BSM prospects at CEPC are included in CEPC snowmass white paper: [arXiv:2205.08553](https://arxiv.org/abs/2205.08553)

BSM Inputs & Status

- BSM Higgs (1709.06103; 1808.02037; 1912.01431; 2008.05492; 2011.04540)
- SUSY Searches
 - Direct SUSY Searches (CPC46(2022)013106; 2101.12131; 2203.10580; 2202.11011)
 - Indirect search of SUSY (2010.09782)
 - Global fit of SUSY (2203.04828)
- Dark Matter and Dark Sector searches
 - Lepton portal DM (JHEP 06 (2021) 149)
 - Asymmetric DM (PRD 104(2021)055008)
 - Dark Sector from exotic Z decay (1712.07237)
 - DM (Millicharged DM, Vector portal DM, DM with EFT interactions): 1903.1211
 - Mono-gamma (2205.05560)
- Long-lived particles (1904.10661, 1911.06576, 2201.08960)
- More exotics:
 - Heavy neutrinos (2102.12826);
 - Axion-like particles (2103.05218, 2204.04702, Jia Liu's [talk](#))
 - Electroweak phase transition (1911.10210, 1911.10206, 2011.04540)
 -

BSM Higgs

- A large class of BSM physics, such as singlet extensions, two Higgs-doublet-models (2HDM), SUSY models, Higgs portals, gauge extensions of the SM, motivates these exotic decay considerations.




Representative topologies of the Higgs exotic decays

- Reference:

- 2HDM searches: 1709.06103; 1808.02037; 1912.01431; 2008.05492; 2011.04540
- Exotic higgs decay: 1612.09284 , 2110.13225 , 2203.08206, 2002.05554 , 2003.01662 , 2006.03527 ...
- Summarized at [2205.08553](#).

Exotic Higgs decay

- Exotic decays of the 125 GeV Higgs boson at future e^+e^- lepton colliders, Z. Liu, L.-T. Wang, and H. Zhang, [1612.09284](#)
- Exotic Higgs Decays to Four Taos at Future Electron-Positron Colliders, J. Shelton and D. Xu, [2110.13225](#)
- CEPC is very sensitive for signals with jets, heavy quarks and taus, which is challenge at LHC





The 95% C.L. upper limit on selected Higgs exotic decay BR

The CEPC* scenario further utilizes the hadronically decaying Z boson and includes an estimated (indicative) improvement of 40%.



SUSY Searches at CEPC

Reference: mainly light EWKino and slepton for CEPC

- Electroweakino (wino, higgsino) search: CPC46(2022)013106
- Bino NLSP at CEP: 2101.12131
- Slepton search: 2203.10580
- Heavy selectron search: 2202.11011
- Indirect search of SUSY: 2010.09782
- Global fit of SUSY: 2203.04828



Higgsino-like EWK processes









CEPC/FCCee(240)

CEPC



ILC 500/CEPC240: discovery in all scenarios up to kinematic limit: $\sqrt{s}/2$

SUSY Searches at CEPC



SUSY global fits with CEPC using GAMBIT




- Study of the impact of the Higgs and electroweak precision measurements at the CEPC with GAMBIT global fits of the SUSY models, such as CMSSM, NUHM1, NUHM2 and pMSSM-7, Yang Zhang etc, [arXiv: 2203.04828](https://arxiv.org/abs/2203.04828)
- CEPC can further test the currently allowed parameter space of these models, advance our understanding of the mass spectrum



Dark Matter and Dark Sector searches



■ Reference:

- Lepton portal DM (JHEP 06 (2021) 149)
- Asymmetric DM (PRD 104(2021)055008)
- Dark Sector from exotic Z decay (1712.07237)
- DM (Millicharged DM, Vector portal DM, DM with EFT interactions):
1903.1211
- Mono-gamma (2205.05560)



DM search at CEPC

- Exposing Dark Sector via exotic Z-boson decay with Future Z-Factories, Jia Liu, Lian-Tao Wang, Xiao-Ping Wang, Wei Xue, [1712.07237](#), PRD 97, 095044 (2018)
- Four models include: Higgs/Vector portal DM, inelastic dark matter and axion like particles.
- Compared with HL-LHC, the reach for the BR of various exotic Z decay modes at Z-factories is sensitive for many decay modes.



[arXiv: 1712.07237](#)


Long-lived particles (LLP)

■ Reference:

- LLP at near Detector: 1904.10661
- LLP at Far Detector: 1911.06576, 2201.0896
- LL Dark Hadrons: 2110.10691
- On-going: Yulei Zhang: [Talk](#)


Long lifetimes result from a few simple physical mechanisms:

- Small couplings (ex. RPV SUSY)
- Limited phase space: small mass splitting (ex. compressed SUSY, ...)
- Heavy intermediate states
-




LLP at Far Detector (FD)

- Physics with Far Detectors at Future Lepton Colliders, Zeren Simon Wang, Kechen Wang, [1911.06576](#), PRD 101, 075046 (2020)
- Search for long-lived axions with far detectors at future lepton colliders, Minglun Tian, Kechen Wang, Zeren Simon Wang, [2201.08960](#)
- FD can extend and complement the sensitivity to the LLPs compared with Near Detector




Light Scalars from
Exotic Higgs Decays



[1911.06576](#)

Light Neutralinos
from Z-boson Decays



Axion-like Particles

More exotics

■ Reference:



- Heavy neutrinos (2102.12826);
- Axion-like particles (2103.05218, 2204.04702, Jia Liu's [talk](#))
- Electroweak phase transition (1911.10210, 1911.10206, 2011.04540,)
-

Axion-like particles (ALP)



- Searching for ALP at future electron-positron colliders, H. Y. Zhang, C.X. Yue, Y.C. Guo, and S. Yang, [2103.05218](#), PRD104 (2021) 096008

→ CEPC is more sensitive to the ALPs couplings $g_{a\gamma\gamma}$ with mass 2-8 GeV than LHC and CLIC.

- Searching for ALP via decay $Z \rightarrow a f \bar{f}$ at future Z factories, [2204.04702](#)
- Axion-like particle solution to muon g-2 and its test at Z-factory, Jia Liu's [talk](#)



M_a (GeV)



Summary and Outlook

- **CEPC has good discovery potential for NP at many scenarios which are challenge for LHC**
- **BSM prospects study at CEPC is going on well, many of the analyses are already public**
- **Plan to organize a workshop by end of this year to collect inputs for CEPC BSM white paper**
- **Please let us know if you would like to contribute to the BSM white paper !**


Thanks for your attention!

Backup

17




Wino & higgsino

- Prospects for chargino pair production at CEPC, Jia-Rong Yuan, Hua-Jie Cheng, Xu-Ai Zhuang, [arXiv:2105.06135](https://arxiv.org/abs/2105.06135).



Chargino pair via
on(off)-shell W decay



Signature:
2 lepton + MET



Discovery in
all scenarios
up to
kinematic
limit: $\sqrt{s}/2$






Bino NLSP at CEPC

- Probing bino NLSP at lepton colliders with Gravitino DM, Junmou Chen, Chengcheng Han, Jin Min Yang, Mengchao Zhang, [arXiv:2101.12131](https://arxiv.org/abs/2101.12131).



Slepton search


■ Prospects for slepton pair production at CEPC, Jia-Rong Yuan, Hua-Jie Cheng, Xu-Ai Zhuang, [arXiv: 2203.10580](https://arxiv.org/abs/2203.10580)




Heavy selectron search

- Probing relatively heavier right-handed selectron in the GmSUGRA, by Waqas Ahmed, Imtiaz Khan, Tianjun Li, Shabbar Raza and Wenxing Zhang, [arXiv: 2202.11011](#)
- There two types of light neutralinos that achieve the correct relic density by Z-resonance and h-resonance.

Higgs-pole $\rightarrow m_{\tilde{\chi}_1^0} \approx \frac{1}{2}m_h$ and Z-pole $\rightarrow m_{\tilde{\chi}_1^0} \approx \frac{1}{2}m_Z$.



$$e^+e^- \rightarrow \tilde{\chi}_1^0(\text{bino}) + \tilde{\chi}_1^0(\text{bino}) + \gamma$$




[arXiv: 2202.11011](#)

DM search at CEPC

- Searching for lepton portal dark matter with colliders and interplay with the gravitational wave (GW) astronomy, Jia Liu, Xiao-Ping Wang, KePan Xie, [2104.06421](#), JHEP 06 (2021) 149
- The phase transition GWs can also be a probe of the model.

$$e^+e^- \rightarrow S^{\pm(*)}S^\mp \rightarrow \ell^+\chi\ell'^-\chi \quad h/Z \rightarrow S^{\pm(*)}S^\mp \rightarrow \ell^+\chi\ell'^-\chi \text{ and } h \rightarrow \chi\chi:$$





[arXiv: 2104.06421](#)

FIG. 10. Figure from Ref. [168], the interplay between GW detection and future e^+e^- collider searches. The gray shaded region is the LISA detectable parameter space. From left to right, the sensitivities for λ_{HS} are shown from future CEPC precision measurements, in which the region above a given m_χ (corresponding to a colored line) can be probed.

DM search at CEPC

- Searching for asymmetric Dark Matter (ADM) at CEPC, Mengchao Zhang, [2104.06988](#), PRD 104, 055008 (2021)
- It is possible to generate dark quark pair through a t-channel process, dark quark q' will be a jet-like object in detector.



- The mass of mediator can be excluded up to $O(10)$ TeV, better than LHC

DM search at CEPC



■ Probing DM particles at CEPC (Millicharged DM, Vector portal DM, DM with EFT interactions): ZL, Yong-Heng Xu, Yu Zhang ,1903.1211

- ✓ CEPC can probe **millicharged DM** that is currently unexplored
- ✓ CEPC can probe the parameter space of vector-portal DM models and EFT DM models that are unconstrained by DMDD

■ Mono- γ Production of a Vector Dark Matter at CEPC, K Ma, 2205.05560

[ZL, Y.-H. Xu, and Y. Zhang,1903.12114]

new physics process: $e^+e^- \rightarrow \bar{\chi}\chi\gamma$



LL Dark Hadrons



- A theory of Dark Pions, Hsin-Chia Cheng, Lingfeng Li, Ennio Salvioni, [2110.10691](#), JHEP 01 (2022) 122, see Lingfeng's [talk](#)
- The dark quarks couple to the SM via irrelevant Z- and Higgs-portal operators. The dark pions, behave as either composite axion-like particles (ALPs) mixing with Z or h

arXiv: 2110.10691

$$\mathcal{L}_{\text{EFT}} = \frac{1}{2} \bar{\psi}_R Y^\dagger M^{-2} Y [|H|^2 i \not{D} + i \gamma^\mu H^\dagger D_\mu H] \psi_R + \text{h.c.} \\ + \frac{1}{2} \bar{\psi}_L \tilde{Y}^\dagger M^{-2} \tilde{Y} [|H|^2 i \not{D} + i \gamma^\mu H^\dagger D_\mu H] \psi_L + \text{h.c.} \\ - \bar{\psi}_L \omega \psi_R + \boxed{\bar{\psi}_L \tilde{Y}^\dagger M^{-1} Y \psi_R |H|^2} + \text{h.c.},$$

Dimension-6 Z portal couplings


Dimension-5 Higgs portal coupling



- $m_\pi < m_{\eta'}$: dimuon mode dominates
- $m_\pi > m_{\eta'}$: PPP modes (mostly SM $\pi^+ \pi^- \pi^0$)

LLP at near Detector (ND)

- Long-lived light neutralinos at future Z-factories (RPV SUSY), Zeren Simon Wang, Kechen Wang, [1904.10661](#), PRD 101, 115018 (2020)
- The model parameter $\lambda'_{112}/m_{\tilde{f}}^2$ can be discovered down to as low as $\sim 1.5 \times 10^{-14}$ (3.9×10^{-14}) GeV^{-2} at the FCC-ee (CEPC)



[arXiv: 1904.10661](#)



Muon Anomalous Magnetic Moment

Present status: Discrepancy between Theory and Experiment at more than three Standard Deviation level

$$\delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{theory}} = 268(63)(43) \times 10^{-11}$$

3.6 σ Discrepancy

New Physics at the Weak scale can fix this discrepancy. Relevant example : Supersymmetry



$$\delta a_\mu \simeq \frac{\alpha}{8\pi s_W^2} \frac{m_\mu^2}{\tilde{m}^2} \text{Sgn}(\mu M_2) \tan \beta \simeq 130 \times 10^{-11} \left(\frac{100 \text{ GeV}}{\tilde{m}} \right)^2 \text{Sgn}(\mu M_2) \tan \beta$$

Grifols, Mendez'85, T. Moroi'95,
Giudice, Carena, C.W.'95, Martin and Wells'00

Here \tilde{m} represents the weakly interacting supersymmetric particle masses.

For $\tan \beta \simeq 10$ (50), values of $\tilde{m} \simeq 230$ (510) GeV would be preferred.

Masses of the order of the weak scale lead to a natural explanation of the observed anomaly !