

## 喷注物理实验综述

*Tuesday, 9 August 2022 14:00 (25 minutes)*

The energy loss of jets (jet quenching) is one of the most important signatures of the deconfined state of quarks and gluons (Quark-Gluon Plasma) created in Pb–Pb collisions at the LHC. The measurement of jets recoiling from a trigger hadron uniquely enables the exploration of medium-induced modification of jet production. Jet deflection via multiple soft scatterings with the medium constituents may result in a broadening of the overall azimuthal correlation between the trigger hadron and the recoiling jet. In addition, the tail of this azimuthal correlation is sensitive to single-hard Molière scatterings off quasi-particles in the medium. The overall yield and  $R$ -dependence of the recoil jets of also offers important information about jet energy loss and intra-jet broadening.

This contribution presents a measurement of charged jets recoiling from a trigger hadron in pp and Pb–Pb collisions at  $\sqrt{s_{\text{NN}}} = 5.02$  TeV. Techniques are employed which allow for a precise data-driven subtraction of the large uncorrelated background contaminating the measurement in Pb–Pb collisions, enabling the exploration of medium-induced modification of jet production and acoplanarity over a wide phase space, including the low jet  $p_{\text{T}}$  region for large jet resolution parameter  $R$ .

**Primary author:** MAO, Yaxian (Central China Normal University)

**Presenter:** MAO, Yaxian (Central China Normal University)

**Session Classification:** Parallel Session I (3): Heavy Ion Physics

**Track Classification:** 重离子物理