

# An Introduction to perturbative QCD

X. Liu

xiliu@bnu.edu.cn

北京师范大学

following Jian's Lecture

2022年理论物理前沿讲习班.

暗物质与新物理暑期学校


④ 北京师范大学

07/04/2022 - 07/22/2022

# Why (perturbative) QCD?

$$\alpha \sim \alpha_w \sim 10^{-2} \ll \alpha_s(M_w) \sim 0.1$$

⇒ More QCD activities at a collider.



a typical event @ LHC

$$\sigma = \sum_{ij} \hat{\sigma}_{ij}^{e^{+}e^{-}} \cdot F_{i/P} F_{j/P} \cdot D_e(u)$$

$$\approx \sigma(\mu_0 \sim 100 \text{ GeV}) \hat{\sigma}(\mu)$$


Possible other scales introduced by kinematics or cuts



evolution, resummation, parton shower deal with log of scales

$$\sim \sigma(\mu_{\text{IR}}) F(\mu_2 = 100 \text{ GeV}) D(\mu_0 = 100 \text{ GeV})$$

- dominant background to New physics Searches,



New physics  $\sim$  Data - QCD prediction.

- Fundamentals to the Monte-Carlo tools

e.g. Madgraph, pythia, Sherpa, ...

- Outline

o thrust as an example

- NLO calculation

- infrared safe

- break-down of fixed order


o Infrared behaviour in QCD/QED

- Coherent branching

- DL Resummation

## Thrust in $e^+e^-$ -annihilation

Consider



$$\sigma = \frac{1}{4} \frac{1}{2S} \int d\Phi_N L_{\mu\nu}(l, \bar{l}) \frac{1}{S^2} H^{\mu\nu}(q, \Phi_N) \Theta(\Phi_N)$$

$\int$  Spin ave.       $\int$  Flux       $\int$  Phase space of  $N$  final particles.

scalar  $\rightarrow$   $\frac{1}{4} \frac{1}{2S^3} L_{\mu\nu}(l, \bar{l}) \int d\Phi_N \left( g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) H(q, \Phi_N) \Theta(\Phi_N)$

gauge

$$= \frac{1}{4} \frac{1}{2S^3} L_{\mu\nu}^{\mu\nu}(l, \bar{l}) \int d\Phi_N H(q, \Phi_N) \Theta(\Phi_N)$$


where

$$H = \frac{1}{d-1} \left( g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) H^{\mu\nu}$$

$$L_{\mu\nu}^{\mu\nu} = 4(2l\bar{l} - d l\bar{l}) = -2(d-2)S$$

For large  $s = Q^2 \gg \Lambda_{QCD}$ , it is well modeled by

perturbative calculation



$L_0$ ,

$$H^{(1)} = 4(\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu - g^{\mu\nu} \gamma^\alpha \gamma^\alpha)$$

$$H^{(2)} = -\frac{1}{d-1} 2(d-2)s$$

$$= \frac{2Q_F^2 N_c e^4}{f} \frac{1}{4} \frac{1}{2S^3} \frac{[4(1-\epsilon)S]^2}{d-1} \int d\Phi_2 \Theta(\Phi_2) = 1$$

$$= \frac{2Q_F^2 N_c e^4}{f} \frac{1}{4} \frac{1}{2S} \frac{16(1-\epsilon)^2}{3-2\epsilon} (2\pi) \frac{1}{4} \frac{\Omega_{d-2}}{(2\pi)^{d-1}} S^{-\epsilon} \int_0^1 dx x^{-\epsilon} (1-x)^{-\epsilon}$$

where  $x \equiv \frac{-t}{s}$ ,  $\Omega_{d-2} = \frac{2\pi^{1-\epsilon}}{\Gamma(1-\epsilon)}$  is the solid angle

$$\Rightarrow G_0 = \frac{2Q_F^2 N_c}{f} \frac{4\pi\alpha^2}{S} \frac{1-\epsilon}{3-2\epsilon} \frac{\Gamma(2-\epsilon)}{\Gamma(2-2\epsilon)} \left(\frac{4\pi\alpha^2}{S}\right)^\epsilon$$

$$= \frac{2Q_F^2 N_c}{f} \frac{4\pi\alpha^2}{3S} + \mathcal{O}(\epsilon)$$

$$\textcircled{*} \int d\Phi_2 = \int \frac{d^d q}{(2\pi)^{d-1}} \delta(q^2) \frac{d\bar{q}}{(2\pi)^{d-1}} \delta(\bar{q}^2) (2\pi)^d \delta^d(Q - q - \bar{q})$$

$$= (2\pi) \int \frac{d^d q}{(2\pi)^{d-1}} \delta(q^2) \delta((Q - q)^2) \stackrel{\text{circled } \infty}{=} \begin{aligned} &= Q^2 - 2Q \cdot q \\ &= s + u + t \end{aligned}$$

$$= (2\pi) \int \frac{d^d q}{(2\pi)^{d-1}} \delta(q^2) \delta(s + u + t)$$

$$\text{let } q^\mu = \frac{2q \cdot \bar{l}}{2l \cdot \bar{l}} l^\mu + \frac{2q \cdot l}{2l \cdot \bar{l}} \bar{l}^\mu + q_\perp^\mu$$

$$= -\frac{u}{s} l^\mu + \frac{-t}{s} \bar{l}^\mu + q_\perp^\mu$$

$$\text{Hence } q^2 = \frac{4t}{s} - q_\perp^2$$

$$\text{and } d^d q = \frac{1}{2} \frac{d(-u)}{s} \frac{d(-t)}{s} d^{d-2} q_\perp$$

$$= \frac{1}{4} \frac{d(-u)}{s} \frac{d(-t)}{s} (q_\perp^2)^{\frac{d-2}{2}} d^{d-2} q_\perp$$

$$\Rightarrow \int d\Phi_2 = \frac{(2\pi)}{(2\pi)^{d-1}} \frac{1}{4} d\Omega_{d-2} S^{-6} \int_0^1 dx x^{-6} (1-x)^{-6}$$

for NLO, we have both virtual and real corrections

Virtual :



+ C.T.



Scaleless  $\frac{1}{\epsilon_{UV}} - \frac{1}{\epsilon_{IR}}$   
renorm.

$$\square = M_0^m \frac{d^d s}{4\pi} C_F \left( \frac{4\pi M^2}{-s} \right)^\epsilon \underbrace{\left( -\frac{2}{\epsilon^2} + \frac{1}{\epsilon} - 2 \right)}_{\text{IR Poles}} \frac{\Gamma^2(1-\epsilon)\Gamma(1+\epsilon)}{\Gamma(2-2\epsilon)} \frac{\epsilon \bar{e}}{e}$$

$\Rightarrow$  Virt

$$= g_0 \frac{d^d s}{2\pi} C_F \left( \frac{4\pi M^2}{s} \right)^\epsilon e^{\gamma_E \epsilon} \cos \pi \epsilon \left( -\frac{2}{\epsilon^2} + \frac{1}{\epsilon} - 2 \right) \frac{\Gamma^2(1-\epsilon)\Gamma(1+\epsilon)}{\Gamma(2-2\epsilon)}$$

$$= g_0 \frac{d^d s}{2\pi} C_F \left( \frac{4\pi M^2}{s} \right)^\epsilon \left( -\frac{2}{\epsilon^2} - \frac{3}{\epsilon} - 8 + \frac{7}{6} \pi^2 \right)$$

⊗

$$\frac{\int d^d x \frac{1}{e^2(p-l)^2}}{p} \rightarrow p-l \rightarrow p \quad p^2=0$$

$$\propto \int d^d x \frac{1}{e^2(p-l)^2} \sim [p^d]^{\frac{d-4}{2}} = 0 = \frac{1}{\epsilon_{UV}} - \frac{1}{\epsilon_{IR}}$$

dim. analysis

scaler

$$\int d^d x \frac{1}{Q^4} \sim UV \text{ div.}$$

$$\text{[Diagram: a loop with a wavy line and a straight line] } = -\frac{\alpha_s}{2\pi} \left( \frac{1}{\epsilon_{UV}} - \frac{1}{\epsilon_{IR}} \right)$$

UV IR

ε

Cancelled by Counter term

⊗ ⊗   $[d^4 l] \equiv \frac{d^4 l}{(2\pi)^4}$

$$= \int d^4 l \bar{u}_i i g_s t_{ik}^a \gamma_5 \frac{i \not{x+l}}{(q+l)^2} i e \not{\epsilon} (-i) \frac{\not{q-x}}{(q-l)^2} i g_s t_{kj}^a \gamma_5 \not{\epsilon} u_j = \frac{-i}{l^2}$$

$$= +i e g_s^2 (t^a t^a)_{ij} \int [d^4 l] \bar{u}_i \not{\epsilon} (x+l) \not{\epsilon} (x-l) \not{\epsilon} u_j \left[ \frac{1}{(q+l)^2} \frac{1}{(q-l)^2} \frac{1}{l^2} \right]$$

Now using the Feyn. Param. we find

$$\left[ \frac{1}{(q+l)^2} \frac{1}{(q-l)^2} \frac{1}{l^2} \right] = \Gamma(3) \int d\alpha_1 d\alpha_2 d\alpha_3 \frac{\delta(\alpha_1 + \alpha_2 + \alpha_3 - 1)}{(L^2 - \alpha_1 \alpha_3 S)^3}$$

$$\hookrightarrow \alpha_1 l^2 + \alpha_2 l^2 + 2\alpha_2 l \cdot q + \alpha_3 l^2 - 2\alpha_3 l \cdot \bar{q}$$

$$= l^2 + 2(\alpha_2 q - \alpha_3 \bar{q}) \cdot l + (\alpha_2 q - \alpha_3 \bar{q})^2 + \alpha_2 \alpha_3 S$$

$$= \underbrace{(l + \alpha_2 q - \alpha_3 \bar{q})^2}_{\equiv L}$$

We simplify the numerator

$$\begin{aligned}
 \square &= \bar{u}_i \delta^\alpha (\epsilon + \alpha_2 \not{x} + \alpha_3 \not{y}) \delta^\mu (\epsilon - \alpha_2 \not{x} - \alpha_3 \not{y}) \gamma_2 \psi_j \\
 &= -2\bar{u}_i (\epsilon - \alpha_2 \not{x} - \alpha_3 \not{y}) \delta^\mu (\epsilon + \alpha_2 \not{x} + \alpha_3 \not{y}) \psi_j \\
 &\quad + 2\epsilon \bar{u}_i (\epsilon + \alpha_2 \not{x} + \alpha_3 \not{y}) \delta^\mu (\epsilon - \alpha_2 \not{x} - \alpha_3 \not{y}) \psi_j
 \end{aligned}$$

$\propto L^{2n}$   
 $\xrightarrow{\text{odd power integrated to zero}}$

$$\begin{aligned}
 &-2\bar{u}_i \epsilon \delta^\mu \epsilon - \alpha_3 \alpha_2 \not{x} \delta^\mu \not{x} \psi_j \\
 &+ 2\epsilon \bar{u}_i \epsilon \delta^\mu \epsilon - \alpha_2 \alpha_3 \not{y} \delta^\mu \not{y} \psi_j
 \end{aligned}$$

$$= -2(1-\epsilon) \bar{u}_i \epsilon \delta^\mu \epsilon \psi_j$$

$$+ (2\alpha_3 \alpha_2 - 2\epsilon \alpha_3 \alpha_2) \bar{u}_i \not{x} \delta^\mu \not{x} \psi_j$$

$L^\mu L^\nu \rightarrow \frac{d}{d} L^\mu g^{\mu\nu}$   
 $\longrightarrow$

$$\begin{aligned}
 &\frac{(d-2)^2}{d} L^2 \bar{u}_i \delta^\mu \psi_j \\
 &+ 2(\alpha_2 \alpha_3 - \epsilon \alpha_2 \alpha_3) (-S) \bar{u}_i \delta^\mu \psi_j
 \end{aligned}$$

$$d = 2 - \epsilon$$

Now we use

$$\int [d\ell] \frac{L^2}{(L^2 + \alpha_2 \alpha_3 S)^3} = \frac{i}{(4\pi)^{d/2}} \frac{d}{2} \frac{\Gamma(\epsilon)}{\Gamma(3)} (\alpha_2 \alpha_3)^{-\epsilon} (-S)^{-\epsilon} \quad (1)$$

$$\int [d\ell] \frac{1}{(L^2 + \alpha_2 \alpha_3 S)^3} = \frac{-i}{(4\pi)^{d/2}} \frac{\Gamma(1+\epsilon)}{\Gamma(3)} (\alpha_2 \alpha_3)^{-\epsilon} (-S)^{-1-\epsilon} \quad (2)$$

⇒

① ⇒

$$g^2 \frac{i}{(4\pi)^{d/2}} \frac{d}{2} \frac{\Gamma(\epsilon)}{\Gamma(\epsilon)} (GS)^{-\epsilon} \frac{(d-2)^2}{d} \int_0^1 dx_2 \int_0^{\bar{x}_2} dx_3 (x_2 x_3)^{-\epsilon} \Gamma(\epsilon)$$

$$= \frac{\alpha_s}{2\pi} C_F \left(\frac{4\pi\mu^2}{-s}\right)^{\epsilon} \left[ \frac{1}{2\epsilon} + \frac{1}{2} \right]$$

UV cancelled by counter term


② ⇒

$$\frac{\alpha_s}{2\pi} C_F \left(\frac{4\pi\mu^2}{-s}\right)^{\epsilon} \left[ -\frac{1}{\epsilon^2} - \frac{2}{\epsilon} - \frac{9}{2} + \frac{\pi^2}{12} \right]$$

↳ Call IR

$$\cancel{\frac{1}{\epsilon^2}} + \cancel{\frac{2}{\epsilon}} + \cancel{\frac{9}{2}} + \cancel{\frac{\pi^2}{12}} + \cancel{\frac{1}{\epsilon^2}} + \cancel{\frac{2}{\epsilon}}$$

$$= \frac{\alpha_s}{2\pi} C_F \left(\frac{4\pi\mu^2}{-s}\right)^{\epsilon} \left[ -\frac{1}{\epsilon^2} - \frac{3}{2\epsilon} - 4 + \frac{\pi^2}{12} \right]$$



$$H^{(4)} = -C_F g_s^2 \delta(1-\epsilon) \frac{1}{d-1}$$

$$\times \left\{ \frac{2}{y_2 y_3} + \frac{-2 + y_3 - \epsilon y_3}{y_2} + \frac{-2 + y_2 - \epsilon y_2}{y_3} - 2\epsilon \right\}$$

Where

$$y_1 = \frac{S_{12}}{Q^2}, \quad y_2 = \frac{S_{13}}{Q^2}, \quad y_3 = \frac{S_{23}}{Q^2}$$

$$\text{and } S_{ij} = 2 p_i \cdot p_j$$

$$d\Phi_3 = \frac{1}{(2\pi)^{2d-3}} \frac{1}{2^{d+1}} S^{d-3} d\Omega_{d-1} d\Omega_{d-2}$$

$$(y_1 y_2 y_3)^{-\epsilon} \int_0^1 dy_1 dy_2 dy_3 \delta(1 - y_1 - y_2 - y_3)$$

$$\sigma_{\text{real}} = \sigma_0 \cdot \frac{d_s}{2\pi} C_F \frac{e^{\delta_{2\epsilon}}}{\Gamma(1-\epsilon)} \left(\frac{4\pi\mu^2}{s}\right)^{-\epsilon}$$

$$\times \int_0^1 dy_1 dy_2 dy_3 (y_1 y_2 y_3)^{-\epsilon} \delta(1-y_1-y_2-y_3)$$

$$\times \left\{ \frac{2}{y_2 y_3} + \frac{-2+y_3-\epsilon y_3}{y_2} + \frac{-2+y_2-\epsilon y_2}{y_3} - 2\epsilon \right\}$$

$$\times \Theta(\Phi_3)$$

Inclusive:  $\Theta(\Phi_3) = 1$

$$\text{let } y_1 = (1-z_1), y_2 = z_1 z_2, y_3 = (1-z_2) z_1$$

$$\sigma_{\text{real}} = \sigma_0 \cdot \frac{d_s}{2\pi} C_F \frac{e^{\delta_{2\epsilon}}}{\Gamma(1-\epsilon)} \left(\frac{4\pi\mu^2}{s}\right)^{-\epsilon}$$

$$\times \int_0^1 dz_1 dz_2 \frac{\bar{z}_1^{-\epsilon}}{z_1} \frac{z_1^{-2\epsilon}}{z_1} \frac{\bar{z}_2^{-\epsilon}}{z_2} \frac{\bar{z}_2^{-\epsilon}}{z_2} z_1 \quad \rightarrow \text{Jacobian}$$

$$\times \left\{ \frac{2}{z_1^2 \bar{z}_2^2} + \frac{-2+z_2\bar{z}_2-\epsilon z_2\bar{z}_2}{z_1 z_2} + \frac{-2+z_1\bar{z}_1-\epsilon z_1\bar{z}_1}{z_1 \bar{z}_2} - 2\epsilon \right\}$$

$$\Rightarrow \Delta_{\text{real}} = \Delta_0 \frac{\alpha_S}{2\pi} C_F \left( \frac{4\pi M^2}{s} \right)^{\epsilon} \left( \frac{2}{\epsilon^2} + \frac{3}{\epsilon} + \frac{19}{2} - \frac{7}{6}\pi^2 \right)$$

$$\Delta_{\text{virt.}} = \Delta_0 \frac{\alpha_S}{2\pi} C_F \left( \frac{4\pi M^2}{s} \right)^{\epsilon} \left( -\frac{2}{\epsilon^2} - \frac{3}{\epsilon} - 8 + \frac{7}{6}\pi^2 \right)$$

$$\Delta = \Delta_0 \left\{ 1 + \frac{\alpha_S}{2\pi} C_F \frac{3}{2} \right\}$$

• all IR poles cancelled, KLN Theorem

• Virt & Real almost completely cancelled with each other:  $V + R \sim 1 \neq \text{Unity}$

• Origin of the IR poles:

all IR poles from the soft & coll. limit

$$1. z_1 \rightarrow 0. \Rightarrow y_2 \neq y_3 \rightarrow 0$$

$$\Rightarrow p_3 \rightarrow 0 \quad \text{soft}$$

$$\cancel{p_4 \bar{E}_g \rightarrow 0}$$

$$2. z_2 \text{ or } \bar{z}_2 \rightarrow 0 \Rightarrow y_2 \text{ or } y_3 \rightarrow 0$$

$$\Rightarrow p_3 \cdot p_1 \rightarrow 0, p_3 \cdot p_1 \neq 0 \Rightarrow p_3 \parallel p_1$$

$$\text{or } p_3 \cdot p_2 \rightarrow 0, p_2 \cdot p_1 \neq 0 \Rightarrow p_3 \parallel p_2$$

$$z_1 \rightarrow 0$$

$$\int_0^1 d\alpha d\alpha' \int_0^1 dz_1 dz_2 \bar{z}_1^{-t} \bar{z}_2^{-t} \bar{z}_2^{-t} \times \left\{ \frac{2}{z_1 + i\epsilon z_2} \right\} = \frac{2}{\epsilon^2} + \dots$$

soft
collinear
soft + coll.

$$z_2 \rightarrow 0$$

$$\int_0^1 d\alpha d\alpha' \int_0^1 dz_1 dz_2 \bar{z}_1^{-t} z_1^{-t} \bar{z}_2^{-t}$$

$$\times \left\{ \frac{2}{z_1 + i\epsilon z_2} + \frac{-2 + t_1 - t_2 z_1}{z_2} \right\} = \frac{3}{2} \frac{1}{\epsilon}$$

Subtract out to avoid double counting

does reproduce all poles

every order you will have

$$\Delta_{\text{real}}^{(n)} \propto \left(\frac{\alpha_s}{2\pi}\right)^n \left( \frac{\#_{1-}}{\epsilon^{2n}} + \dots + \frac{\#_{1-}}{\epsilon} + \text{Finite terms} \right)$$

n-emission

n soft + coll.


cancel against the IR poles  
in the virtual. For inclusive  
processes

for exclusive processes,




e.g.  $pp \rightarrow \dots$   
 $e^- \rightarrow \text{hadron}^+$

remaining poles to be absorbed  
into PDFs or FFs.

thrust:  $T = \frac{\sum_i |\vec{p}_i \cdot \vec{n}|}{\sum_i |\vec{p}_i|}$




massless case  $\sum_i |\vec{p}_i| = Q$



measure the shape of an event

as  $T \rightarrow 0$ , only soft or coll. radiations are allowed



Used for discovery of gluon  
+ gluon spin determination  
+ precision determination of  $\alpha_s$

restrictions on the phase space


$$\Theta(\mathbb{I}_2) = \delta\left(T - \sum_{\max \vec{n}} \frac{|\vec{q}_1 \cdot \vec{n}| + |\vec{q}_2 \cdot \vec{n}|}{Q}\right)$$

$$= \delta\left(T - \sum_{\max \vec{n}} \frac{|\vec{q}_2 \cdot \vec{n}| + |\vec{q}_1 \cdot \vec{n}|}{Q}\right)$$

$$= \delta\left(T - \sum_{\max \vec{n}} \frac{Q/2 \cos \theta + Q/2 \cos \theta}{Q}\right)$$

$$= \delta\left(T - \sum_{\max \vec{n}} \cos \theta\right)$$

$$= \delta(1 - T) \quad \text{always.}$$



No restrictions

does not contribute to  $T \neq 1$  cases,

which starts from 3-body final


state

$$\delta^{(6)} + \delta_{\text{virt}}^{(1)} \rightarrow \left(\delta^{(6)} + \delta_{\text{virt}}^{(1)}\right) \delta(1 - T)$$

$$\Theta(\bar{\mathcal{E}}_3) = \mathcal{SCT} - \sum_{\max \vec{n}} \frac{(\vec{q}_1 \cdot \vec{n} + \vec{q}_2 \cdot \vec{n} + |\vec{q}_3 \cdot \vec{n}|)}{Q}$$

too complicated for analytic calculation

- for an arbitrary given  $\vec{n}$ , we can separate the phase space to 2 hemispheres, with one covers only one particle  $q_i$



- the max of  $\vec{n}$  for such config. is given by

$$\begin{aligned} & \max \left| \frac{\vec{q}_i \cdot \vec{n}}{Q} \right| + \left| \frac{\vec{q}_j \cdot \vec{n}}{Q} \right| + \left| \frac{\vec{q}_k \cdot \vec{n}}{Q} \right| \\ &= \max \left| \frac{\vec{q}_i \cdot \vec{n}}{Q} \right| + \left| \frac{\vec{q}_j \cdot \vec{n}}{Q} + \frac{\vec{q}_k \cdot \vec{n}}{Q} \right| \\ &= \max \left| \frac{\vec{q}_i \cdot \vec{n}}{Q} \right| + \left| -\frac{\vec{q}_i \cdot \vec{n}}{Q} \right| = \max \frac{2|\vec{q}_i \cdot \vec{n}|}{Q} = \frac{2E_i}{Q} = x_i \\ &= \frac{2q_i \cdot Q}{Q^2} = 1 - y_i \end{aligned}$$

- We need to maximize all possible configs

$$\Rightarrow T = \max(x_1, x_2, x_3) = \max(1 - y_1, 1 - y_2, 1 - y_3)$$

$$\Rightarrow 1 - T = \min(y_1, y_2, y_3) \equiv \tau$$

Which leads to the phase space restriction

$$\begin{aligned}\Theta(\Phi_3) &= \sum_i \delta(z - y_i) \Theta(y_i < y_j) \Theta(y_i < y_k) \\ &= \sum_i \delta(z - y_i) \Theta(z < y_k < 1 - 2z) \Theta(z < 1/3)\end{aligned}$$

Plug into the real-emission cross section


$$\begin{aligned}\sigma_{\text{real}} &= \sigma_0 \cdot \frac{\alpha_s}{2\pi} C_F \frac{e^{\gamma_{\text{E}} \epsilon}}{\Gamma(1-\epsilon)} \left(\frac{4\pi\mu^2}{s}\right)^{-\epsilon} \\ &\times \int_0^1 dy_1 dy_2 dy_3 (y_1 y_2 y_3)^{-\epsilon} \delta(1 - y_1 - y_2 - y_3) \\ &\times \left\{ \frac{2}{y_2 y_3} + \frac{-2 + y_3 - \epsilon y_3}{y_2} + \frac{-2 + y_2 - \epsilon y_2}{y_3} - 2\epsilon \right\} \\ &\Theta(\Phi_3)\end{aligned}$$

You can work out the integration with  $\epsilon$  kept using Mathematica

For simplicity, we set  $\epsilon=0$  to find

$$\frac{d\sigma_{\text{real}}}{d\tau} = \frac{\alpha_s G_F^2}{2\pi} \left\{ \frac{4}{\tau} \log\left(\frac{1-2\tau}{\tau}\right) - \frac{3(1+\tau)(1-3\tau)}{\tau} - 6 \log\left(\frac{1-2\tau}{\tau}\right) \right\}$$

for  $0 < \tau < \frac{1}{3}$ , otherwise vanishes.



We can slightly improve our result to cover the " $\epsilon=0$ " bin.

Since we know  $\sigma_{tot} = \sigma_0 \left(1 + \frac{\alpha_s}{2\pi} C_F \frac{3}{2}\right)$

hence  $\int_0^{\frac{1}{3}} \frac{d\sigma}{d\epsilon} = \sigma_{tot}$

We assume our real calculation is only "valid" for  $\epsilon > \delta$ , with  $\delta \rightarrow 0$ . Therefore

$$\frac{\sigma_{tot}}{\sigma_0} = \int_{\delta}^{\frac{1}{3}} \frac{d\sigma_{real}}{d\epsilon} + \int_0^{\delta} \frac{d\sigma_{virt}}{d\epsilon}$$

an effective virtual correction

$$= \frac{\alpha_s}{2\pi} C_F \left( \frac{5}{2} - \frac{\pi^2}{3} + 3 \log[\delta] + 2 \log^2[\delta] \right) + \Delta_{virt}$$

$$= 1 + \frac{3}{2} \frac{\alpha_s}{2\pi} C_F \quad * \quad \Delta_{real} + \Delta_{virt} \approx 1$$

this feature is useful for deriving Resummation

$$\Rightarrow \frac{\delta_{\text{virt}}}{\delta_0} = 1 + \frac{\alpha_s}{2\pi} C_F \left[ -1 + \frac{\pi^2}{3} - 3 \log[\delta] - 2 \log^2[\delta] \right]$$

Hence the  $\tau$ -distribution covers  $\tau > 0$  is given by

$$\frac{1}{\delta_0} \frac{d\delta}{d\tau} = \frac{1}{\delta_0} \left\{ \lim_{\delta \rightarrow 0} \frac{d\delta_{\text{real}}}{d\tau} \theta(\tau - \delta) + \frac{\delta_{\text{virt}}}{\delta} \theta(\delta - \tau) \right\}$$

$$= \lim_{\delta \rightarrow 0} \frac{1}{\delta_0} \frac{d\delta_{\text{real}}}{d\tau} \theta(\tau - \delta) - \frac{1}{\delta} \theta(\delta - \tau) \left( 3 \log[\delta] + 2 \log^2[\delta] \right) \frac{\alpha_s}{2\pi} C_F$$

$$\left[ 1 + \frac{\alpha_s}{2\pi} C_F \left( -1 + \frac{\pi^2}{3} \right) \right] \delta(\tau)$$

Now we introduce the "+"-prescription

$$\left( \frac{\log^n z}{z} \right)_+ = \lim_{\delta \rightarrow 0} \theta(\tau - \delta) \frac{\log^n z}{z} + \frac{1}{n+1} \log^n(\delta) \theta(\delta - \tau)$$

You can show that it is equivalent to what you are familiar with

$$\int_0^1 \left( \frac{\log^m z}{z} \right)_+ f(z) dz = \int_0^1 \frac{\log^m z}{z} (f(z) - f_0) dz$$

by noting that

$$\delta_{\text{real}} + \delta_{\text{virt}} = \delta_{\text{tot}}$$

$$\Rightarrow \delta_{\text{virt}} = \delta_{\text{tot}} \delta(1-z) - \int_0^1 \frac{d\delta_{\text{real}}}{dz} dz \delta(1-z)$$

$$= \delta_{\text{tot}} \delta(1-z) - \int_0^1 \frac{d\delta_{\text{real}}^{\text{neg.}}}{dz} \delta(1-z) - \int_0^1 \frac{d\delta_{\text{real}}^{\text{sig.}}}{dz} \delta(1-z)$$

finite as  $z \rightarrow 0$       div. as  $z \rightarrow 0$

combine with  $\frac{d\delta_{\text{real}}}{dz}$

$$\frac{d\delta_{\text{real}}}{dz} - \int_0^1 \frac{d\delta_{\text{real}}^{\text{sig.}}}{dz} \delta(1-z) \rightarrow \text{plus prescription}$$

eventually we find

$$\frac{1}{b_0} \frac{db}{dz} = \delta(z) + \frac{\alpha_s C_F}{2\pi} \int -\frac{4}{1-z} \left(\frac{\log z}{z}\right)_+ - 3(1+z)(1-3z) \left(\frac{1}{z}\right)_+ \\ + (-1 + \pi^2/3) \delta(z) \\ + \left. \frac{4}{1-z} \frac{\log(1-2z)}{z} - 6 \log\left(\frac{1-2z}{z}\right) \right\}$$

The result is equivalent to keep  $\varepsilon$  in real  
do the calc, Laurent expand in  $\varepsilon$ , + virt,

We can thus have the cumulant cross-section

$$\frac{\Delta(\tau < \delta)}{\Delta_0} = 1 + \frac{\alpha_S}{2\pi} C_F$$


$$\left\{ -2L^2 - 3L - 1 - 4\text{Li}_2(-1+2\delta) \right.$$

$$+ 6\delta \left[ \log\left(\frac{\delta}{1-2\delta}\right) + 1 \right] + \frac{9}{2}\delta^2 + 3\log(1-2\delta)$$

$$- 4 \left[ \log(1-2\delta) \log(1-\delta) - \log(1-\delta) \log \delta \right.$$

$$\left. \left. - \text{Li}_2(\delta) + \text{Li}_2(2\delta) \right] \right\}$$

for  $\delta \leq \frac{1}{3}$ ,  $L \equiv \log \delta$



⊙ in the " $\tau=0$ "-bin, there is an exact cancellation between real & virt. poles

$$\frac{1}{\sigma_0} \frac{d\sigma_{\text{virt}}}{d\tau} = \frac{\alpha_S}{2\pi} C_F \left(\frac{4\pi\mu^2}{s}\right)^{\epsilon} \left(-\frac{2}{\epsilon^2} - \frac{3}{\epsilon} - 8 + \frac{7}{6}\pi^2\right) \delta(1-\tau)$$

$$\frac{1}{\sigma_0} \frac{d\sigma_{\text{real}}}{d\tau} = \frac{\alpha_S}{2\pi} C_F \left(\frac{4\pi\mu^2}{s}\right)^{\epsilon} \left(\frac{2}{\epsilon^2} + \frac{3}{\epsilon} + \dots\right) \delta(1-\tau) + \dots$$


The cancellation ensures the predictive power of a F.O. calculation.

Not all observable has this feature !!)

recall that

- all poles come from the soft & coll. limit of the emitted partons,
- virt correction does not change multiplicities

⇒ to cancel poles, soft & coll. of real should fall into the same bin of virt.



IR safe observables:

$$\mathcal{O}_N(p_1, p_2, p_3, \dots, p_i, \dots, p_N) \xrightarrow{P_i \to 0} \mathcal{O}_{N-1}(p_1, p_2, p_3, \dots, p_{i-1}, p_{i+1}, \dots, p_N)$$

$$\mathcal{O}_N(p_1, p_2, p_3, \dots, p_i, \dots, p_j, \dots, p_N) \xrightarrow{P_i \parallel P_j} \mathcal{O}_{N-1}(p_1, p_2, \dots, p_i, p_j, \dots, p_{i+j}, \dots, p_N)$$

You can check that thrust is IR safe.

IRC unsafe:

e.g. particle numbers

$$\mathcal{O}_N(p_1 \dots p_N) = N \xrightarrow{p_i \rightarrow 0} N \neq \mathcal{O}_{N-1}(p_1 \dots p_N) = N-1$$

We need IRC safe observables to make  
F.O. predictive.

①  $L$  indicates the incomplete cancellation between real. & virt.

as  $\delta \rightarrow 1/3$ ,  $G^{(1)} \rightarrow \frac{\alpha_s}{2\pi} C_F^{3/2}$ , "Complete"


as  $\delta \rightarrow 0$ ,  $L \rightarrow \infty$ , recovers the divergence


↓  
recall only soft and coll. radiations are allowed



## ⊙ Break down of F.O.

The entire  $\frac{1}{\sigma} \frac{dB}{d\tau}$  spectrum does not agree with data, all due to large logs in the small  $\tau$  region






Now we focus on the  $\delta \ll 1$  region,


When  $\delta \ll 1$ , only soft & coll. radiations are allowed, and we have

$$\frac{\sigma(\delta)}{\sigma_0} = 1 - \frac{\alpha_S}{2\pi} C_F \left( \frac{4}{2} L^2 + 3 \cdot L + 1 - \frac{\pi^2}{3} \right)$$

recall the poles



$$\frac{\sigma_{\text{rad}}}{\sigma_0} \sim \frac{\alpha_S}{2\pi} C_F \left( \frac{2}{\epsilon^2} + \frac{3}{\epsilon} \dots \right)$$



soft & collinear nature.

Indeed, recall  $\ln = \int \frac{dx}{x}$

Hence, the logs in

$$\text{Real} = G_0 \cdot \frac{ds}{2\pi} \text{CF} \frac{e^{\delta_{\text{sc}}}}{\Gamma(1-\epsilon)} \left(\frac{4\pi\alpha'}{s}\right)^{-\epsilon}$$

$$\times \int_0^1 dy_1 dy_2 dy_3 (y_1 y_2 y_3)^{-\epsilon} \delta(1-y_1-y_2-y_3)$$

$$\times \left\{ \frac{2}{y_2 y_3} + \frac{-2+y_3-\epsilon y_3}{y_2} + \frac{-2+y_2-\epsilon y_2}{y_3} - 2\epsilon \right\}$$

$\Theta(\mathbb{E}_3)$   $\rightarrow$  double log. Single log.

Comes from  $y_2 \rightarrow 0, y_3 \rightarrow 0,$


but  $y_1 \rightarrow 0$  does not generate any

logs

Pictureially  $\circ$

$\gamma_2 \rightarrow$   + soft g  $\Rightarrow \tau = \gamma_2$

real  $\tau = \min(\gamma_i)$

$\gamma_3 \rightarrow$   + soft g  $\Rightarrow \tau = \gamma_3$

$\gamma_1 \rightarrow$   + soft quark  $\Rightarrow \tau = \gamma_1$

In this limit  $\circ$

$$\sigma_{\text{real}} = \sigma_0 \cdot \frac{\alpha_s}{2\pi} C_F \frac{e^{\gamma_{\text{soft}}}}{\Gamma(1-\epsilon)} \left(\frac{4\pi\mu^2}{s}\right)^{-\epsilon}$$

$$\times \int_0^1 dy_1 dy_2 dy_3 (y_1 y_2 y_3)^{-\epsilon} \cancel{\delta(1-y_1-y_2-y_3)}$$

$$\times \left\{ \frac{2}{y_2 y_3} + \frac{-2+\gamma_3-\epsilon\gamma_3}{y_2} + \frac{-2+\gamma_2-\epsilon\gamma_2}{y_3} - 2\epsilon \right\}$$

if keep only double log.

$\Theta(\Phi_3)$

$$\hookrightarrow \left[ \delta(\tau-\gamma_2) \Theta(\tau < \gamma_3) + \gamma_2 \leftrightarrow \gamma_3 + \text{log suppressed} \right]$$

$$G_{\text{IR}}^{\text{tree}} = -b_0 \cdot \frac{\alpha_s}{2\pi} C_F \frac{2 \times 2}{\epsilon} \log \epsilon + \log \text{Suppressed}$$

$$\rightarrow -b_0 \frac{\alpha_s}{2\pi} C_F \left( \frac{\log \epsilon}{2} \right)_+ \quad \text{by "Unitarity"}$$

cumulative  $\rightarrow -b_0 \frac{\alpha_s}{2\pi} C_F \cdot 2 \ln^2 \delta$  one power from soft  
one power from coll.

soft-collinear limits

reproduce the leading log structure

To all orders, we have structures

$$\Delta \sim 1$$

$$\text{NLO} \quad + \alpha_s L^2 \quad + \alpha_s L \quad + \#$$

$$\text{NNLO} \quad + \alpha_s^2 L^4 \quad + \alpha_s^2 L^3 \quad + \alpha_s^2 L^2 \quad + \alpha_s^2 L \quad + \#$$

$$\text{N}^3\text{LO} \quad + \alpha_s^3 L^6 \quad + \alpha_s^3 L^5 \quad + \alpha_s^3 L^4 \quad + \alpha_s^3 L^3 \quad + \alpha_s^3 L^2 \dots$$

$$\vdots \quad \quad \quad \vdots$$

When  $\alpha_s L^2 \sim 1$ , we can not truncate the series at fixed  $\alpha_s$

since the  $\alpha_s^n L^{2n}$  for  $n > 1$  are equally important. We need to resum  $\alpha_s L^2$  to all orders.

$$\Delta \sim 1$$

$$NLO \rightarrow \alpha_s L^2 + \alpha_s L + \#$$

$$NNLO \rightarrow \alpha_s^2 L^4 + \alpha_s^2 L^3 + \alpha_s^2 L^2 + \alpha_s^2 L + \#$$


$$N^3LO \rightarrow \alpha_s^3 L^6 + \alpha_s^3 L^5 + \alpha_s^3 L^4 + \alpha_s^3 L^3 + \alpha_s^3 L^2 + \dots$$

⋮  
⋮  
⋮

dominant  
logs, DL

Or equivalently.

We sum up the most singular behavior  
of the soft & coll. radiations to  
all orders



## There are different Approaches to Resummation

### • CSS Formalism

Collins, Soper, Sterman *Nucl. Phys. B* 250 (1985) 199-224  
hep-ph/0409313 (1929)

### • Coherent branching algorithm

Catani et al. *nucl. Phys. B* 327 (1989) 323-352  
*nucl. Phys. B* 407 (1993) 3-42

### • Soft collinear effective Theory

Bauer et al. hep-ph/0202088, hep-ph/0109045  
hep-ph/0107001, hep-ph/0011376  
Beneke et al. hep-ph/0206152


### • CAESAR

Banfi et al. hep-ph/0112156

All Approaches Now have the ability to go beyond NLL.

## • Infrared behaviour in QCD

→ a lesson from QED  $\text{gluon} \rightarrow \text{photon}$




photon phase space:

$$\frac{d^3 q}{2q(2\pi)^3} = \frac{q^2 dq d\Omega}{2q(2\pi)^3} = \boxed{\frac{1}{2(2\pi)^2} q dq d\cos\theta}$$

we are interested in terms  $\propto \frac{1}{q^2}$  in the matrix element square.

## • possible single emission


1. photon from internal line



$$\propto \frac{1}{(P_{ij} + q)^2} = \frac{1}{2P_{ij}q + P_{ij}^2} \xrightarrow{q \rightarrow 0} \frac{1}{P_{ij}^2}$$

regular as  $q \rightarrow 0$

## 2. photon from external line



$$\bar{u}(p_i) (-ie) \gamma^\mu \frac{p_i + q}{2p_i \cdot q} \dots$$


$$= \bar{u}(p_i) e \frac{\cancel{p_i} + \cancel{p_i}}{2p_i \cdot q}$$

No  $\cancel{q}$  enhancement

0, since  $\cancel{p_i} \cdot \cancel{p_i} = 0$

$$= \bar{u}(p_i) e \frac{p_i^\mu}{p_i \cdot q}$$


will give  $\cancel{q}$  enhancement we need



$$\bar{u}(p_i) (ie) \gamma^\mu \frac{-p_i}{2p_i \cdot q}$$

$$= \bar{u}(p_i) (-e) \frac{p_i^\mu}{p_i \cdot q}$$

absorb "-" sign into the electric charge.





$$- q^\mu \mathcal{M}_\mu = \sum_i e_i \frac{q \cdot p_i}{p_i \cdot q} = \sum_i e_i = 0$$

charge conservation, gauge invariance.

- factorized s-ft current  $J_i^\mu = e_i \frac{p_i^\mu}{p_i \cdot q}$   
independent of spin, Low theorem.

- a factorization of the long & short distance physics.

$$\frac{1}{q} \sim \lambda_q \gg \frac{1}{E} \sim \lambda_i$$



Can not resolve the internal structure

Q: How about radiate a soft fermion?

→ Soft leads to logs?

Now we square the Matrix element

$$\left| \text{Diagram} \right|^2 \rightarrow -\sum_{i \neq j} e_i e_j \frac{p_i \cdot p_j}{p_i \cdot q p_j \cdot q} \left| \mathcal{M}_0(\{p_i\}) \right|^2$$

for single emission

$$\begin{aligned} \text{define } d\omega(q) &= -d\Phi_1 \sum_{i \neq j} e_i e_j \frac{p_i \cdot p_j}{p_i \cdot q p_j \cdot q} \\ &= -\sum_{i \neq j} \frac{e_i e_j}{8\pi^2} \frac{p_i \cdot p_j}{p_i \cdot q p_j \cdot q} q dq d\omega d\theta \end{aligned}$$

$$\frac{1}{2(2\pi)^2} q dq d\omega d\theta$$

Single photon emission prob.

Suppose we do a measurement  $\nu$  on the photon. that imposes the phase space restriction  $\Theta_1(\nu)$  the prob. for the photon emission is

$$W_1(\nu)$$




$$= \int d\nu \left[ \underbrace{\frac{dW_1(\nu)}{d\nu}}_{\text{real}} \Theta_1(\nu) - \underbrace{\int d\nu' \frac{dW_1(\nu')}{d\nu'}}_{\text{virt.}} S(\nu) \right]$$

since  $\int d\nu (\text{virt.} + \text{real}) = 1 + \text{small corrections}$

$\Rightarrow$

$$W_1(\nu) = \int d\nu' [\Theta_1(\nu') - 1] \frac{dW_1}{d\nu'}$$

Multiple emissions :



$$= \frac{1}{n!} \sum_{\text{All permutations}} e_i \frac{p^m}{p \cdot q_1} e_i \frac{p^m}{p \cdot (q_1 + q_2)} \dots e_i \frac{p^m}{p \cdot (q_1 + \dots + q_n)}$$

$$= \frac{1}{n!} \prod_{j=1}^n e_i \frac{p^m}{p \cdot q_j} \rightarrow dW_n = \frac{1}{n!} \prod_{j=1}^n dW_j$$

"independent" emission

$$\Rightarrow \sum_n \frac{1}{n!} \prod_i \int d\omega_i |M(\{p_i\})|^2$$


The diagram shows a central circle representing a particle with several lines radiating outwards, representing emitted photons. A bracket on the right side of the lines is labeled 'n photons'. An arrow points from this diagram to the mathematical expression on the right.

- We note that the emission probability also depends on the phase space restriction which is not necessarily independent

$$\Theta_n(\omega) = \Theta_n(V(\{q\}))$$

- if the phase space restriction is also independent then

$$\Theta \rightarrow \Theta_0 \exp[W_1(\omega)]$$

where

$\omega_1$  defined before including the "virtual".

$$W_1(\omega) = \int dV \{ \Theta_1(\omega) - 1 \} \frac{dW_1(q)}{dV}$$

you get the resummed form in QED for  $\sigma$ .

\*  $\Theta_c(\infty) = 1 \Rightarrow$  Inclusive  $\Rightarrow W_c(\infty) = 0$   
 $\Rightarrow \sigma \rightarrow \sigma_0$ , Unitarity,

\*  $\Theta_c(\infty) = 0 \Rightarrow$  No emission

$\Rightarrow$

Sudakov factor

$$\delta \rightarrow \sigma_0 \exp \left[ - \int_{\delta}^{\infty} dv \frac{dw(v)}{dv} \right] \rightarrow 0$$

$\int_{-\infty}^{\infty}$  virtual only

$\Delta(\Theta_{\max}, 0)$

prob. of no emission below  $\Theta_{\max}$

\* 2 photons,

$$\frac{P_i^h}{P_i \cdot \xi_i} \frac{P_i^a}{P_i \cdot (\xi_i + \eta_i)} + \frac{P_i^h}{P_i \cdot \eta_i} \frac{P_i^m}{P_i \cdot (\xi_i + \eta_i)}$$

$$= P_0^m P_1^r \frac{P_0^r q_2 + P_1^r q_1}{P_0^r q_1 P_1^r q_2 P_2^r (q_1 + q_2)} = \frac{P_0^m}{P_1^r q_1} \frac{P_1^m}{P_2^r q_2} \checkmark$$

$$\text{suppose } \sum_{\text{perm.}} \frac{P_0^r}{P_1^r q_1} \frac{P_1^r}{P_2^r (q_1 + q_2)} + \dots + \frac{P_0^m}{P_2^r (q_1 + q_2)}$$

$$= \frac{P_0^r}{P_1^r q_1} \dots \frac{P_1^m}{P_2^r q_n}$$

$$\text{then } \sum_{\text{perm.}} \frac{P_0^r}{P_1^r q_1} \dots \frac{P_1^r}{P_2^r (q_1 + q_2)} \frac{P_2^r}{P_3^r (q_1 + q_2)}$$

$$= \sum_{\text{perm. } j \neq n} \frac{P_1^r}{P_2^r q_j} \dots \frac{P_1^m}{P_2^r (q_1 + q_2)}$$

$$+ \sum_{\text{perm. } j \neq 1} \frac{P_1^m}{P_2^r q_2} \dots \frac{P_1^r}{P_2^r (q_1 + q_2)} \frac{P_2^r}{P_3^r (q_1 + q_2)}$$


⊥ ...

$$= \left\{ \prod_{j \neq 1} \frac{P_1^m}{P_2^r q_j} + \prod_{j \neq 2} \frac{P_1^r}{P_2^r q_j} + \dots + \prod_{j \neq n} \frac{P_1^r}{P_2^r q_j} \right\} \frac{P_0^m}{P_2^r (q_1 + q_2)}$$

$$= \prod_{j \neq 1} \left( \frac{P_1^m}{P_2^r q_j} (P_2^r q_1 + P_2^r q_2 + \dots + P_2^r q_n) \right) \frac{P_0^m}{P_2^r (q_1 + q_2)}$$

$$= \prod_{j \neq 1} \frac{P_1^m}{P_2^r q_j}$$

## QED coherence



suppose a  $\theta_{pk}$  is the smallest among all angles.

$$= - \sum_{i \neq j} e_i e_j \frac{2 P_i \cdot P_j}{P_i \cdot q P_j \cdot q}$$

$$= -e_k \sum_{j \neq k} e_j \frac{2 P_k \cdot P_j}{E_k E_j (1 - \cos \theta_{kj})} \frac{1}{E_j E_j (1 - \cos \theta_{kj})} + \text{less singular}$$

$$= -\frac{e_k}{E_q} \frac{2}{1 - \cos \theta_{kj}} \sum_{j \neq k} e_j \frac{P_k \cdot P_j}{E_j E_k (1 - \cos \theta_{kj})} + \text{less singular}$$

$$\approx -\frac{2 e_k}{E_q^2} \frac{1}{1 - \cos \theta_{kj}} \sum_{j \neq k} e_j \frac{E_k E_j (1 - \cos \theta_{kj})}{E_k E_j (1 - \cos \theta_{kj})} + \text{less singular}$$

~~$e_k$~~  Since charge cons.

$$\approx \frac{2e_K^2}{E_q^2} \frac{1}{1 - \cos \theta_{qk}} \quad (\otimes)$$

Here we used the coll. limit to derive the result but it can be derived as long as  $\theta_{qk}$  is the smallest angle, no need for  $\theta_{qk} \rightarrow 0$ . See below.

Now we integrate over the photon phase space to find


$$d\omega_i(q) = \frac{\alpha}{\pi} \underbrace{\sum_k e_k^2 \frac{dq}{q}}_{\text{from a single independent emitter}} \underbrace{\frac{d\theta_{qk}^2}{\theta_{qk}^2} \Theta(\theta_{\max} - \theta_{qk})}_{\text{QED coherence due to destructive interference}}$$

from a single independent emitter

QED coherence due to destructive interference

Here  $\theta_{\max} = \min\{\theta_{ij}\}$





$\Rightarrow$  see the overall charge  
 does not carry charge  $\rightarrow$  hence the wide angle  
 emission is suppressed

$$(-e) \frac{p_j^m}{p_j \cdot r} + e \frac{p_i^m}{p_i \cdot r} = (-e + e) \frac{p^m}{p \cdot r} = 0$$

Therefore we find

$$\sigma = \sigma_0 \exp \left[ \int dV (\theta_i(\omega) - 1) \frac{dW_i}{dV} \right]$$


$$= \sigma_0 \prod_k \exp \left\{ \frac{\alpha}{\pi} e_k^2 \int \frac{dq}{q} \int_{\theta_{qk}^{\min}}^{\theta_{qk}^{\max}} [\theta_i(\omega) - 1] \right\}$$

Shows each leg

$$= \sigma_0 \prod_k \sum_{n=0}^{\infty} \frac{1}{n!} \prod_n \frac{\alpha^n}{\pi^n} \left( e_k^2 \int \frac{dq}{q} [\theta_i(\omega) - 1] \right)^n \int_{\theta_1^{\min}}^{\theta_1^{\max}} \int_{\theta_2^{\min}}^{\theta_2^{\max}} \dots \int_{\theta_n^{\min}}^{\theta_n^{\max}}$$

$$= \sigma_0 \prod_k \sum_{n=0}^{\infty} \prod_n \frac{\alpha^n}{\pi^n} \left( e_k^2 \int \frac{dq}{q} [\theta_i(\omega) - 1] \right)^n \int_{\theta_n^{\min}}^{\theta_n^{\max}} \int_{\theta_{n-1}^{\min}}^{\theta_{n-1}^{\max}} \int_{\theta_{n-2}^{\min}}^{\theta_{n-2}^{\max}} \dots \int_{\theta_1^{\min}}^{\theta_1^{\max}}$$

angular order



angular ordering to  
account for the interference

Very useful  
for Monte Carlo  
event-generator

(\*)

$$\frac{P_i \cdot P_j}{P_i \cdot q \cdot P_j \cdot q} = S_{ij}^{(1)} + S_{ij}^{(2)}$$

$$S_{ij}^{(1)} = \frac{1}{2} \left[ \frac{P_i \cdot P_j}{P_i \cdot q \cdot P_j \cdot q} + \frac{1}{c} \frac{1}{1 - \cos \theta_{iq}} - \frac{1}{c^2} \frac{1}{1 - \cos \theta_{jq}} \right]$$

where

$$E_i \cdot c \left( \sin \theta_{iq} \cos \phi_{iq} \sin \theta_{iq} \sin \phi_{iq} \cos \theta_{iq} \right)$$

$$E_j \cdot c \left( \sin \theta_{ij} \quad , \quad 0 \quad , \quad \cos \theta_{ij} \right)$$



$$1 - \cos \theta_{jq} = \underbrace{1 - \cos \theta_{iq} \cos \theta_{ij}}_c - \underbrace{\sin \theta_{iq} \sin \theta_{ij} \cos \phi_{iq}}_s$$

Therefore

$$\int_0^{2\pi} \frac{d\phi_{iq}}{2\pi} \frac{1}{1 - \cos \theta_{jq}} = \int_0^{2\pi} \frac{d\phi_{iq}}{2\pi} \frac{1}{c - s \cos \phi_{iq}} \quad \text{let } z = e^{i\phi_{iq}}$$

$$= \frac{-i}{2\pi} \oint_{|z|=1} \frac{dz}{z} \frac{1}{c - s(z + 1/z)} = \frac{i}{2\pi} \oint_{|z|=1} dz \frac{1}{sz^2 - cz + s}$$

$$= \frac{i}{2\pi s} \oint_{|z|=1} dz \frac{1}{(z - z_+)(z - z_-)} \quad \text{where } z_{\pm} = \frac{1}{2} \frac{c}{s} \pm \frac{1}{2} \sqrt{\frac{c^2}{s^2} - 1}$$



$$\Rightarrow \text{result} = -\frac{1}{s} \frac{1}{z-z_+}$$

$$= \frac{1}{\sqrt{c^2-s^2}}$$

$$\Rightarrow \text{result} = 0$$

Therefore

$$\int \frac{d\theta_{ij}}{2\pi} S_{ij}^{(c)} = \frac{1}{2} \frac{1}{c^2} \int \frac{d\theta_{ij}}{2\pi} \left\{ \frac{1-\cos\theta_{ij}}{(1-\cos\theta_{ij})(1-\cos\theta_{iq})} + \frac{1}{1-\cos\theta_{iq}} - \frac{1}{1-\cos\theta_{ij}} \right\}$$

$$= \frac{1}{2} \frac{1}{c^2} \left\{ \left[ \frac{1-\cos\theta_{ij}}{1-\cos\theta_{iq}} - 1 \right] \frac{1}{\sqrt{c^2-s^2}} + \frac{1}{1-\cos\theta_{iq}} \right\}$$

$$= \frac{1}{2c^2} \left\{ \frac{\cos\theta_{iq} - \cos\theta_{ij}}{1-\cos\theta_{iq}} \frac{1}{|\cos\theta_{iq} - \cos\theta_{ij}|} + \frac{1}{1-\cos\theta_{iq}} \right\}$$

$$= \begin{cases} \frac{1}{2c^2} \frac{1}{1-\cos\theta_{iq}} & \theta_{iq} < \theta_{ij} \\ 0 & \theta_{iq} > \theta_{ij} \end{cases}$$

$z_{i,0}$

$\frac{z_{i,0}^2}{z_{i,0}}$

$z_{i,0} \sim z_{i,0} \frac{1}{z_{i,0}}$   
 $\sigma_b = \frac{\theta_{ij}}{1-\cos\theta_{ij}} \gg 1 \sim \frac{1}{z_{i,0}}$

$\theta > \theta$   
 $\theta > \theta$

① summarize for  $\Delta E D$

$$- W_1 = -4\pi d \sum_{\vec{e}_j} e_j \int \frac{p_i \cdot \vec{e}_j}{r_{ij}^2} (\Theta_i(\nu) - 1) d\nu d\Omega_j$$

$\nu$   
virtual included

= if phase space factorize, then

$$\delta G = \delta G_0 \exp[W_1]$$

- Interference well approximated by time ordering

$$W_1 \rightarrow \frac{d}{dt} \sum_{\vec{k}} e_k^2 \int \frac{d^3 q}{q} \frac{d\omega}{\omega^2} \Theta(\omega_{\max} - \omega_{qk}) (\Theta_i(\nu) - 1) d\nu$$

and

$$\delta G = \delta G_0 \exp[W_1]$$

$$= \delta G_0 \prod_{k=1}^n \prod_n \left[ \frac{d}{dt} \exp \left[ \int \frac{d^3 q}{q} (\Theta_i(\nu) - 1) d\nu \right] \right]^n \underbrace{\int_{\omega_{\min}}^{\omega_{\max}} \frac{d\omega_n}{\omega_n} \int_{\omega_{\min}}^{\omega_n} \frac{d\omega_{n-1}}{\omega_{n-1}} \dots \frac{d\omega_1}{\omega_1}}_{\text{angular ordering}}$$

## → soft emissions in QCD.

### • single emission

Similar to the QED soft photon emission, we have the soft gluon radiated off a quark line.


$$\text{quark line} \rightarrow \text{gluon } (\mu, a, q) = g_s t_{ij}^a \frac{p_i^\mu}{p_i \cdot q} \mathcal{M}_0^{j, \dots}$$

↳ with  $e_i$  replaced by the color charge,  $g_s t^a$ , a matrix.


$$\text{anti-quark line} \rightarrow \text{gluon } (\mu, a, q) = g_s (-t_{ij}^a) \frac{p_i^\mu}{p_i \cdot q} \mathcal{M}_0^{j, \dots}$$

↳ From anti-quark

More than that, since now the gluon carries charge



soft gluon can be emitted from hard gluons.



$$\frac{g_s^2}{(P_i + q_1 + q_2)^2} \sim \frac{g_s^2}{2P_i(q_1 + q_2) + 2q_1 \cdot q_2} \sim \frac{g_s^2}{2P_i \cdot (q_1 + q_2)}$$


$\underbrace{\hspace{10em}}_S \quad \underbrace{\hspace{10em}}_{E_1 + E_2} \quad \underbrace{\hspace{10em}}_{\cancel{E_1 + E_2}} \quad \underbrace{\hspace{10em}}_{\text{not singular}}$

Since from phase space we have

$q_1 dq_1 q_2 dq_2$ , which returns

suppressed results for either

$q_1 \rightarrow 0$  or  $q_2 \rightarrow 0$



$$\begin{aligned}
 &\propto \frac{1}{2p_i \cdot q} g_s f^{abc} \left\{ \overset{\text{gauge}}{\cancel{+2(p_i+q)^\mu}} g^{\mu\nu} - \underset{\substack{\text{non-singular} \\ \downarrow q}}{(p_i+q)^\mu} g^{\nu\mu} + \overset{\text{gauge}}{\cancel{+ (p_i-q)^\nu}} g^{\mu\nu} \right\} \epsilon_\nu^c \ell^\mu \\
 &= -\frac{1}{2p_i \cdot q} g_s f^{abc} 2p_i^\mu \epsilon_\nu^c \ell^\mu \\
 &= g_s f^{abc} \frac{p_i^\mu}{p_i \cdot q} \ell^\mu
 \end{aligned}$$

C.F. shown from quark:  $g_s t_{ij}^a \frac{p_i^\mu}{p_i \cdot q}$

- same eikonal factor, spin-independent
- with the color charge  $g_s F^{abc} \equiv g_s f_{cb}^a$

We can Unify the soft approximation for both quark and gluon emission by introducing the color charge  $\vec{T}_i^a$  where  $i$  is the  $i$ -th parton

$$\vec{T}_i^a \rightarrow t^a \text{ if } i \text{ is a quark } (\neq)$$

$$\rightarrow \bar{t}^a \text{ if } i \text{ is an anti-quark}$$

$$\rightarrow f_{cb}^a \text{ if } i \text{ is a gluon.}$$

Color algebra: 

We note that  $\sum_i \vec{T}_i^a = 0$  color charge conservation

cf.  $\sum_i e_i = 0$  electric charge conservation

and if we define  $\vec{T}_i^2 = \sum_a \vec{T}_i^a \vec{T}_i^a$ , then

$$\vec{T}_i^2 = \begin{cases} CF & \text{For quark } \neq \text{ anti-quark} \\ CA & \text{For gluon.} \end{cases}$$

⊗

$$= g_s t_{c_i c_i'}^a \mathcal{M}^{\dots c_i \dots}$$

Color operator acting on the color space, like a rotation

$$\equiv g_s t_{c_i c_i'}^a |\dots c_i \dots\rangle$$

$$= g_s T_c^a |\dots c_i \dots\rangle$$





multiple emissions

recall QED :

$$d\sigma \sim d\sigma_0 \exp[W_1]$$

$$W_1(\nu) = -4\pi^2 \sum_{i \neq j} e_i e_j \int \frac{p_i \cdot p_j}{p_i \cdot q p_j \cdot q} (Q_1(\omega) - 1) d\nu d\Phi_q$$

direct exponentiation of 1-photo emission. (if the observable  $\nu$  is factorizable), since in QED,

$$|M|^2 \sim |M_0|^2 e^{\frac{p_i \cdot p_j}{p_i \cdot q p_j \cdot q}} \text{ entirely factorizable,}$$

How about QCD ?

Can we have


$$d\sigma \sim d\sigma_0 \exp[W_1]$$

$$W_1(\nu) = \int -4\pi^2 \sum_{i \neq j} \vec{t}_i \cdot \vec{t}_j \int \frac{p_i \cdot p_j}{p_i \cdot q p_j \cdot q} (Q_1(\omega) - 1) d\nu d\Phi_q \quad ?$$

No. **cannot factorize!!**

- color coherence

similar to QED, the interference can be simplified in the leading IR limit by angular ordering



$$= -g_s^2 \frac{p_i \cdot p_j}{p_i \cdot q p_j \cdot q} \langle M_0 | \epsilon \cdot T_j | M_0 \rangle$$

$$\approx g_s^2 \sum_i T_i^2 \frac{2}{E^2} \frac{1}{\theta_{iq}^2} \Theta(\theta_{max} - \theta_{iq}) |M|^2$$

$\left\{ \begin{array}{l} \text{diagonal in the color space, C-A} \\ \text{does not "change" color} \end{array} \right.$


- QED-like factorization

- angular ordering for interference

$$- T_i^2 = \begin{cases} CF & \text{for } q \\ CA & \text{for } g \end{cases}$$

small angle radiation does not "change" the color charge.






$$T_i \frac{P^m}{P \cdot q} + T_j \frac{P^m}{P \cdot q} = \{T_i + T_j\} \frac{P^m}{P \cdot q} = T_{ij} \frac{P^m}{P \cdot q}$$

as if radiated from the hard emitter  $\tilde{q}_j$

wide angle soft radiation only sees  
the overall color charge of the partons  
at smaller angles



QCD multiple  
emission picture  $\Rightarrow$   
coherent branching



2CD

$$W_k(v) = 1 + \frac{1}{n_i} \prod_{n=1}^{\infty} \left[ \frac{\alpha_s}{\pi} T_k^2 \frac{dE_g}{E_g} \int_{\theta_{qk}^2}^{\theta_{max}} \left\{ \theta_k(v) W_q(v) - 1 \right\} \right]^n$$


$$= \exp \left[ \frac{\alpha_s}{\pi} T_k^2 \int \frac{dE_g}{E_g} \int_{\theta_{qk}^2}^{\theta_{max}} \left\{ \theta_k(v) W_q(v) - 1 \right\} \right]$$

- the soft gluon as the new emitter.
- small angle  $\theta_{max} \sim \theta_{qk}$  now

$$W(v) = \prod_k W_k(v), \quad \delta = \delta_0 W(v)$$

- the fundament to parton shower

e.g. Prob. for one-emission



$$W_c(v_{c0}, \theta_j, \theta_c) \frac{\alpha_s}{\pi} \frac{2 dE_g}{E_g} \frac{d\theta_g^2}{d\theta_g^2} W_g(v_{g0}, \theta_g, 0) W_c(v_{c0}, \theta_j, 0)$$

Sudakov  $\textcircled{1}$ 
1-emission prob.  $\textcircled{2}$   $\textcircled{3}$

1-emission generated using Markov chain

- can be improved to sub-leading logs

$$\frac{\alpha_s}{\pi} \frac{2 dE}{E} \frac{d\theta^2}{\theta^2} \rightarrow \frac{\alpha_s}{2\pi} \frac{d\theta^2}{\theta^2} P_{\text{ang}}(z) dz$$

(splitting function)

- applied to analytic Resummation

## DL resummation for the thrust.



we have two branchings, and we consider  $y_2 \rightarrow 0$ .  $y_3 \rightarrow 0$  can be included in the same way

We first consider the 1-gluon emission

Recall that for the thrust  $z \ll 1$

$$\Theta_1(\nu) = \Theta(s-z) \delta(\tau - y_2) \Theta(y_3 - y_2) + 2 \leftrightarrow 3$$

↓

+  ~~$1 \leftrightarrow 2$~~   $\rightarrow 0 \rightarrow$  does not lead to LL.

$$\approx \Theta\left(s - \underbrace{\frac{\sqrt{E_1 E_3}}{4E^2} \frac{\theta^2}{2}}_z\right) \Theta\left(\frac{4EE_3}{4E^2} - \tau\right)$$

$$= \Theta(s - z) \Theta\left(\frac{E_3}{E} - \tau\right)$$

Therefore we have for one-gluon case:

$$\frac{dE_3}{E_3} \frac{d\theta_3^2}{\theta_3^2} \Theta_1(\nu) = \int_0^1 \frac{d\tau}{\tau} \int_0^1 \frac{dz}{z} \Theta(s - z)$$

For  $n$ -emissions we have

$$\Theta_n(V) = \Theta(\delta - \sum_0^n \tau_i) \prod_0^n \delta(\tau_i - \tau_w) \Theta(y_{i1} - y_{i2})$$

$$= \Theta(\delta - \sum_0^n \tau_i) \prod_0^n (\tau_i - \tau)$$

Does not factorize! Factorized

Hence we have for  $n$ -emission

$$\frac{dE_i}{E_i} \frac{d\theta_i^2}{\theta_i^2} \Theta_n(V)$$

$$= \left\{ \prod_0^n \int_0^1 \frac{d\tau_i}{\tau_i} \int_{\tau_i}^1 \frac{dz_i}{z_i} \right\} \Theta(\delta - \sum_0^n \tau_i)$$

To factorize  $\Theta(\delta - \sum_0^n \tau_i)$ , we note

that

$$\Theta(\delta - \sum_0^n \tau_i) = \frac{1}{2\pi i} \int \frac{dv}{v} e^{v\delta} e^{-\sum_0^n \tau_i v}$$

to find

$$\frac{d\bar{E}_0}{E_0} \frac{d\theta_0^2}{\theta_0^2} \Theta_n(\nu)$$

$$= \frac{1}{2\pi i} \int \frac{d\nu}{\nu} e^{\nu S} \prod_{\tilde{c}=1}^n \int \frac{d\tau_{\tilde{c}}}{\tau_{\tilde{c}}} \int_{\tau_{\tilde{c}}}^1 \frac{dz_{\tilde{c}}}{z_{\tilde{c}}} e^{-\tau_{\tilde{c}} \nu}$$

$\int \Theta_1(\nu)$

Now we plug it into  $W(\nu)$


$$W(S) = \frac{1}{2\pi i} \int \frac{d\nu}{\nu} e^{\nu S} \exp \left[ 2 \int \frac{d\tau}{\tau} \int_{\tau}^1 \frac{dz}{z} \frac{d\tau}{\tau} G_F \left\{ e^{-\tau \nu} - 1 \right\} \right]$$

$e^{-\tau \nu}$  virtual

We further use the fact that

$$e^{-\tau \nu} - 1 \simeq -\Theta(\tau - e^{-\delta E \nu^{-1}})$$

which holds up to NLL.



If we ignore the  $\tau$  &  $z$  dependence in  $\alpha_s$  which is sub-leading, we can have

$$W(s) = \frac{1}{2\pi i} \int \frac{dv}{v} e^{vs} e^{-\frac{\alpha_s}{\pi} C_F (\chi_E + \log v)^2}$$

Now we let  $u = vs$  to have


$$\begin{aligned} W(s) &= \frac{1}{2\pi i} \int du e^u e^{-\log u} e^{-\frac{\alpha_s}{\pi} C_F (\log u + \log e^{\frac{u}{s}})^2} \\ &= e^{-\frac{\alpha_s}{\pi} C_F \log^2 e^{\frac{u}{s}}} \underbrace{\frac{1}{2\pi i} \int du e^u e^{-\log u} e^{-\frac{\alpha_s}{\pi} C_F (\log u + \dots)^2}}_{\mathcal{O}(u)} \end{aligned}$$

sub-leading in logs

$$\Rightarrow \frac{\mathcal{O}(s)}{\mathcal{O}(t)} = e^{-\frac{\alpha_s}{\pi} C_F \log^2 s} + \text{sub-leading terms} \dots$$

$$= 1 - \frac{\alpha_s C_F}{\pi} \log^2 s + \underbrace{\left[ \frac{\alpha_s}{\pi} \right]^2 C_F^2 \log^4 s + \dots}_{\text{predict higher orders}}$$

↓ reproduce the full NLO



Comment on the results:

- Can be improved to NLL by using the coherent branching to derive a RGE of the "jet" function

$$\Rightarrow DL \approx \exp[\text{NLO double log}]$$

$$DL = \exp \left[ \frac{2\alpha_s}{\pi} C_F \int_0^1 \frac{d\theta^2}{\theta^2} \int_{\tau}^1 \frac{dz}{z} \left\{ \Theta(s-z) - 1 \right\} \right]$$

$$\approx 1 + \frac{\alpha_s}{n\pi} \left[ \frac{2\alpha_s}{\pi} C_F \int_0^1 \frac{d\theta^2}{\theta^2} \int_{\tau}^1 \frac{dz}{z} \left\{ \Theta(s-z) - 1 \right\} \right]^n$$

$$\text{with } \tau = \frac{z\theta^2}{4}$$

Therefore at DL, we can replace

$$\Theta(s - \sum_{i=1}^n \tau_i) \rightarrow \prod_{i=1}^n \Theta(s - \tau_i)$$

The emissions are all independent!

This holds in general for many observables

