



中国科学院高能物理研究所
Institute of High Energy Physics Chinese Academy of Sciences

Probing Higgs CP properties at the CEPC

Qiyu Sha , Abdualazem Fadol, Fangyi Guo, Jiayin Gu, Gang Li, Xinchou Lou, Yaquan Fang

Institute of High Energy Physics CAS

Fudan University

HIGGS POTENTIAL 2022

Content

Introduction

Theory framework

Optimal variable

Monte Carlo samples

Event selection

Fitting strategy and results

Summary

[arXiv:2203.11707](https://arxiv.org/abs/2203.11707)

1 Probing Higgs CP properties at the CEPC

2 Qiyu sha,^{*} Abdualazem Fadol,[†] Fangyi Guo,[‡] Gang Li,[§] and Yaquan Fang,[¶]

3 *Institute of High Energy Physics, 19B, Yuquan Road,*

4 *Shijingshui District, Beijing, China, 100049 and*

5 *University of Chinese Academy of Sciences (CAS)*

6 Jiayin Gu^{**}

7 *Department of Physics and Center for Field Theory and Particle Physics,*

8 *Fudan University, Shanghai 200438, China and*

9 *Key Laboratory of Nuclear Physics and Ion-beam Application (MOE),*

10 *Fudan University, Shanghai 200433, China*

11 Xinchou Lou^{††}

12 *Institute of High Energy Physics, 19B, Yuquan Road,*

13 *Shijingshui District, Beijing, China, 100049*

14 *University of Chinese Academy of Sciences (CAS) and*

15 *University of Texas at Dallas, Richardson, Texas 75080-3021, USA*

16 Abstract

17 In the Circular Electron Positron Collider (CEPC), a measurement of the Higgs CP mixing
18 through $e^+e^- \rightarrow ZH \rightarrow l^+l^-(e^+e^-/\mu^+\mu^-)H \rightarrow b\bar{b}/c\bar{c}/gg$ process is presented, with 5.6 ab^{-1}
19 e^+e^- collision data at the center-of-mass energy of 240 GeV. In this study, the CP -violating
20 parameter $\tilde{c}_{Z\gamma}$ is constrained between the region of -0.30 and 0.27 and \tilde{c}_{ZZ} between -0.06 and
21 0.06 at 68% confidence level. This study demonstrates the great potential of probing Higgs CP
22 properties at the CEPC.

23 Keywords: the Higgs Boson, CP violation, CEPC

^{*} shaqiyu@ihep.ac.cn

[†] amohammed@alims.ac.tz

[‡] guofangyi@ihep.ac.cn

[§] li.gang@ihep.ac.cn

[¶] fangyq@ihep.ac.cn

^{**} jiayin_gu@fudan.edu.cn

^{††} xinchou@ihep.ac.cn

Introduction

Properties of Higgs in Standard Model: $m_H = 125.10\text{GeV}$, $J^{PC} = 0^{++}$

Related experiments in LHC:

- The hypothesis of spin-1 or spin-2 Higgs has been excluded by the ATLAS and CMS at >99% CL in $\sqrt{s} = 7\&8 \text{ TeV}$, 25 fb^{-1} data. [Eur. Phys. J. C75 \(2015\) 476](#)
- The results of the study on the CP properties of the Higgs boson interactions with gauge bosons by the ATLAS and CMS show no deviations from the SM predictions.

Higgs-gauge vector boson interaction lacks precise measurement in all inclusive Higgs production mode(i.e. ggF dominant).


Any observation of CP violation in Higgs would be New Physics!

What we want to study is the Higgs CP mixing model aiming to find the CP-odd Higgs.

Introduction

Future e^+e^- collider experiment as Higgs factory :


- At a center of mass energy of $\sqrt{s} \sim 240\text{GeV}$ which maximizes the Higgs boson production cross section through $e^+e^- \rightarrow ZH$ process.
- Cleaner environment and more events produced than (HL)-LHC.
- More precise Higgs-gauge boson coupling study.





CEPC, 5.6 ab^{-1} @ 240GeV



FCC-ee, 5 ab^{-1} @240GeV



ILC, 2 ab^{-1} @250Gev



Theory framework

JHEP 03(2016) 050 JHEP 11(2014) 028

Consider a 6-dimension EFT model: $\mathcal{L}_{eff} = \mathcal{L}_{SM}^{(4)} + \frac{1}{\Lambda^2} \sum_{k=1}^{59} \alpha_k \mathcal{O}_k (\mathcal{L}_{BSM})$

$$\begin{aligned}\mathcal{L}_{eff} \supset & c_{ZZ}^{(1)} H Z_\mu Z^\mu + c_{ZZ}^{(2)} H Z_{\mu\nu} Z^{\mu\nu} + c_{Z\tilde{Z}} H Z_{\mu\nu} \tilde{Z}^{\mu\nu} + c_{AZ} H Z_{\mu\nu} A^{\mu\nu} + c_{A\tilde{Z}}^{HZ\mu\nu} \tilde{A}^{\mu\nu} \\ & + H Z_\mu \bar{\ell} \gamma^\mu (c_V + c_A \gamma_5) \ell + Z_\mu \bar{\ell} \gamma^\mu (g_V - g_A \gamma_5) \ell - g_{em} Q_\ell A_\mu \bar{\ell} \gamma^\mu \ell\end{aligned}$$

Where: $c_{ZZ}^{(1)} = m_Z^2 (\sqrt{2} G_F)^{1/2} (1 + \hat{\alpha}_{ZZ}^{(1)})$, $c_{ZZ}^{(2)} = (\sqrt{2} G_F)^{1/2} \hat{\alpha}_{ZZ}$, $c_{Z\tilde{Z}} = (\sqrt{2} G_F)^{1/2} \hat{\alpha}_{Z\tilde{Z}}$,
 $c_{AZ} = (\sqrt{2} G_F)^{1/2} \hat{\alpha}_{AZ}$, $c_{A\tilde{Z}} = (\sqrt{2} G_F)^{1/2} \hat{\alpha}_{A\tilde{Z}}$.

- In this base, the experimental observables G_F, m_Z, α_{em} could be presented:

$$m_Z = m_{Z0}(1 + \delta_Z), \quad G_F = G_{F0}(1 + \delta_{G_F}), \quad \alpha_{em} = \alpha_{em0}(1 + \delta_A)$$

$$\text{where: } \delta_Z = \hat{\alpha}_{ZZ} + \frac{1}{4} \hat{\alpha}_{\Phi D}, \quad \delta_{G_F} = -\hat{\alpha}_{4l} + 2\hat{\alpha}_{\Phi l}^{(3)}, \quad \delta_A = 2\hat{\alpha}_{AA}.$$

$m_{Z0} = 91.1876$
$G_{F0} = 1.166367 \times 10^{-5}$
$\alpha_{em0} = 1/127.940$

Theory framework

[JHEP 03\(2016\) 050](#)

[JHEP 11\(2014\) 028](#)

The $H \rightarrow Zll$ matrix element:

$$\mathcal{M}_{HZ\ell\ell}^\mu = \frac{1}{m_H} \bar{u}(p_3, s_3) \left[\gamma^\mu (H_{1,V} + H_{1,A}\gamma_5) + \frac{q^\mu p}{m_H^2} (H_{2,V} + H_{2,A}\gamma_5) + \frac{\epsilon^{\mu\nu\sigma\rho} p_\nu q_\sigma}{m_H^2} \gamma_\rho (H_{3,V} + H_{3,A}\gamma_5) \right] v(p_4, s_4)$$

- Where $\epsilon_{0123} = +1$ and $q = p_3 + p_4$.

And the parameters in the function are following:

$$H_{1,V} = -\frac{2m_H(\sqrt{2}G_F)^{1/2}r}{r-s} g_V \left(1 + \hat{\alpha}_1^{\text{eff}} - \frac{\kappa}{r} \hat{\alpha}_{ZZ} - \frac{\kappa}{2r} \frac{Q_\ell g_{em}(r-s)}{sg_V} \hat{\alpha}_{AZ} \right)$$

$$H_{1,A} = \frac{2m_H(\sqrt{2}G_F)^{1/2}r}{r-s} g_A \left(1 + \hat{\alpha}_2^{\text{eff}} - \frac{\kappa}{r} \hat{\alpha}_{ZZ} \right),$$

$$H_{2,V} = -\frac{2m_H(\sqrt{2}G_F)^{1/2}}{r-s} g_V \left[2\hat{\alpha}_{ZZ} + \frac{Q_\ell g_{em}(r-s)}{sg_V} \hat{\alpha}_{AZ} \right]$$

$$H_{2,A} = \frac{4m_H(\sqrt{2}G_F)^{1/2}}{r-s} g_A \hat{\alpha}_{ZZ}$$

$$H_{3,V} = -\frac{2m_H(\sqrt{2}G_F)^{1/2}}{r-s} g_V \left[2\hat{\alpha}_{ZZ} + \frac{Q_\ell g_{em}(r-s)}{sg_V} \hat{\alpha}_{AZ} \right]$$

$$H_{3,A} = \frac{4m_H(\sqrt{2}G_F)^{1/2}}{r-s} g_A \hat{\alpha}_{ZZ}$$

$$\hat{\alpha}_1^{\text{eff}} \equiv \hat{\alpha}_{ZZ}^{(1)} - \frac{m_H(\sqrt{2}G_F)^{1/2}(r-s)}{2\sqrt{r}} \frac{\hat{\alpha}_{\Phi l}^V}{g_V}$$

$$\hat{\alpha}_2^{\text{eff}} \equiv \hat{\alpha}_{ZZ}^{(1)} + \frac{m_H(\sqrt{2}G_F)^{1/2}(r-s)}{2\sqrt{r}} \frac{\hat{\alpha}_{\Phi l}^A}{g_A}$$

 : SM term
 Others : EFT contribution


Theory framework

[JHEP 03\(2016\) 050](#) [JHEP 11\(2014\) 028](#)

Differential cross section for $e^+e^- \rightarrow ZH \rightarrow llH$:

$$\frac{d\sigma}{dcos\theta_1 dcos\theta_2 d\phi} = \frac{\mathcal{N}_\sigma(q^2)}{m_H^2} \mathcal{J}(q^2, \theta_1, \theta_2, \phi),$$

$$\mathcal{N}_\sigma(q^2) = \frac{1}{2^{10}(2\pi)^3} \cdot \frac{1}{\sqrt{r}\gamma_Z} \cdot \frac{\sqrt{\lambda(1,s,r)}}{s^2}$$



$$\begin{aligned} \mathcal{J}(q^2, \theta_1, \theta_2, \phi) = & J_1(1 + \cos^2 \theta_1 \cos^2 \theta_2 + \cos^2 \theta_1 + \cos^2 \theta_2) \\ & + J_2 \sin^2 \theta_1 \sin^2 \theta_2 + J_3 \cos \theta_1 \cos \theta_2 \\ & + (J_4 \sin \theta_1 \sin \theta_2 + J_5 \sin 2\theta_1 \sin 2\theta_2) \sin \phi \\ & + (J_6 \sin \theta_1 \sin \theta_2 + J_7 \sin 2\theta_1 \sin 2\theta_2) \cos \phi \\ & + J_8 \sin^2 \theta_1 \sin^2 \theta_2 \sin 2\phi + J_9 \sin^2 \theta_1 \sin^2 \theta_2 \cos 2\phi. \end{aligned}$$

Variables for studying distribution: θ_1, θ_2, ϕ


Theory framework

[JHEP 03\(2016\) 050](#) [JHEP 11\(2014\) 028](#)

Differential cross section for $e^+e^- \rightarrow ZH \rightarrow llH$:

$$\frac{d\sigma}{dcos\theta_1 dcos\theta_2 d\phi} = \frac{\mathcal{N}_\sigma(q^2)}{m_H^2} J(q^2, \theta_1, \theta_2, \phi),$$

$$\mathcal{N}_\sigma(q^2) = \frac{1}{2^{10}(2\pi)^3} \cdot \frac{1}{\sqrt{r}\gamma_Z} \cdot \frac{\sqrt{\lambda(1,s,r)}}{s^2}$$



$$\begin{aligned} J(q^2, \theta_1, \theta_2, \phi) = & J_1(1 + \cos^2 \theta_1 \cos^2 \theta_2 + \cos^2 \theta_1 + \cos^2 \theta_2) \\ & + J_2 \sin^2 \theta_1 \sin^2 \theta_2 + J_3 \cos \theta_1 \cos \theta_2 \\ & + (J_4 \sin \theta_1 \sin \theta_2 + J_5 \sin 2\theta_1 \sin 2\theta_2) \sin \phi \\ & + (J_6 \sin \theta_1 \sin \theta_2 + J_7 \sin 2\theta_1 \sin 2\theta_2) \cos \phi \\ & + J_8 \sin^2 \theta_1 \sin^2 \theta_2 \sin 2\phi + J_9 \sin^2 \theta_1 \sin^2 \theta_2 \cos 2\phi. \end{aligned}$$

Variables for studying distribution: θ_1, θ_2, ϕ

Assumption for simplification:

- $\hat{\alpha}_{A\tilde{Z}}$ and $\hat{\alpha}_{Z\tilde{Z}}$ contribute to cp-odd. (useful parameters)
- Others are set to 0, so $H_{2,V/A} = 0$.

$$\begin{aligned} J_1 &= 2rs(g_A^2 + g_V^2)(|H_{1,V}|^2 + |H_{1,A}|^2), \\ J_2 &= \kappa(g_A^2 + g_V^2)[\kappa(|H_{1,V}|^2 + |H_{1,A}|^2) + \cancel{\lambda} \text{Re}(H_{1,V}H_{2,V}^* + H_{1,A}H_{2,A}^*)], \\ J_3 &= 32rs g_A g_V \text{Re}(H_{1,V}H_{1,A}^*), \\ J_4 &= 4\kappa \sqrt{rs\lambda} g_A g_V \text{Re}(H_{1,V}H_{3,A}^* + H_{1,A}H_{3,V}^*), \\ J_5 &= \frac{1}{2}\kappa \sqrt{rs\lambda}(g_A^2 + g_V^2) \text{Re}(H_{1,V}H_{3,V}^* + H_{1,A}H_{3,A}^*), \\ J_6 &= 4\sqrt{rs} g_A g_V [4\kappa \text{Re}(H_{1,V}H_{1,A}^*) + \cancel{\lambda} \text{Re}(H_{1,V}H_{2,A}^* + H_{1,A}H_{2,V}^*)], \\ J_7 &= \frac{1}{2}\sqrt{rs}(g_A^2 + g_V^2)[2\kappa(|H_{1,V}|^2 + |H_{1,A}|^2) + \cancel{\lambda} \text{Re}(H_{1,V}H_{2,V}^* + H_{1,A}H_{2,A}^*)], \\ J_8 &= 2rs\sqrt{\lambda}(g_A^2 + g_V^2) \text{Re}(H_{1,V}H_{3,V}^* + H_{1,A}H_{3,A}^*), \\ J_9 &= 2rs(g_A^2 + g_V^2)(|H_{1,V}|^2 + |H_{1,A}|^2). \end{aligned}$$

6 of these 9 functions are independent

$\cancel{\lambda}$	0 in assumption
	EFT CP-odd term
Others	CP-even contribution

Optimal variable approach

Differential cross section could be represent as:

$$\frac{d\sigma}{d\cos\theta_1 d\cos\theta_2 d\phi} = N \times \left(J_{\text{even}}(\theta_1, \theta_2, \phi) + \hat{\alpha}_{A\tilde{Z}} \times J_{\text{odd}_1}(\theta_1, \theta_2, \phi) + \hat{\alpha}_{Z\tilde{Z}} \times J_{\text{odd}_2}(\theta_1, \theta_2, \phi) \right)$$







where $\hat{\alpha}_{A\tilde{Z}}$ and $\hat{\alpha}_{Z\tilde{Z}}$ are CP-violating parameters.

Optimal variable approach

Differential cross section could be represent as:

$$\frac{d\sigma}{dcos\theta_1 dcos\theta_2 d\phi} = N \times (J_{even}(\theta_1, \theta_2, \phi) + \hat{\alpha}_{A\tilde{Z}} \times J_{odd_1}(\theta_1, \theta_2, \phi) + \hat{\alpha}_{Z\tilde{Z}} \times J_{odd_2}(\theta_1, \theta_2, \phi))$$

where $\hat{\alpha}_{A\tilde{Z}}$ and $\hat{\alpha}_{Z\tilde{Z}}$ are CP-violating parameters.



Optimal variable approach

[PLB 306 \(1993\) 411-417](#)

$$\frac{d\sigma}{dcos\theta_1 dcos\theta_2 d\phi} = N \times \left(J_{even}(\theta_1, \theta_2, \phi) + \hat{\alpha}_{A\tilde{Z}} \times J_{odd_1}(\theta_1, \theta_2, \phi) + \hat{\alpha}_{Z\tilde{Z}} \times J_{odd_2}(\theta_1, \theta_2, \phi) \right)$$



In this formation, we could define **Optimal Variable ω** which combines the information from $\{\theta_1, \theta_2, \phi\}$:

$$\omega_1 = \frac{J_{odd_1}(\theta_1, \theta_2, \phi)}{J_{even}(\theta_1, \theta_2, \phi)} \text{ to measure } \hat{\alpha}_{A\tilde{Z}}$$

$$\omega_2 = \frac{J_{odd_2}(\theta_1, \theta_2, \phi)}{J_{even}(\theta_1, \theta_2, \phi)} \text{ to measure } \hat{\alpha}_{Z\tilde{Z}}$$

Benefits:

- Combine the information from 3-dimension phase space
- Easier to study



Monte Carlo samples

Samples:

- The SM Higgs and background samples: generate with Whizard 1.95 and full simulated based on the CEPC baseline detector design. (Used to calculate the selection efficiencies and study background.)
- CP-mixing Higgs samples: generate according to differential cross section for $e^+e^- \rightarrow ZH \rightarrow llH$:

$$\frac{d\sigma}{dcos\theta_1 dcos\theta_2 d\phi} = \frac{\mathcal{N}_\sigma(q^2)}{m_H^2} \mathcal{J}(q^2, \theta_1, \theta_2, \phi),$$

- $\sqrt{s} = 240\text{GeV}$
- The mass of Higgs boson is set to be 125GeV and the couplings are set to the SM predictions.
- All the generations are normalized to the expected yields in data with an integrated luminosity of 5.6ab^{-1} .

Event selection

(Using the SM Higgs and background samples)

- **Signal:** $e^+e^- \rightarrow ZH \rightarrow \mu^+\mu^-H(\rightarrow b\bar{b}/c\bar{c}/gg)$ channel
- **Background:** Irreducible background which contains the same final states as that in signal.

Choose selections by the best significance:

- Lepton pair selection:

$$|\cos\theta_{\mu^+\mu^-}| < 0.81 ; \quad \text{Mass}_{\mu\mu} \in (77.5\text{GeV}, 104.5\text{GeV}) ; \quad M_{\text{recoil}\,\mu\mu} \in (124\text{GeV}, 140\text{GeV}).$$

$$\text{Where } M_{\text{recoil}\,\mu\mu}^2 = (\sqrt{s} - E_{\mu\mu})^2 - p_{\mu\mu}^2 = s - 2E_{\mu\mu}\sqrt{s} + m_{\mu\mu}^2$$

- Jets pair selection:

$$|\cos\theta_{jet}| < 0.96 ; \quad \text{Mass}_{jj} \in (100\text{GeV}, 150\text{GeV}).$$

-
- **Signal:** $e^+e^- \rightarrow ZH \rightarrow e^+e^-H(\rightarrow b\bar{b}/c\bar{c}/gg)$ channel
 - **Background:** Irreducible background which contains the same final states as that in signal.

Choose selections by the best significance:

- Lepton pair selection:

$$|\cos\theta_{e^+e^-}| < 0.81 ; \quad \text{Mass}_{ee} \in (85\text{GeV}, 95\text{GeV}) ; \quad M_{\text{recoil}\,ee} \in (124\text{GeV}, 140\text{GeV}); \quad |\cos\varphi_{e^+}| < 0.95 ; \quad |\cos\varphi_{e^-}| < 0.95 ;$$

- Jets pair selection:

$$|\cos\theta_{jet}| < 0.96 ; \quad \text{Mass}_{jj} \in (110\text{GeV}, 145\text{GeV}).$$

Event selection

$ZH \rightarrow \mu^+ \mu^- + b\bar{b}/c\bar{c}/gg$ channel

	Signal	Irreducible Background
Original	2.62×10^4	1.25×10^6
Lepton pair selection	1.59×10^4 (efficiency:60.67%)	9.91×10^3 (efficiency:0.79%)
All selection	1.48×10^4 (efficiency:56.42%)	5.60×10^3 (efficiency:0.45%)



$ZH \rightarrow e^+ e^- + b\bar{b}/c\bar{c}/gg$ channel

	Signal	Irreducible Background
Original	2.72×10^4	1.77×10^6
Lepton pair selection	8.76×10^3 (efficiency:32.2%)	8.77×10^4 (efficiency:0.50%)
All selection	7.15×10^3 (efficiency:26.3%)	4.59×10^3 (efficiency:0.25%)

Event selection

Correlation:

- We can see that θ_1, θ_2, ϕ have little correlation with $\cos\theta_{l^+l^-}$, Mass_{ll} , M_{recoil_ll} , $\cos\theta_{\text{jet}}$, Mass_{jj} .



- So we can ignore the impact of event selections to $\theta_1, \theta_2, \text{and } \phi$.



Fitting strategy and result

Fit strategy: Maximum-likelihood fit

$$f^{\vec{\alpha}}(\omega) = N_{\text{sig}} * f_{\text{sig}}^{\vec{\alpha}}(\omega) + N_{\text{bkg}} * f_{\text{bkg}}^{\vec{\alpha}}(\omega)$$

where $\vec{\alpha}$ means $\hat{\alpha}_{A\bar{Z}}$ and $\hat{\alpha}_{Z\bar{Z}}$, ω represents ω_1 and ω_2 .

- Fit ω to get $f_{\text{sig}}^{\vec{\alpha}}(\omega)$ and $f_{\text{bkg}}^{\vec{\alpha}}(\omega)$
- Fit $M_{\text{recoil_}\mu\mu}$ to get N_{sig} and N_{bkg}
- Evaluate likelihood function for each $\vec{\alpha}$ value hypothesis, and construct a ΔNLL as a function of $\vec{\alpha}$.






Fitting strategy and result

Fit ω :

- Use histogram pdf to fit **MC signal and background sample**.
- The red curve is global fit, the green curve is signal events, the blue curve is background events.

Fit $M_{\text{recoil_}\mu\mu}$:

- The signal modeled by the Crystal Ball function.
- The background modeled by a second-order polynomial.
- Using **ISR sample** can simulate the small exponential tail (which corresponding to the expected distribution.)



Indvidual Fit

Extract maximum-likelihood fit p-value and interval


- In $\mu^+ \mu^- H$ channel.
- Fit ΔNLL curve with a quadratic function $\Delta NLL(\vec{\alpha}) = a \cdot (\vec{\alpha} - \vec{\alpha}_0)^2$
- 68%(95%) CL interval corresponds to $\Delta NLL=0.5(1.96)$.
- Set: fit to $\hat{\alpha}_{A\tilde{Z}}, \hat{\alpha}_{Z\tilde{Z}} = 0$.

$$\Delta NLL(\hat{\alpha}_{A\tilde{Z}} | \omega_1) = 2.93 \times 10^{-4} (\hat{\alpha}_{A\tilde{Z}} + 8.68 \times 10^{-1})^2$$

For $\hat{\alpha}_{A\tilde{Z}}$:

$$68\% \text{ CL: } [-4.16 \times 10^{-2}, 3.88 \times 10^{-2}]$$

$$95\% \text{ CL: } [-8.10 \times 10^{-2}, 7.82 \times 10^{-2}]$$




- Set: fit to $\hat{\alpha}_{A\tilde{Z}}, \hat{\alpha}_{Z\tilde{Z}} = 0$.

$$\Delta NLL(\hat{\alpha}_{Z\tilde{Z}} | \omega_2) = 4.51 \times 10^{-3} (\hat{\alpha}_{Z\tilde{Z}} + 6.36 \times 10^{-1})^2$$

For $\hat{\alpha}_{Z\tilde{Z}}$:

$$68\% \text{ CL: } [-1.06 \times 10^{-2}, 1.00 \times 10^{-3}]$$

$$95\% \text{ CL: } [-2.06 \times 10^{-2}, 2.01 \times 10^{-2}]$$



Fit to phi

ϕ has the most information among the three kinematic variables (θ_1, θ_2, ϕ)

straight-forward to fit ϕ .

In $\mu^+ \mu^- H$ channel.

	$\hat{\alpha}_{A\tilde{Z}} (\times 10^{-2})$	$\hat{\alpha}_{Z\tilde{Z}} (\times 10^{-2})$
ω -fitting		
68% CL(1σ)	[-4.16, 3.88]	[-1.06, 1.00]
95% CL(2σ)	[-8.10, 7.82]	[-2.06, 2.01]
ϕ -fitting		
68% CL(1σ)	[-4.42, 4.21]	[-1.35, 1.24]
95% CL(2σ)	[-8.66, 8.45]	[-2.62, 2.51]

The results of ϕ -fitting is slight worse than those of the ω -fitting.

- θ_1 and θ_2 have less information.

Combined results

$5.6ab^{-1}$	$\hat{\alpha}_{A\tilde{Z}}(1\sigma)$	$\hat{\alpha}_{Z\tilde{Z}}(1\sigma)$	$\hat{\alpha}_{A\tilde{Z}}(2\sigma)$	$\hat{\alpha}_{Z\tilde{Z}}(2\sigma)$
$\mu^+\mu^-$	$[-4.16 \times 10^{-2}, 3.88 \times 10^{-2}]$	$[-1.06 \times 10^{-2}, 1.00 \times 10^{-2}]$	$[-8.10 \times 10^{-2}, 7.82 \times 10^{-2}]$	$[-2.06 \times 10^{-2}, 2.01 \times 10^{-2}]$
e^+e^-	$[-5.97 \times 10^{-2}, 5.43 \times 10^{-2}]$	$[-1.48 \times 10^{-2}, 1.48 \times 10^{-2}]$	$[-1.16 \times 10^{-1}, 1.10 \times 10^{-1}]$	$[-2.93 \times 10^{-2}, 2.93 \times 10^{-2}]$
<i>combine</i>	$[-3.47 \times 10^{-2}, 3.10 \times 10^{-2}]$	$[-8.50 \times 10^{-3}, 8.27 \times 10^{-3}]$	$[-6.69 \times 10^{-2}, 6.32 \times 10^{-2}]$	$[-1.67 \times 10^{-2}, 1.65 \times 10^{-2}]$

Combined results

$5.6ab^{-1}$	$\hat{\alpha}_{A\bar{Z}}(1\sigma)$	$\hat{\alpha}_{Z\bar{Z}}(1\sigma)$	$\hat{\alpha}_{A\bar{Z}}(2\sigma)$	$\hat{\alpha}_{Z\bar{Z}}(2\sigma)$
$\mu^+\mu^-$	$[-4.16 \times 10^{-2}, 3.88 \times 10^{-2}]$	$[-1.06 \times 10^{-2}, 1.00 \times 10^{-2}]$	$[-8.10 \times 10^{-2}, 7.82 \times 10^{-2}]$	$[-2.06 \times 10^{-2}, 2.01 \times 10^{-2}]$
e^+e^-	$[-5.97 \times 10^{-2}, 5.43 \times 10^{-2}]$	$[-1.48 \times 10^{-2}, 1.48 \times 10^{-2}]$	$[-1.16 \times 10^{-1}, 1.10 \times 10^{-1}]$	$[-2.93 \times 10^{-2}, 2.93 \times 10^{-2}]$
combine	$[-3.47 \times 10^{-2}, 3.10 \times 10^{-2}]$	$[-8.50 \times 10^{-3}, 8.27 \times 10^{-3}]$	$[-6.69 \times 10^{-2}, 6.32 \times 10^{-2}]$	$[-1.67 \times 10^{-2}, 1.65 \times 10^{-2}]$

Consider CEPC Higgs operation can be upgraded to $20ab^{-1}$.

- Normalize samples to the expected yields in data with an integrated luminosity of $20ab^{-1}$.
- This result is about **two times** better than that of $5.6ab^{-1}$

$20ab^{-1}$	$\hat{\alpha}_{A\bar{Z}}(1\sigma)$	$\hat{\alpha}_{Z\bar{Z}}(1\sigma)$	$\hat{\alpha}_{A\bar{Z}}(2\sigma)$	$\hat{\alpha}_{Z\bar{Z}}(2\sigma)$
$\mu^+\mu^-$	$[-2.19 \times 10^{-2}, 2.03 \times 10^{-2}]$	$[-5.38 \times 10^{-3}, 5.16 \times 10^{-3}]$	$[-4.26 \times 10^{-2}, 4.04 \times 10^{-2}]$	$[-1.05 \times 10^{-2}, 1.03 \times 10^{-2}]$
e^+e^-	$[-3.22 \times 10^{-2}, 2.75 \times 10^{-2}]$	$[-7.63 \times 10^{-3}, 7.85 \times 10^{-3}]$	$[-6.15 \times 10^{-2}, 5.68 \times 10^{-2}]$	$[-1.52 \times 10^{-2}, 1.54 \times 10^{-2}]$
combine	$[-1.85 \times 10^{-2}, 1.59 \times 10^{-2}]$	$[-4.20 \times 10^{-3}, 4.30 \times 10^{-3}]$	$[-3.54 \times 10^{-2}, 3.27 \times 10^{-2}]$	$[-8.36 \times 10^{-3}, 8.46 \times 10^{-3}]$

Result compare --> Compared with HL-LHC

In order to compare our study with HL-LHC, some conversion is necessary. (show in backup.)

In HL-LHC: (1sigma)

Analysis \ Parameter	$\bar{c}_{Z\gamma}$	\bar{c}_{ZZ}	Case
Analysis	$\bar{c}_{Z\gamma}$	\bar{c}_{ZZ}	
HL-LHC (4ℓ , incl.)	[-0.22,0.22]	[-0.33,0.33]	IP
	[-0.25,0.25]	[-0.27,0.27]	IP _{marg.}
HL-LHC (4ℓ , diff.)	[-0.10,0.10]	[-0.31,0.31]	IP
	[-0.13,0.13]	[-0.22,0.22]	IP _{marg.}
HE-LHC (4ℓ , incl.)	[-0.18,0.18]	[-0.17,0.17]	IP
	[-0.23,0.23]	[-0.20,0.20]	IP _{marg.}
HE-LHC (4ℓ , diff.)	[-0.05,0.05]	[-0.13,0.13]	IP
	[-0.06,0.06]	[-0.10,0.10]	IP _{marg.}

[arXiv:1902.00134](https://arxiv.org/abs/1902.00134)

Result compare --> Compared with HL-LHC

In order to compare our study with HL-LHC, some conversion is necessary. (show in backup.)

In HL-LHC: (1sigma)

Analysis \ Parameter	$\tilde{c}_{Z\gamma}$	\tilde{c}_{ZZ}	Case
Analysis	$\tilde{c}_{Z\gamma}$	\tilde{c}_{ZZ}	
HL-LHC (4ℓ , incl.)	$[-0.22, 0.22]$ $[-0.25, 0.25]$	$[-0.33, 0.33]$ $[-0.27, 0.27]$	IP IP _{marg.}
HL-LHC (4ℓ , diff.)	$[-0.10, 0.10]$ $[-0.13, 0.13]$	$[-0.31, 0.31]$ $[-0.22, 0.22]$	IP IP _{marg.}
HE-LHC (4ℓ , incl.)	$[-0.18, 0.18]$ $[-0.23, 0.23]$	$[-0.17, 0.17]$ $[-0.20, 0.20]$	IP IP _{marg.}
HE-LHC (4ℓ , diff.)	$[-0.05, 0.05]$ $[-0.06, 0.06]$	$[-0.13, 0.13]$ $[-0.10, 0.10]$	IP IP _{marg.}

[arXiv:1902.00134](https://arxiv.org/abs/1902.00134)

Collider	pp	e^+e^-	e^+e^-
E (GeV)	14000	240	240
\mathcal{L} (fb $^{-1}$)	3000	5600	20000
$\tilde{c}_{Z\gamma}$ (1 σ)	$[-0.22, 0.22]$	$[-0.30, 0.27]$	$[-0.16, 0.14]$
\tilde{c}_{ZZ} (1 σ)	$[-0.33, 0.33]$	$[-0.06, 0.06]$	$[-0.03, 0.03]$

The results in $5.6ab^{-1}$ are significantly better than HL-LHC on the \tilde{c}_{ZZ} and comparable on the $\tilde{c}_{Z\gamma}$.

◦ For $5.6ab^{-1}$: $f_{CP}^{HZZ} < 9.16 \times 10^{-6}$

Summary

An EFT based Higgs CP-mixing test is performed.

- Set up some basic assumptions to have a simplest CP-mixing model.
- Introduced optimal variable with better performance.
- Used ML-fit in ω and ϕ distribution to extract $\hat{\alpha}_{A\tilde{Z}}$ and $\hat{\alpha}_{Z\tilde{Z}}$.
- Result: 68% CL $\tilde{c}_{Z\gamma} \in [-0.30, 0.27]$ and $\tilde{c}_{ZZ} \in [-0.06, 0.06]$; $f_{CP}^{HZZ} < 9.16 \times 10^{-6}$.
- Consider CEPC Higgs operation can be upgraded to $20ab^{-1}$, the results is about **two times** better.

In \tilde{c}_{ZZ} , the sensitivities is improved by about **one order of magnitude** compared with pp collider.

Collider	pp	e^+e^-	e^+e^-
E (GeV)	14000	240	240
\mathcal{L} (fb^{-1})	3000	5600	20000
$\tilde{c}_{Z\gamma}$ (1σ)	$[-0.22, 0.22]$	$[-0.30, 0.27]$	$[-0.16, 0.14]$
\tilde{c}_{ZZ} (1σ)	$[-0.33, 0.33]$	$[-0.06, 0.06]$	$[-0.03, 0.03]$

Thank you!

Backup

Result compare --> Compared with HL-LHC

In HL-LHC: [arXiv:1902.00134](https://arxiv.org/abs/1902.00134)

$$\mathcal{L}_{\text{CPV}} = \frac{H}{v} \left[\tilde{c}_{\gamma\gamma} \frac{e^2}{4} A_{\mu\nu} \tilde{A}^{\mu\nu} + \tilde{c}_{Z\gamma} \frac{e\sqrt{g_1^2 + g_2^2}}{2} Z_{\mu\nu} \tilde{A}^{\mu\nu} + \tilde{c}_{ZZ} \frac{g_1^2 + g_2^2}{4} Z_{\mu\nu} \tilde{Z}^{\mu\nu} + \tilde{c}_{WW} \frac{g_2^2}{2} W_{\mu\nu}^+ \tilde{W}^{\mu\nu} \right]$$

Compare theory model in P5, we can get that the value in red frame are same:

$$(g1=0.358, g2=0.648, e=0.313, v = 1/\sqrt{\sqrt{2}G_F^0} = 2M_W/g \approx 246.22\text{GeV})$$

$$(\sqrt{2}G_F)^{1/2} \boxed{\hat{\alpha}_{Z\tilde{Z}}} H Z_{\mu\nu} \tilde{Z}^{\mu\nu} = \frac{H}{v} \boxed{\tilde{c}_{ZZ} \frac{g_1^2 + g_2^2}{4}} Z_{\mu\nu} \tilde{Z}^{\mu\nu}$$
$$\frac{g_1^2 + g_2^2}{4} = 0.137$$

$$(\sqrt{2}G_F)^{1/2} \boxed{\hat{\alpha}_{A\tilde{Z}}} H Z_{\mu\nu} \tilde{A}^{\mu\nu} = \frac{H}{v} \boxed{\tilde{c}_{Z\gamma} \frac{e\sqrt{g_1^2 + g_2^2}}{2}} Z_{\mu\nu} \tilde{A}^{\mu\nu}$$
$$\frac{e\sqrt{g_1^2 + g_2^2}}{2} = 0.116$$