

# Precision Higgs Physics at Current and Future Colliders



Shufang Su • U. of Arizona

Higgs Potential 2022

Peking University

July 26, 2022

Time flies ....

10 Years into Higgs Discovery !

Time for Precision Higgs Era !

# Where Are We Now?

---

- ◎ Our wish list has not change much from 10 years ago.
- ◎ Discovery of Higgs and measurements of its property
  - ➔ Exclude certain models (technicolor,...)
  - ➔ Narrow down parameter space
- ◎ Non-discovery of anything else
  - ➔ New physics gets heavier
  - ➔ A bit uncomfortable, big picture unchanged

# Then What?

---

Where is New Physics?

larger mass? Small Coupling? Too much BG?

# Then What?

---

Where is New Physics?

larger mass? Small Coupling? Too much BG?

- ◎ Direct search for new particles

# Then What?

---

## Where is New Physics?

larger mass? Small Coupling? Too much BG?

- ◎ Direct search for new particles

Need colliders with larger energies (pp or e+e- with large  $E_{\text{cm}}$ )

# Then What?

---

## Where is New Physics?

larger mass? Small Coupling? Too much BG?

- ◎ Direct search for new particles

Need colliders with larger energies (pp or e+e- with large  $E_{\text{cm}}$ )

# Then What?

## Where is New Physics?

larger mass? Small Coupling? Too much BG?

- ◎ Direct search for new particles

Need colliders with larger energies (pp or e+e- with large  $E_{\text{cm}}$ )

- ◎ Indirect search for imprints on W, Z, top and Higgs



# Then What?

## Where is New Physics?

larger mass? Small Coupling? Too much BG?

- ◎ Direct search for new particles

Need colliders with larger energies (pp or e+e- with large  $E_{\text{cm}}$ )

- ◎ Indirect search for imprints on W, Z, top and Higgs

Need colliders/measurements with unprecedented accuracy

# Then What?

## Where is New Physics?

larger mass? Small Coupling? Too much BG?

- ◎ Direct search for new particles

Need colliders with larger energies (pp or e+e- with large  $E_{cm}$ )

- ◎ Indirect search for imprints on W, Z, top and Higgs

Need colliders/measurements with unprecedented accuracy

(e+e- or pp with high luminosity)

# Then What?

## Where is New Physics?

larger mass? Small Coupling? Too much BG?

- ◎ Direct search for new particles

Need colliders with larger energies (pp or e+e- with large  $E_{cm}$ )



- ◎ Indirect search for imprints on W, Z, top and Higgs


Need colliders/measurements with unprecedented accuracy

(e+e- or pp with high luminosity)

# LHC / HL-LHC Plan



## LHC / HL-LHC Plan



**We are here.**

### HL-LHC CIVIL ENGINEERING:

DEFINITION	EXCAVATION	BUILDINGS
------------	------------	-----------

**LHC is a Higgs factory: 15 M Higgs**  
**HL-LHC: 170 M Higgs, 120 K HH pair**

# Current and Future Colliders

---

**pp**

**e<sup>+</sup>e<sup>-</sup>**

# Current and Future Colliders


---

**pp**


**e<sup>+</sup>e<sup>-</sup>**



# Current and Future Colliders




# Current and Future Colliders







# Current and Future Colliders




# Current and Future Colliders




# Current and Future Colliders



# Current and Future Colliders




# Current and Future Colliders




**LHC**  
**HL-LHC**  
**SPPC**  
**FCC-hh**  
**...**

# Current and Future Colliders




LHC  
HL-LHC  
SPPC  
FCC-hh  
...

# Current and Future Colliders



LHC  
HL-LHC  
SPPC  
FCC-hh  
...


# Current and Future Colliders



LHC  
HL-LHC  
SPPC  
FCC-hh  
...




# Current and Future Colliders



LHC  
HL-LHC  
SPPC  
FCC-hh  
...


# Current and Future Colliders




larger mass  $E_{cm}$

LHC  
HL-LHC  
SPPC  
FCC-hh  
...


# Current and Future Colliders




LHC  
HL-LHC  
SPPC  
FCC-hh  
...



# Current and Future Colliders



LHC  
HL-LHC  
SPPC  
FCC-hh  
...



CEPC  
FCC-ee  
ILC  
...

# A Light Higgs is Puzzling...

- ⊙ Light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$ 
  - ➔ spin 0, a new kind of fundamental particle, no charge, no structure
  - ➔ Nothing protects its mass  $\Rightarrow$  New physics beyond the SM

# A Light Higgs is Puzzling...

- ◎ Light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$ 
  - ➔ spin 0, a new kind of fundamental particle, no charge, no structure
  - ➔ Nothing protects its mass  $\Rightarrow$  New physics beyond the SM

Then What?

# A Light Higgs is Puzzling...

- ◎ Light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$ 
  - ➔ spin 0, a new kind of fundamental particle, no charge, no structure
  - ➔ Nothing protects its mass  $\Rightarrow$  New physics beyond the SM

Then What? Still a lot of hard, but fun work to do!

# A Light Higgs is Puzzling...

- ⊙ Light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$ 
  - spin 0, a new kind of fundamental particle, no charge, no structure
  - Nothing protects its mass  $\Rightarrow$  New physics beyond the SM


Then What?    Theoretically ...



# A Light Higgs is Puzzling...

- ⊙ Light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$ 
  - spin 0, a new kind of fundamental particle, no charge, no structure
  - Nothing protects its mass  $\Rightarrow$  New physics beyond the SM

Then What?    Theoretically ...



$$V(\phi) = \frac{1}{2}\mu_h^2\phi^2 + \frac{\lambda}{4}\phi^4$$

$$\langle\phi\rangle \equiv v \neq 0 \quad \rightarrow \quad m_W = g_W \frac{v}{2}$$


$$M_H^2 = -2\mu^2 = 2\lambda v^2$$

# A Light Higgs is Puzzling...

- Light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$ 
  - spin 0, a new kind of fundamental particle, no charge, no structure
  - Nothing protects its mass  $\Rightarrow$  New physics beyond the SM

Then What? Theoretically ...

- $\lambda \sim 1/8$ , origin of  $\lambda$  ?
- extended Higgs sector?
- stabilization of EW scale?
- ...



$$V(\phi) = \frac{1}{2}\mu_h^2\phi^2 + \frac{\lambda}{4}\phi^4$$

$$\langle\phi\rangle \equiv v \neq 0 \quad \rightarrow \quad m_W = g_W \frac{v}{2}$$

$$M_H^2 = -2\mu^2 = 2\lambda v^2$$

# Then What?

light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$

Then What? experimentally...

# Then What?

light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$

Then What? experimentally...

- © Is it a SM Higgs? Mass, width, spin, coupling, CP,...

# Then What?

light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$

Then What? experimentally...

- Is it a SM Higgs? Mass, width, spin, coupling, CP, ...
- Is there more than one Higgs boson?

# Then What?

light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$

Then What? experimentally...

- ⦿ Is it a SM Higgs? Mass, width, spin, coupling, CP, ...
- ⦿ Is there more than one Higgs boson?
- ⦿ Does this H decay to other things unexpected?

# Then What?

light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$

Then What? experimentally...

- Is it a SM Higgs? Mass, width, spin, coupling, CP, ...
- Is there more than one Higgs boson?
- Does this H decay to other things unexpected?
- Can we use H to look for new physics?

# Then What?

light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$

Then What? experimentally...

- Is it a SM Higgs? Mass, width, spin, coupling, CP, ...
- Is there more than one Higgs boson?
- Does this H decay to other things unexpected?
- Can we use H to look for new physics?
- Where is new physics? Top partners? Dark matter?



# Then What?

light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$

Then What? experimentally...

- Is it a SM Higgs? Mass, width, spin, coupling, CP, ...
- Is there more than one Higgs boson?
- Does this H decay to other things unexpected?
- Can we use H to look for new physics?
- Where is new physics? Top partners? Dark matter?
- ...

# Then What?

light, weakly coupled boson:  $m_h = 125\text{-}126\text{ GeV}$ ,  $\Gamma \ll 1\text{ GeV}$

Then What? experimentally...

- Is it a SM Higgs? Mass, width, spin, coupling, CP, ...
- Is there more than one Higgs boson?
- Does this H decay to other things unexpected?
- Can we use H to look for new physics?
- Where is new physics? Top partners? Dark matter?
- ...

**This talk focuses on the Higgs precision measurements.**

# Outline


---

- Introduction
- Precision Higgs measurements: current/future
- Implication of Higgs precision measurements

# Precision Higgs Measurements





## Precision Higgs Measurements


- Mass, width, spin, CP
- Higgs couplings
- differential distributions, STXS, Global fits



from Ed Scott



# LHC Higgs Observation

Channel categories	Br	ggF  ~4 M vets produced	VBF  ~300 k vets produced	VH  ~200 k vets produced	ttH  ~40 k evts produced
Cross Section 13 TeV (8 TeV)		48.6 (21.4) pb*	3.8 (1.6) pb	2.3 (1.1) pb	0.5 (0.1) pb
Observed modes	$\gamma\gamma$	0.2 %	✓	✓	✓
	ZZ	3%	✓	✓	✓
	WW	22%	✓	✓	✓
	$\tau\tau$	6.3 %	✓	✓	✓
	bb	55%	✓	✓	✓
Remaining to be observed	Z $\gamma$ and $\gamma\gamma^*$	0.2 %	✓	✓	✓
	$\mu\mu$	0.02 %	✓	✓	✓
Limits	Invisible	0.1 %	✓ (monojet)	✓	✓



# LHC Precision Higgs Measurements

Mass



LHC

**ATLAS**

$$m_H = 124.92 \pm 0.19(\text{stat.})_{-0.06}^{+0.09}(\text{syst.}) \text{ GeV.}$$

PLB 784 (2018) 345  
ATLAS-CONF-2020-005

**CMS**

$$m_H = 125.38 \pm 0.14 \text{ GeV}$$

JHEP11 (2017) 047  
PLB 805 (2020) 135425

HL-LHC: 10-20 MeV

e+e- Higgs factory: < 6 MeV

# LHC Precision Higgs Measurements

## Width

### LHC

• On-shell (CMS 4I)  $\Gamma_H < 1.1 \text{ GeV}$  at 95% CL

JHEP 11 (2017) 047

• Off-shell

$$\frac{\sigma_{vv \rightarrow H \rightarrow 4\ell}^{\text{off-shell}}}{\sigma_{vv \rightarrow H \rightarrow 4\ell}^{\text{on-shell}}} \propto \Gamma_H$$

ATLAS

$$\Gamma_H < 14.4 \text{ MeV}$$

PLB 786 (2018) 223


CMS

$$\Gamma_H = 3.2^{+2.4}_{-1.7} \text{ MeV}$$

CMS-PAS-HIG-21-013

HL-LHC  $\Gamma_H = 4.1^{+1.0}_{-1.1}$

e+e- Higgs factory: 0.1 MeV



# LHC Precision Higgs Measurements

## Couplings

$$\mathcal{L}_{\text{SM}} = \dots + |D_\mu \phi|^2 + \psi_i y_{ij} \psi_j \phi - V(\phi)$$


Gauge interactions  
studied for many decades  
now with a scalar

Higgs self-interactions  
unobserved

Yukawa interactions  
new  
study started in 2018



# LHC Precision Higgs Measurements



ATLAS-CONF-2021-053

S. Su

$$\kappa_f = \frac{g(hff)}{g(hff; \text{SM})}, \quad \kappa_V = \frac{g(hVV)}{g(hVV; \text{SM})}$$


	ATLAS - CMS Run 1 combination	ATLAS Run 2	CMS Run 2	Current precision
$\kappa_\gamma$	13%	$1.04 \pm 0.06$	$1.01^{+0.09}_{-0.14}$	6%
$\kappa_W$	11%	$1.06 \pm 0.06$	$-1.11^{+0.14}_{-0.09}$	6%
$\kappa_Z$	11%	$0.99 \pm 0.06$	$0.96 \pm 0.07$	6%
$\kappa_g$	14%	$0.92^{+0.07}_{-0.06}$	$1.16^{+0.12}_{-0.11}$	7%
$\kappa_t$	30%	$0.92 \pm 0.10$	$1.01 \pm 0.11$	11%
$\kappa_b$	26%	$0.87 \pm 0.11$	$1.18^{+0.19}_{-0.27}$	11%
$\kappa_\tau$	15%	$0.92 \pm 0.07$	$0.94 \pm 0.12$	8%

JHEP 08 (2016) 045

ATLAS-CONF-2021-053

CMS-PAS-HIG-19-005

# HL-LHC



- 2-4% for most couplings,  $Z\gamma$  10%
- $\mu\mu$ ,  $Z\gamma$  statistical limited
- Others dominated by theoretical uncertainties

# Theoretical Uncertainties

from Gavin Salam

$$\sigma = \sum_{i,j} \int dx_1 dx_2 f_{i/p}(x_1) f_{j/p}(x_2) \hat{\sigma}(x_1 x_2 s) \times [1 + \mathcal{O}(\Lambda/M)^p]$$

Parton distribution functions (PDFs)  
(non-perturbative, universal)


non-perturbative effects  
(power suppressed)

hard scattering  
(perturbative)

$$\sigma_{ggF} = 48.68 \pm 3.9 \text{ (scales)} \pm 1.9 \text{ (PDF)} \pm 2.6 \text{ } (\alpha_S) \text{ Pb}$$

Lots of hard work to be done to reduce the theoretical uncertainty.

# Precision Measurements @ Higgs Factory




- Determine all Higgs couplings (model-independent)
- Infer Higgs total decay width
- probe invisible/exotic Higgs decay

collider	CEPC	FCC-ee			ILC				
$\sqrt{s}$	240 GeV	240 GeV	365 GeV	250 GeV	350 GeV	500 GeV			
$\int \mathcal{L} dt$	5.6 ab <sup>-1</sup>	5 ab <sup>-1</sup>	1.5 ab <sup>-1</sup>	2 ab <sup>-1</sup>	200 fb <sup>-1</sup>	4 ab <sup>-1</sup>			
production	<i>Zh</i>	<i>Zh</i>	<i>Zh</i>	$\nu\bar{\nu}h$	<i>Zh</i>	<i>Zh</i>	$\nu\bar{\nu}h$	<i>Zh</i>	$\nu\bar{\nu}h$
$\Delta\sigma/\sigma$	0.5%	0.5%	0.9%	–	0.71%	2.0%	–	1.05	–
decay	$\Delta(\sigma \cdot BR)/(\sigma \cdot BR)$								
$h \rightarrow b\bar{b}$	0.27%	0.3%	0.5%	0.9%	0.46%	1.7%	2.0%	0.63%	0.23%
$h \rightarrow c\bar{c}$	3.3%	2.2%	6.5%	10%	2.9%	12.3%	21.2%	4.5%	2.2%
$h \rightarrow gg$	1.3%	1.9%	3.5%	4.5%	2.5%	9.4%	8.6%	3.8%	1.5%
$h \rightarrow WW^*$	1.0%	1.2%	2.6%	3.0%	1.6%	6.3%	6.4%	1.9%	0.85%
$h \rightarrow \tau^+\tau^-$	0.8%	0.9%	1.8%	8.0%	1.1%	4.5%	17.9%	1.5%	2.5%
$h \rightarrow ZZ^*$	5.1%	4.4%	12%	10%	6.4%	28.0%	22.4%	8.8%	3.0%
$h \rightarrow \gamma\gamma$	6.8%	9.0%	18%	22%	12.0%	43.6%	50.3%	12.0%	6.8%
$h \rightarrow \mu^+\mu^-$	17%	19%	40%	–	25.5%	97.3%	178.9%	30.0%	25.0%
$(\nu\bar{\nu})h \rightarrow b\bar{b}$	2.8%	3.1%	–	–	3.7%	–	–	–	–

# Precision Measurements @ Higgs Factory


CEPC, 1810.09037



# EFT Description

$$\mathcal{L}_{\text{Eff}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{C_i^{(6)} O_i^{(6)}}{\Lambda^2} + \mathcal{O}(\Lambda^{-4})$$


- EFT: Operators with coefficient suppressed by NP scale  $\Lambda$
- standard tool to study large exp data set
- correlate Higgs, top, EW sector
- model independent
- caveat... See Tim Cohen talk @ Higgs 2021



# EFT

## LHC


Ellis et. al., 2012.02779





## HL-LHC/Higgs factory

Blas et. al., 1907.04311


95% CL reach from the full EFT fit




# Higgs self-coupling



- small CS: 1000 times smaller than Higgs
- 120 K HH events at HL-LHC




**LHC**  $-1.0 < \kappa_\lambda < 6.6$



**HL-LHC**  $0.5 < \kappa_\lambda < 1.5$



# Higgs self-coupling




Higgs@FC WG September 2019

di-Higgs	single-Higgs
HL-LHC 50%	HL-LHC 50%
HE-LHC [10-20]%	HE-LHC 50%
FCC-ee/eh/hh 5%	FCC-ee/eh/hh 25%
LE-FCC 15%	LE-FCC n.a.
FCC-eh <sub>3500</sub> -17+24%	FCC-eh <sub>3500</sub> n.a.
	FCC-ee <sup>4IP</sup> <sub>365</sub> 24%
	FCC-ee <sub>365</sub> 33%
	FCC-ee <sub>240</sub> 49%
ILC <sub>1000</sub> 10%	ILC <sub>1000</sub> 36%
ILC <sub>500</sub> 27%	ILC <sub>500</sub> 38%
	ILC <sub>250</sub> 49%
	CEPC 49%
CLIC <sub>3000</sub> -7%+11%	CLIC <sub>3000</sub> 49%
CLIC <sub>1500</sub> 36%	CLIC <sub>1500</sub> 49%
	CLIC <sub>380</sub> 50%




All future colliders combined with HL-LHC

# New Physics Implication



# MSSM: $m_A$ vs. $\tan\beta$


H. Li, SS, W. Su, J. Yang, 2010.09782




**complementary to  
LHC direct search**

# Tree-level 2HDM fit


## 2HDM, LHC/CEPC fit



# Distinguish different types of 2HDMs



T. Han, S. Li, SS, W. Su, Y. Wu, 2008.05492



# Conclusion

- ◎ The discovery of Higgs is a remarkable triumph in particle physics
- ◎ A light weakly coupled Higgs argues for new physics beyond SM
- ◎ Search for new physics calls for both high precision machine and high energy machine
- ◎ LHC Run II and beyond
  - Higgs precision measurements: mass, width, couplings, CP,...
- ◎ Future Higgs factories: FCC-ee, CEPC, ILC/CLIC...
  - Higgs coupling to sub-percent level
  - Higgs self-coupling 10% @ ILC, CLIC
- ◎ Implication: model independent ( $\kappa$ , EFT), model dependent
- ◎ Higgs precision measurements complementary to direct search/Z pole precision