

A new electron peaks counting algorithm based on a running pulse template

Walaa Elmetenawee

INFN of Bari, Italy

On behalf of Bari and Lecce group

Meeting on cluster counting in drift chambers

7 Apr 2022


Introduction

- ❑ Offline analysis on November test beam data taken with 165 GeV/c muons beams from Nov 2021.
- ❑ Different configurations are used (gas mixture, HV, trigger, track incident angle w.r.t. drift tube, sampling frequency).
- ❑ Dealing with 11 drift tubes having cell sizes of 1 cm, 2 cm, and 3 cm:

- Channels 0,1,2,3 are **Trigger Counters**
- Channels 4,5,6,7,8,9 are the **6 Drift Tubes of 1 cm cell size**
 - ✓ Channel 4 with a wire diameter of 10 micrometer
 - ✓ Channel 5 with a wire diameter of 15 micrometer
 - ✓ Channel 6 and 7 with a wire diameter of 20 micrometer
 - ✓ Channel 8 and 9 with a wire diameter of 25 micrometer

- Channels 10,11,12 are the **3 Drift Tubes of 2 cm cell size**
 - ✓ Channel 10 with a wire diameter of 20 micrometer
 - ✓ Channel 11 with a wire diameter of 25 micrometer
 - ✓ Channel 12 with a wire diameter of 40 micrometer

- Channels 13,14 are the **2 Drift Tubes of 3 cm cell size** (Signal acquisition window is out of the signal range)
 - ✓ Channel 13 with a wire diameter of 25 micrometer
 - ✓ Channel 14 with a wire diameter of 40 micrometer






https://github.com/bdanzi/drifftubes_offline_analysis/

Find Electron Peaks strategy

- Define an **electron pulse** template based on experimental data.
- Raising and falling exponential** over a fixed number of bins (K_{tot}).
- Digitize it ($A(k)$)** according to the data sampling rate.
- Run over K_{tot} bins by comparing it to the subtracted and normalized data (**build a sort of χ^2**).
- Define a cut on χ^2 .
- Subtract the found peak to the signal spectrum.
- Iterate the search.
- Stop when no new peak is found.

k	A(k)
0	0.0
1	0.269
2	1.0
3	0.744
4	0.545
5	0.390
6	0.269
7	0.175
8	0.102
9	0.044
10	0.0



Choose peak function **A(k)** and **Krise** and **Kfall**

Maximum Number of electron **Npeaks** = 500

Store in a temporary buffer the pulse spectrum **Snew(i)** and its standard deviation **sigSnew (i)**

Initialize peak search
jmaxrel bin of presumed peak; **jlast** last possible bin
Search again after peaks subtraction

Loop over all bins of signal spectrum

Define raising or falling ramp **Delta** per bin and the normalization factor of the presumed peak **Norm** with their propagated errors **sigDelta** and **sigNorm**

Loop over **Ktot** bins of presumed signal peak

Peak function **F(k)** and its error **sigF(k)** to be compared to the normalized peak function chosen
Chi square, Chi2, build up

Chi2 condition **chicut** for peak finding

Increment the number of peaks found **Npeaks** and store the corresponding bin position **jmaxrel** and the normalized amplitude **Amp** and **sigAmp**

Loop over **Ktot** bins of found signal peak

Current found peak **subtraction**

Clusterization counting strategy

The strategy make the clusterization of the electron peaks into ionization clusters:

- 1) Association of electron peaks in consecutive bins (difference in time == 1 bin) electrons to a single electron.
- 2) Contiguous electrons peaks which are compatible with the electrons diffusion time (2.5 ns or 3 bins) are considered belonging to the same ionization cluster.
- 3) Position of the clusters is taken as the position of the last electron in the cluster.
- 4) The distributions of the number of clusters must follow a Poisson distribution!



Electron peaks counting & clustering

1 cm drift tubes

Run: run_96.root; Track angle: 30° ; Gas mixture: 90%He10% iC_4H_{10} ; HV = +20

tmpSignal_afterFlt_Ch6_ev51_run_96

tmpSignal_afterFlt_Ch6_ev51_run_96




Electron peaks counting & clustering


2 cm drift tubes

Run: run_96.root; Track angle: 30° ; Gas mixture: 90%He10% i C $_4$ H $_{10}$; HV = +20

tmpSignal_afterFlt_Ch10_ev101_run_96



tmpSignal_afterFlt_Ch10_ev101_run_96



Expected number of Electron Peaks & Clusters

The expected numbers of the electron peaks and clusters are estimated to check our electron peak (cluster) algorithm:

➤ Expected number of electron peaks =
 $\delta \text{ cluster/cm (M.I.P.)} * \text{drift tube size [cm]} * 1.3 \text{ (relativistic rise)} * 1.6 \text{ electrons/cluster} * 1/\cos(\alpha)$

➤ Expected number of clusters =
 $\delta \text{ cluster/cm (M.I.P.)} * \text{drift tube size [cm]} * 1.3 \text{ (relativistic rise)} * 1/\cos(\alpha)$

α = angle of the muon track w.r.t. normal to sense wire.
 $\delta \text{ cluster/cm (mip)}$ = 12 for 90He (18 for 80He) gas mixtures.
 drift tube size = 0.8 for 1 cm (1.8 for 2 cm) drift tube.







Electron peaks counting

N electrons = 32.4

1 cm drift tubes

Expected = 23.1

Run: run_96.root; Track angle: 30^0 ; Gas mixture: 90%He10% iC_4H_{10} ; HV = +20






Electron peaks counting

N electrons = 52.8

2 cm drift tubes

Expected = 51.9


Run: run_96.root; Track angle: 30°; Gas mixture: 90%He10%iC4H10; HV = +20



Electron peaks counting


1 cm drift tubes

Electrons Finding Efficiency 1 cm cell size Drift Tubes



2 cm drift tubes

Electrons Finding Efficiency 2 cm cell size Drift Tubes



- **Electrons overcounting** due to fake electron peaks in adjacent bins.
- (easily corrected in the clusterization algorithm)
- **Undercounting for 2 cm $\alpha < 30^\circ$** due to space charge effects.







Electron Clustering

N Clusters = 12.6

1 cm drift tubes

Expected = 14.4

Run: run_96.root; Track angle: 30° ; Gas mixture: 90%He10% iC_4H_{10} ; HV = +20



*The Fit applied is Poisson fit




Electron Clustering

N Clusters = 21.5

2 cm drift tubes

Expected = 32.4

Run: run_96.root; Track angle: 30° ; Gas mixture: 90%He10% i C₄H₁₀; HV = +20




*The Fit applied is Gauss fit

Electron Clustering


1 cm drift tubes

Clusters Finding Efficiency 1 cm cell size Drift Tubes



2 cm drift tubes



Clusters Finding Efficiency 2 cm cell size Drift Tubes




- Undercounting at $\alpha < 30^\circ$ due to space charge effects.
- Undercounting at $\alpha > 45^\circ$ due to high electron peaks density.
- 10 % average inefficiency for 1 cm drift tubes ($\alpha < 30^\circ$ & $\alpha > 45^\circ$)
- 30% average inefficiency for 2 cm drift tubes (electron inefficiency) ($\alpha < 30^\circ$ & $\alpha > 45^\circ$)

Testing different templates

Template 1





Template 2



Template 3

k	A(k)
0	0,000
1	0,102
2	0,269
3	0,545
4	1,000
5	0,864
6	0,744
7	0,638
8	0,545
9	0,463
10	0,390
11	0,326
12	0,269
13	0,219
14	0,175
15	0,136
16	0,102
17	0,071
18	0,044
19	0,021
20	0,000




Comparison between the different templates

1 cm drift tubes

Template 1


Template 3

Clusters Finding Efficiency 1 cm cell size Drift Tubes




Template 2

Clusters Finding Efficiency 2 cm cell size Drift Tubes



Clusters Finding Efficiency 1 cm cell size Drift Tubes




Comparison between the different templates

Template 1


2 cm drift tubes

Template 3

Clusters Finding Efficiency 2 cm cell size Drift Tubes




Clusters Finding Efficiency 2 cm cell size Drift Tubes



Template 2

Clusters Finding Efficiency 2 cm cell size Drift Tubes



Decided to keep with Template 1

Comparison between DERIV & RTA Algorithms


1 cm drift tubes

- Comparison between the derivatives (DERIV) and Running Template Algorithm (RTA): "Electron peaks counting & clustering"

Run: run_96.root; Track angle: 30° ; Gas mixture: 90%He10% iC_4H_{10} ; HV = +20


RTA algorithm

tmpSignal_afterFlt_Ch6_ev51_run_96



DERIV algorithm

tmpSignal_afterFlt_Ch6_ev51_run_96



Comparison between DERIV & RTA Algorithms


2 cm drift tubes

- Comparison between the derivatives (DERIV) and Running Template Algorithm (RTA):
"Electron peaks counting & clustering"

Run: run_96.root; Track angle: 30° ; Gas mixture: 90%He10% iC_4H_{10} ; HV = +20


RTA algorithm

tmpSignal_afterFlt_Ch10_ev101_run_96



DERIV algorithm

tmpSignal_afterFlt_Ch10_ev101_run_96




Comparison between DERIV & RTA Algorithms

1 cm drift tubes

- Comparison between the derivatives (DERIV) and Running Template Algorithm (RTA):
"Electron peaks counting"


RTA algorithm

Electrons Finding Efficiency 1 cm cell size Drift Tubes



DERIV algorithm

Electrons Finding Efficiency 1 cm cell size Drift Tubes




Comparison between DERIV & RTA Algorithms

2 cm drift tubes

- Comparison between the derivatives (DERIV) and Running Template Algorithm (RTA):
"Electron peaks counting"


RTA algorithm

Electrons Finding Efficiency 2 cm cell size Drift Tubes



DERIV algorithm

Electrons Finding Efficiency 2 cm cell size Drift Tubes




Comparison between DERIV & RTA Algorithms

1 cm drift tubes

- Comparison between the derivatives (DERIV) and Running Template Algorithm (RTA):
"Electrons cluster density"


RTA algorithm

Epc 1 cm cell size Drift Tubes



DERIV algorithm

Epc 1 cm cell size Drift Tubes




Comparison between DERIV & RTA Algorithms

2 cm drift tubes

- Comparison between the derivatives (DERIV) and Running Template Algorithm (RTA):
"Electrons cluster density"


RTA algorithm

Epc 2 cm cell size Drift Tubes



DERIV algorithm

Epc 2 cm cell size Drift Tubes




Comparison between DERIV & RTA Algorithms

1 cm drift tubes

- Comparison between the derivatives (DERIV) and Running Template Algorithm (RTA):
"Electrons Clustering"


RTA algorithm

Clusters Finding Efficiency 1 cm cell size Drift Tubes



DERIV algorithm

Clusters Finding Efficiency 1 cm cell size Drift Tubes



~ 5% inefficiency in the RTA in comparison with DERIV algorithm


Comparison between DERIV & RTA Algorithms

2 cm drift tubes

- Comparison between the derivatives (DERIV) and Running Template Algorithm (RTA):
"Electrons Clustering"


RTA algorithm

Clusters Finding Efficiency 2 cm cell size Drift Tubes



DERIV algorithm

Clusters Finding Efficiency 2 cm cell size Drift Tubes



~ 10% inefficiency in the RTA in comparison with DERIV algorithm

Conclusions

- Further optimization will be done to the Running Template Algorithm (RTA) in order to recover the inefficiency observed.
- The application of the two different algorithms will be very useful for understanding the pathologies of both algorithms, therefore, it will be extremely useful to have a third algorithm like the one being developed at IHEP with NN.