

Next-to-next-to-leading order matching of B_c and B_c^* decay constants

**Reporter: Tao Wei
2022.11.24**

**Based on:
Wei Tao, Ruilin Zhu, Zhen-Jun Xiao, arXiv:2209.15521**

Outline

1

Introduction

2

Calculation Procedure

3

Results

4

Summary

Introduction

- **Beauty-charmed meson family discovered in particle physics experiment**
 - $B_c(1S)$, CDF collaboration, 1998
 - $B_c(2S)$, ATLAS collaboration, 2014
 - $B_c^*(2S)$, CMS and LHCb collaborations, 2019

- **Difficulty for experiment measurement**
 - composed of two different heavy flavor quarks
 - the ground state $B_c(1S)$ only weak decays
 - Absolute branching ratios are hardly measured

Introduction

➤ Theoretical investigation

- Need study decay constants to obtain leptonic branching ratios
- Decay constants are essentially nonperturbative and universal
- But lattice QCD studies lesser due to doubly heavy flavors

➤ Nonrelativistic QCD (NRQCD) effective theory

scale : $M \gg M v \sim \Lambda_{\text{QCD}}$

perturbative nonperturbative

- Decay constants \sim Short-distance perturbative matching coefficients \times long-distance nonperturbative NRQCD matrix elements (LDMEs)
- Systematical calculation order by order

Introduction

➤ Review high order calculation for B_c and B_c^* decay constants with NRQCD

- systematical study of B_c at the leading order (LO) [C.H.Chang,Y.Q.Chen,PRD(1994)]
- NLO for B_c [E.Braaten,S.Fleming,PRD(1995)]
- NLO for B_c^* [D.S.Hwang,S.Kim,PRD(1999)]
- High order relativistic corrections resummed [J.Lee,W.Sang,S.Kim,JHEP(2011)]
- Approximate NNLO for B_c [A.I.Onishchenko,O.L.Veretin,EPJC(2007)]
- Full analytical NNLO for B_c [L.B.Chen,C.F.Qiao,PLB(2015)]
- NNNLO for B_c [F.Feng,et al.,2208.04302]
- NNLO for B_c^* [W.Tao,R.Zhu,Z.J.Xiao,2209.15521]
- NNNLO for B_c^* [W.Sang,H.F.Zhang,M.Z.Zhou,2210.02979]

Outline

- 1 Introduction
- 2 Calculation Procedure
- 3 Results
- 4 Summary

Definition

➤ QCD definition for decay constants

$$\langle 0 | \bar{b} \gamma^\mu \gamma_5 c | B_c(P) \rangle = i f_{B_c}^a P^\mu,$$

$$\langle 0 | \bar{b} \gamma_5 c | B_c(P) \rangle = i f_{B_c}^p m_{B_c},$$

$$\langle 0 | \bar{b} \gamma^\mu c | B_c^*(P, \varepsilon) \rangle = f_{B_c^*}^v m_{B_c^*} \varepsilon^\mu,$$

$a \equiv (a, 0)$:timelike component of axial current

p :pseudoscalar current

v :vector current

$$f_{B_c}^a \equiv f_{B_c}^p$$

➤ NRQCD definition for decay constants

$$f_{B_c}^p = \sqrt{\frac{2}{m_{B_c}}} \underbrace{C_p(m_b, m_c, \mu_f)}_{\text{matching coefficients}} \underbrace{\langle 0 | \chi_b^\dagger \psi_c | B_c(\mathbf{P}) \rangle}_{\text{NRQCD LDMEs}} (\mu_f) + O(v^2)$$

$$f_{B_c^*}^v = \sqrt{\frac{2}{m_{B_c^*}}} \underbrace{C_v(m_b, m_c, \mu_f)}_{\text{matching coefficients}} \underbrace{\langle 0 | \chi_b^\dagger \sigma \cdot \varepsilon \psi_c | B_c^*(\mathbf{P}) \rangle}_{\text{NRQCD LDMEs}} (\mu_f) + O(v^2)$$

Matching&Renormalization

➤ Perturbative matching formulae

$$Z_J Z_{2,b}^{\frac{1}{2}} Z_{2,c}^{\frac{1}{2}} \Gamma_J = C_J \tilde{Z}_J^{-1} \tilde{Z}_{2,b}^{\frac{1}{2}} \tilde{Z}_{2,c}^{\frac{1}{2}} \tilde{\Gamma}_J$$

Z_J :QCD on-shell current renormalization constants

$$Z_a = Z_v = 1, Z_p = \frac{m_b Z_{m,b} + m_c Z_{m,c}}{m_b + m_c}$$

\tilde{Z}_J :NRQCD $\overline{\text{MS}}$ current renormalization constants

Z_2/Z_m :QCD on-shell field/mass renormalization constants

\tilde{Z}_2 : NRQCD on-shell field renormalization constants

$$\tilde{Z}_{2,b} = \tilde{Z}_{2,c} = 1$$

$\Gamma_J/\tilde{\Gamma}_J$:unrenormalized QCD/NRQCD current vertex function

$$\tilde{\Gamma}_J=1$$

Matching & Renormalization

➤ NRQCD current renormalization constants and anomalous dimensions

$$\tilde{Z}_J = 1 - \left(\frac{\alpha_s^{(n_l)}(\mu_f)}{\pi} \right)^2 \frac{\gamma_J^{(2)}(x)}{4\epsilon} + \mathcal{O}(\alpha_s^3),$$

$$\gamma_J = \frac{d \ln \tilde{Z}_J}{d \ln \mu_f} = \frac{-2 \partial \tilde{Z}_J^{(1)}}{\partial \ln \alpha_s^{(n_l)}(\mu_f)} = \left(\frac{\alpha_s^{(n_l)}(\mu_f)}{\pi} \right)^2 \gamma_J^{(2)}(x) + \mathcal{O}(\alpha_s^3)$$

$$\gamma_p^{(2)}(x) = -\pi^2 \left(\frac{C_F C_A}{2} + \frac{(1 + 6x + x^2) C_F^2}{2(1 + x)^2} \right),$$

$$\gamma_v^{(2)}(x) = -\pi^2 \left(\frac{C_F C_A}{2} + \frac{(3 + 2x + 3x^2) C_F^2}{6(1 + x)^2} \right)$$

Matching & Renormalization

➤ Matching and Calculation of α_s

- Decoupling $\alpha_s^{(n_f=n_b+n_c+n_l)}(\mu)$ to $\alpha_s^{(n_l)}(\mu)$

$$\alpha_s^{(n_f)}(\mu) = \alpha_s^{(n_l)}(\mu) \left(1 + \frac{\alpha_s^{(n_l)}(\mu)}{\pi} T_F \left(\frac{n_b}{3} \ln \frac{\mu^2}{m_b^2} + \frac{n_c}{3} \ln \frac{\mu^2}{m_c^2} + O(\epsilon) \right) + O(\alpha_s^2) \right)$$

- Renormalization group running equations

$$\alpha_s^{(n_l)}(\mu_f) = \left(\frac{\mu}{\mu_f} \right)^{2\epsilon} \alpha_s^{(n_l)}(\mu),$$

μ : renormalization scale
 μ_f : NRQCD factorization scale

$$\alpha_s^{(n_l)}(\mu) = \frac{4\pi}{\beta_0^{(n_l)} \ln \frac{\mu^2}{\Lambda_{QCD}^{(n_l)2}}} \left(1 - \frac{\beta_1^{(n_l)} \ln \ln \frac{\mu^2}{\Lambda_{QCD}^{(n_l)2}}}{\beta_0^{(n_l)2} \ln \frac{\mu^2}{\Lambda_{QCD}^{(n_l)2}}} \right)$$
















Matching & Renormalization

➤ Renormalization at NNLO

- two loop diagrams
- tree diagram inserted with one α_s^2 -order counter-term vertex
- tree diagram inserted with two α_s -order counter-term vertexes (vanishing)
- one loop diagram inserted with one α_s -order counter-term vertex

Diagrams

➤ Tree,1loop,2loop



Calculation steps

➤ Higher order calculation steps

- **FeynCalc** obtains diagrams and corresponding amplitudes,
\$Apart decomposes every amplitude into several Feynman integral families
- **FIRE / Kira / FiniteFlow** reduces every Feynman integral family to master integral family
- **Kira+FIRE+Mathematica code** reduces all of master integral families to the minimal master integral families
- **AMFlow** with **Kira/FiniteFlow** calculates the minimal master integral families one by one
- Renormalization

Outline

- 1 Introduction
- 2 Calculation Procedure
- 3 Results
- 4 Summary

Matching coefficients results

➤ Matching coefficient formula

$$\begin{aligned} C_J(\mu_f, \mu, m_b, x) = & 1 + \frac{\alpha_s^{(n_f)}(\mu)}{\pi} C_J^{(1)}(x) \\ & + \left(\frac{\alpha_s^{(n_f)}(\mu)}{\pi} \right)^2 \left(C_J^{(1)}(x) \frac{\beta_0^{(n_f)}}{4} \ln \frac{\mu^2}{m_b^2} + \frac{\gamma_J^{(2)}(x)}{2} \ln \frac{\mu_f^2}{m_b^2} \right. \\ & + C_F^2 C_J^{FF}(x) + C_F C_A C_J^{FA}(x) + C_F T_F n_l C_J^{FL}(x) + C_F T_F C_J^{FH}(x) \Big) \\ & + O(\alpha_s^3). \end{aligned}$$

- $x = \frac{m_c}{m_b}$





- NLO matching coefficients:

$$C_p^{(1)}(x) = \frac{3}{4} C_F \left(\frac{x-1}{x+1} \ln x - 2 \right)$$

$$C_v^{(1)}(x) = \frac{3}{4} C_F \left(\frac{x-1}{x+1} \ln x - \frac{8}{3} \right)$$

Matching coefficients results

➤ x dependence for C_J^{FF} , C_J^{FA} , C_J^{FL} , and C_J^{FH}





Matching coefficients results

➤ numerical results for matching coefficients

- $\mu_f \in [1.5, 1.2, 1] \text{ GeV}$, $\mu \in [6.25, 4.75, 3] \text{ GeV}$, $m_b \in [5.25, 4.75, 4.25] \text{ GeV}$, $m_c \in [2, 1.5, 1] \text{ GeV}$

	LO	NLO	NNLO
C_p	1	$0.9117^{+0.0072+0.0061-0.0156}_{-0-0.0160-0.0064+0.0263}$	$0.7897^{+0.0206+0.0119+0.0149}_{-0.0310-0.0482-0.0133-0.0141}$
C_v	1	$0.8697^{+0.0107+0.0061-0.0156}_{-0-0.0236-0.0064+0.0263}$	$0.7363^{+0.0230+0.0106+0.0117}_{-0.0234-0.0526-0.0117-0.0121}$

- μ dependence for matching coefficients



Decay constants results

➤ Bc and Bc* decay constants

- LDMEs $\langle 0 | \chi_b^\dagger \sigma \cdot \varepsilon \psi_c | B_c^*(\mathbf{P}) \rangle \approx \langle 0 | \chi_b^\dagger \psi_c | B_c(\mathbf{P}) \rangle \approx \sqrt{2N_c} \psi_{B_c}(0)$.
- $|\psi_{B_c}(0)|^2 \in [0.13, 0.12, 0.10] \text{GeV}^3$, $\mu_f \in [1.5, 1.2, 1] \text{GeV}$,
 $\mu \in [6.25, 4.75, 3] \text{GeV}$, $m_b \in [5.25, 4.75, 4.25] \text{GeV}$, $m_c \in [2, 1.5, 1] \text{GeV}$

	$\frac{f_{B_c}^p}{10^{-1} \text{GeV}}$	$\frac{f_{B_c^*}^v}{10^{-1} \text{GeV}}$
LO	$4.79^{+0.20+0+0+0+0}_{-0.42-0-0-0-0}$	$4.78^{+0.20+0+0+0+0}_{-0.42-0-0-0-0}$
NLO	$4.37^{+0.18+0+0.03+0.03-0.07}_{-0.38-0-0.08-0.03+0.13}$	$4.15^{+0.17+0+0.05+0.03-0.07}_{-0.36-0-0.11-0.03+0.13}$
NNLO	$3.78^{+0.15-0.15+0.10+0.06+0.07}_{-0.33+0.12-0.23-0.06-0.07}$	$3.52^{+0.14-0.11+0.11+0.05+0.06}_{-0.31+0.09-0.25-0.06-0.06}$

Decay widths & branching ratios

➤ Bc and Bc* leptonic decay widths and branching ratios

- Formulae

$$\Gamma(B_c^+ \rightarrow l^+ + \nu_l) = \frac{|V_{bc}|^2}{8\pi} G_F^2 m_{B_c} m_l^2 \left(1 - \frac{m_l^2}{m_{B_c}^2}\right)^2 f_{B_c}^p{}^2,$$

$$\Gamma(B_c^{*+} \rightarrow l^+ + \nu_l) = \frac{|V_{bc}|^2}{12\pi} G_F^2 m_{B_c^*}^3 \left(1 - \frac{m_l^2}{m_{B_c^*}^2}\right)^2 \left(1 + \frac{m_l^2}{2m_{B_c^*}^2}\right) f_{B_c^*}^v{}^2$$

- Decay widths

	$\frac{\Gamma(B_c^+ \rightarrow e^+ + \nu_e)}{10^{-21} \text{ GeV}}$	$\frac{\Gamma(B_c^{*+} \rightarrow e^+ + \nu_e)}{10^{-13} \text{ GeV}}$
LO	$3.39^{+0.28+0+0+0+0}_{-0.56-0-0-0-0}$	$3.45^{+0.29+0+0+0+0}_{-0.57-0-0-0-0}$
NLO	$2.82^{+0.23+0+0.04+0.04-0.10}_{-0.47-0-0.10-0.04+0.16}$	$2.61^{+0.22+0+0.06+0.04-0.09}_{-0.43-0-0.14-0.04+0.16}$
NNLO	$2.11^{+0.18-0.16+0.11+0.06+0.08}_{-0.35+0.14-0.25-0.07-0.07}$	$1.87^{+0.16-0.12+0.12+0.05+0.06}_{-0.31+0.10-0.26-0.06-0.06}$
	$\frac{\Gamma(B_c \rightarrow \mu^+ + \nu_\mu)}{10^{-16} \text{ GeV}}$	$\frac{\Gamma(B_c^* \rightarrow \mu^+ + \nu_\mu)}{10^{-13} \text{ GeV}}$
LO	$1.45^{+0.12+0+0+0+0}_{-0.24-0-0-0-0}$	$3.45^{+0.29+0+0+0+0}_{-0.57-0-0-0-0}$
NLO	$1.20^{+0.10+0+0.02+0.02-0.04}_{-0.20-0-0.04-0.02+0.07}$	$2.61^{+0.22+0+0.06+0.04-0.09}_{-0.43-0-0.14-0.04+0.16}$
NNLO	$0.90^{+0.08-0.07+0.05+0.03+0.03}_{-0.15+0.06-0.11-0.03-0.03}$	$1.87^{+0.16-0.12+0.12+0.05+0.06}_{-0.31+0.10-0.26-0.06-0.06}$

Decay widths & branching ratios

- Decay widths

	$\frac{\Gamma(B_c^+ \rightarrow \tau^+ + \nu_\tau)}{10^{-14} \text{GeV}}$	$\frac{\Gamma(B_c^{*+} \rightarrow \tau^+ + \nu_\tau)}{10^{-13} \text{GeV}}$
LO	$3.47^{+0.29+0+0+0+0}_{-0.58-0-0-0-0}$	$3.04^{+0.25+0+0+0+0}_{-0.51-0-0-0-0}$
NLO	$2.88^{+0.24+0+0.05+0.04-0.10}_{-0.48-0-0.10-0.04+0.17}$	$2.30^{+0.19+0+0.06+0.03-0.08}_{-0.38-0-0.12-0.03+0.14}$
NNLO	$2.16^{+0.18-0.17+0.11+0.07+0.08}_{-0.36+0.14-0.26-0.07-0.08}$	$1.65^{+0.14-0.10+0.10+0.05+0.05}_{-0.27+0.09-0.23-0.05-0.05}$

- Decay branching ratios

	$e^+ \nu_e$	$\mu^+ \nu_\mu$	$\tau^+ \nu_\tau$
B_c	$\left(1.64^{+0.44}_{-0.71}\right) \times 10^{-9}$	$\left(7.00^{+1.89}_{-3.01}\right) \times 10^{-5}$	$\left(1.68^{+0.45}_{-0.72}\right) \times 10^{-2}$
B_c^*	$\left(2.34^{+0.61}_{-1.01}\right) \times 10^{-6}$	$\left(2.34^{+0.61}_{-1.01}\right) \times 10^{-6}$	$\left(2.06^{+0.54}_{-0.89}\right) \times 10^{-6}$

Outline

- 1 Introduction
- 2 Calculation Procedure
- 3 Results
- 4 **Summary**

Summary

- Verify B_c decay constant from pseudoscalar current is identical with the B_c decay constant from axial-vector current
- Obtain the novel anomalous dimension for the flavor-changing heavy quark vector current
- Obtain NNLO result for the decay constant of B_c^*
- The obtained branching ratio of $B_c^{*+} \rightarrow \mu^+ + \nu_\mu$ isn't small, which can be a good channel to detect B_c^*

Thank you!