



$H \rightarrow \gamma\gamma$ fiducial and differential cross-section measurements with full Run2 dataset at ATLAS

Fábio Lucio Alves (Nanjing University)
on behalf of the ATLAS Collaboration

The 8th China LHC Physics Workshop (CLHCP2022)
November 23-27, Nanjing, China

Introduction

◎ It's 10 years since the Higgs boson discovery (2012):

- ◎ Measurements of its properties using full Run2 dataset (2015-2018) are in excellent agreement with the SM.


◎ Inclusive and Differential Fiducial Higgs measurements:

- ◎ Test the SM Higgs boson properties and probe for BSM contributions:
 - ◎ measured differential cross-sections distributions are compared to state-of-the-art SM predictions
- ◎ Less model-dependent measurements: small extrapolations and SM assumptions
- ◎ $H \rightarrow \gamma\gamma$ channel measurements [JHEP08\(2022\)027](#):
 - ◎ No separation of production modes is aimed
 - ◎ Observables sensitive to New Physics, contributions of different Higgs boson production modes, CP-properties and QCD effects
 - ◎ Cross-sections measured are used in the interpretation via EFT theory and setting limits on Yukawa bottom- and charm-quarks using di-photon p_T shape information

July 2012



July 2022



$H \rightarrow \gamma\gamma$ analysis in a nutshell

● **Signature:** two reconstructed isolated photons

▶ **diphoton vertex:** NN algorithm, improves mass resolution

▶ **kinematic selections:**

▶ $pT(\gamma_1) > 0.35m_{\gamma\gamma}$; $pT(\gamma_2) > 0.25m_{\gamma\gamma}$

▶ $|\eta| < 2.37$ (exclude 1.37-1.52 region)

▶ Jets: $pT > 30$ GeV, $|\eta| < 4.4$

▶ $105 \text{ GeV} < m_{\gamma\gamma} < 160 \text{ GeV}$

▶ **Signal Modelling:**

▶ double-sided Crystal Ball function: MC simulation

● **Background sources and modelling:**


▶ SM $\gamma\gamma$ production (irreducible) and $\gamma - jet$ and $jet - jet$ (reducible ones):

▶ built background-only templates, GPR approach to smooth statistical fluctuations in low-yields templates

▶ choice of the background model: signal+background fits to $m_{\gamma\gamma}$
background-only templates:


▶ function's choice uncertainty ('spurious signal')

Fits to $m_{\gamma\gamma}$ to extract the signal yields



Measurement methodology

- ▶ **Fiducial region:** defined to closely match the detector-level analysis and object selections
- ▶ **Differential fiducial cross-section** are measured in bins of the studied observable (bin i of a variable x)
 - ✓ N_i^{sig} (**measured signal yield**): extracted signal events in data
 - ✓ Δx (**bin width**): choice based on significance (close to or greater than 2σ) and minimize migrations
 - ✓ c_i (**correction factor**): accounts for detector inefficiencies and resolutions effects as well as migrations in and out of fiducial region
 - ▶ MC response matrix




$$\frac{d\sigma_i}{dx} = \frac{N_i^{sig}}{c_i \Delta x \mathcal{L}}$$

Differential fiducial cross-section

Fiducial cross-section measurement

Fiducial cross-sections measurements in regions enriched in various production modes



Diphoton fiducial region

Source	Uncertainty [%]
Statistical uncertainty	7.5
Systematic uncertainties	6.4
Background modelling (spurious signal)	3.8
Photon energy scale & resolution	3.6
Photon selection efficiency	2.6
Luminosity	1.8
Pile-up modelling	1.4
Trigger efficiency	1.0
Theoretical modelling	0.4
Total	9.8

Diphoton fiducial cross-section =
 $67 \pm 5(\text{stat.}) \pm 4(\text{sys.}) \text{ fb}$
SM = $64 \pm 4 \text{ fb}$

- Measurement is statistical dominated
- Background modelling is the largest systematic uncertainty



Differential cross-section vs $p_T^{\gamma\gamma}$ and $|y_{\gamma\gamma}|$

default MC: Powheg
NNLOPS, normalized to N3LO
(QCD)+NLO(EW)

- **Low $p_T^{\gamma\gamma}$** : sensitive to bottom- and charm-quark Yukawa couplings
- **High $p_T^{\gamma\gamma}$** : top quark coupling and BSM effects

- $|y_{\gamma\gamma}|$ is sensitive to light-quark Yukawa coupling and gluon distribution in the proton

pT(H) > 350 GeV: sensitivity to BSM effects




Good agreement with the SM predictions within the uncertainties

Cross-section vs N_{jets} and N_{b-jets}


- Inclusive N_{jets} ($p_T > 30$ GeV and $|y| < 4.4$):
 - sensitive to different Higgs boson production modes and QCD modelling (ggF)
- N_{b-jets} : at least 1 central jet and veto on electrons/muons (suppress ttH contribution)
 - sensitive to Higgs production in association with heavy particles

Good agreement with the SM predictions

Jet-related systematic reaching up to 24% in highest jet multiplicities



Inclusive N_{jets}
 $p(\chi^2) = 95\%$
 (default)



N_{b-jets}
 $p(\chi^2) = 60\%$
 (default)

Differential cross-section vs p_T^{j1} , m_{jj} and $\Delta\phi_{jj}$

- p_T^{j1} : same range as $p_T^{\gamma\gamma}$ but within coarser bins at low p_T




► $p(\chi^2) = 78\%$ (default MC)

- m_{jj} : highest bin is most sensitive to the VBF production mechanism

► $p(\chi^2) = 79\%$ (default MC)

- $\Delta\phi_{jj}$: sensitive to CP properties of the Higgs boson



► $p(\chi^2) = 91\%$ (default MC)



Good agreement with the SM predictions within the uncertainties

Double Differential cross-section vs $p_T^{\gamma\gamma}$ vs $|y_{\gamma\gamma}|$ and $p_T^{\gamma\gamma}$ vs $\tau_{C,j1}$

- 2D observables provides further look into the Higgs boson properties and correlations between the observables



$p_T^{\gamma\gamma}$ vs $|y_{\gamma\gamma}|$
 $p(\chi^2) = 75\%$ (default MC)

Good agreement with the SM predictions within the uncertainties

$p_T^{\gamma\gamma}$ vs $\tau_{C,j1}$
 $p(\chi^2) = 39\%$ (default MC)

Interpretation using the p_T spectrum information

shape only

shape+XS+BR

- $p_T^{\gamma\gamma}$ spectrum provides indirect measurement of the c- and b-quarks Yukawa couplings (κ_c and κ_b):
- shape-only and shape+normalization information are used for the fitting

- Limits on κ_c and κ_b are set from a profile likelihood method:
 - Most sensitive region of $p_T^{\gamma\gamma}$ spectrum < 200 GeV
 - Stronger constraints from shape+normalization
 - Shape+normalization constrain on κ_b is comparable to direct searches while κ_c provides stronger constraints (a factor of 3 more stringent for the observed result and a factor of 4 for the expected result)

Fit set-up	κ	Observed 95% CL	Expected 95% CL
Shape-only	κ_c	$[-13.0, 18.9]$	$[-10.1, 17.3]$
	κ_b	$[-3.7, 10.4]$	$[-2.6, 8.1]$
Shape+normalisation (with branching ratio variations)	κ_c	$[-2.7, 2.6]$	$[-3.1, 3.2]$
	κ_b	$[-1.2, -0.8] \cup [0.8, 1.1]$	$[-1.2, -0.9] \cup [0.8, 1.2]$

Direct search limits on κ_c @ 95%CL:

observed: $|\kappa_c| < 8.5$

expected: $|\kappa_c| < 12.4$


ATLAS and CMS Run2 brief comparisons

- Comparison between the two analysis (diphoton fiducial region):
 - ATLAS/CMS unfolding: matrix response
 - S+B unbinned fit to $m_{\gamma\gamma}$ (ATLAS); S+B binned fit to $m_{\gamma\gamma}$ (CMS)
 - Signal Model: Double Sided Crystall Ball (ATLAS); Sum of Gaussians (CMS)
 - Background modelling: from spurious signal studies (ATLAS); Discrete profiling method (CMS)
 - Uncertainties:
 - statistical component: $\sim 7.0\%$ (similar)
 - Systematic component: spurious signal is about 3.8% (ATLAS) and 0%(included in the statistical error) in CMS

CMS, submit. to JHEP


ATLAS

Diphoton fiducial cross-section =
 $67 \pm 5(\text{stat.}) \pm 4(\text{sys.}) \text{ fb}$
SM = $64 \pm 4 \text{ fb}$



CMS

Diphoton fiducial cross-section =
 $73.40^{+6.1}_{-5.9} \text{ fb}$
SM = $75.44 \pm 4.1 \text{ fb}$



Summary

✓ *$H \rightarrow \gamma\gamma$ channel fiducial and differential cross sections measurements using full Run2 dataset in ATLAS experiment have been presented:*

- Higgs boson properties and probe to new physics contributions in many observables exploring *Higgs kinematic and jet-kinematic activity in the events*
- *Very good agreement between the measurements and SM predictions:*
 - ▶ Statistical uncertainty still the dominant uncertainty source
- *Interpretation of the measurements using $p_T(H)$ shape information to set constraints on charm- and bottom-quarks Yukawa couplings (κ_c and κ_b):*
 - ▶ No significant BSM contributions are observed
 - ▶ κ_b limits are comparable to the direct searches
 - ▶ Stronger limits on κ_c are set compared to direct searches

Stay tuned for Run3 news!

谢谢大家的关注!

Back-up slides


Fraction of signal process in fiducial regions

◎ Baseline fiducial region (particle level):

- ▶ $E_T/m_{\gamma\gamma} > 0.35$; $E_T/m_{\gamma\gamma} > 0.25$; isolation requirements;
 $|\eta| < 2.37$ excluding transition region

- ▶ **Sub-sets of the baseline fiducial region (sensitive to different Higgs boson production modes):** selections are applied on the electrons, muons, jets and MET

- ◎ **VBF-enhanced region:** at least two jets, $m_{jj} \geq 600 \text{ GeV}$, $|\Delta y_{jj}| \geq 3.5$
- ◎ **$N_{\text{lepton}} \geq 1$ region:** additional charged lepton with $p_T > 15 \text{ GeV}$. Sensitive to VH (V= W/Z), tH or ttH production modes
- ◎ **High MET region:** large MET $> 80 \text{ GeV}$ and $p_T^{\gamma\gamma} > 80 \text{ GeV}$. Sensitive to VH and ttH production mechanisms and BSM effects (WIMPS)
- ◎ **ttH-enhanced region:** at least 1 b-jets/no leptons/4jets or at least 1 lepton and at least 3 jets





Differential cross-section vs $\tau_{C,j1}$ and $\sum \tau_{C,j}$

$$\tau_{C,j} = \frac{\sqrt{p_T^2 + m^2}}{2\cosh(y_j - y_{\gamma\gamma})}$$

m is the jet mass and y_j its rapidity

- ▶ Beam-thrust event-shape variable (hadronic observable): boost-invariant along the beam axis
- ▶ $\tau_{C,j1}$ is the highest value of $\tau_{C,j}$ among all jets in the event while $\sum \tau_{C,j}$ is the scalar sum of τ for all jets with $\tau > 5$ GeV





$\tau_{C,j1}$
 $p(\chi^2) = 27\%$
(default)

$\sum \tau_{C,j}$
 $p(\chi^2) = 39\%$
(default)

Good agreement with the
SM predictions

Systematic uncertainties in $p_T^{\gamma\gamma}$ and N_{jets} bins

- ▶ Systematics uncertainties sources on the signal and background modeling
- ▶ Experimental and theoretical uncertainties uncertainties




Binning choice for different observables

Variable	Bin Edges	N_{bins}
$p_{\text{T}}^{\gamma\gamma}$	0, 5, 10, 15, 20, 25, 30, 35, 45, 60, 80, 100, 120, 140, 170, 200, 250, 300, 450, 650, 13000	20
$ y_{\gamma\gamma} $	0, 0.15, 0.3, 0.45, 0.6, 0.75, 0.9, 1.2, 1.6, 2.0, 2.5	10
$p_{\text{T}}^{\gamma 1}/m_{\gamma\gamma}$	0.35, 0.45, 0.5, 0.55, 0.6, 0.65, 0.75, 0.85, 0.95, 10	9
$p_{\text{T}}^{\gamma 2}/m_{\gamma\gamma}$	0.25, 0.35, 0.4, 0.45, 0.5, 0.55, 0.65, 0.75, 0.85, 10	9
N_{jets}	0, 1, 2, ≥ 3	4
$N_{b\text{-jets}}$	$N_{\text{jets}}^{\text{central}} = 0$ or $N_{\text{lep}} > 0$, $N_{b\text{-jets}} = 0, \geq 1$	3
$p_{\text{T}}^{j_1}$	30, 60, 90, 120, 350, 13000	5
H_{T}	30, 60, 140, 200, 500, 13000	5
$p_{\text{T}}^{\gamma\gamma j}$	0, 30, 60, 120, 13000	4
$m_{\gamma\gamma j}$	120, 220, 300, 400, 600, 900, 13000	6
$\tau_{C,j1}$	0, 5, 15, 25, 40, 13000	5
$\sum \tau_{C,j}$	5, 15, 25, 40, 80, 13000	5
$p_{\text{T}}^{\gamma\gamma, \text{jet veto } 30 \text{ GeV}}$	0, 5, 10, 15, 20, 30, 40, 50, 100, 13000	9
$p_{\text{T}}^{\gamma\gamma, \text{jet veto } 40 \text{ GeV}}$	0, 5, 10, 15, 20, 30, 40, 50, 60, 100, 13000	10
$p_{\text{T}}^{\gamma\gamma, \text{jet veto } 50 \text{ GeV}}$	0, 5, 10, 15, 20, 30, 40, 50, 60, 70, 100, 13000	11
$p_{\text{T}}^{\gamma\gamma, \text{jet veto } 60 \text{ GeV}}$	0, 5, 10, 15, 20, 30, 40, 50, 60, 70, 80, 100, 13000	12
m_{jj}	0, 120, 450, 3000, 13000	4
$\Delta\phi_{jj}$	$-\pi, -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi$	4
$\pi - \Delta\phi_{\gamma\gamma,jj} $	0, 0.15, 0.65, π	3
$p_{\text{T},\gamma\gamma jj}$	0, 30, 60, 120, 13000	4
VBF-enhanced: $p_{\text{T}}^{j_1}$	30, 120, 13000	2
VBF-enhanced: $\Delta\phi_{jj}$	$-\pi, -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi$	4
VBF-enhanced: $ \eta^* $	0, 1, 2, 10	3
VBF-enhanced: $p_{\text{T},\gamma\gamma jj}$	0, 30, 13000	2

Binning choice for different observables

Variable	Bin Edges		N_{bins}
$p_T^{\gamma\gamma}$ vs $ y_{\gamma\gamma} $	$0.0 < y_{\gamma\gamma} < 0.5$	$p_T^{\gamma\gamma}: 0, 45, 120, 350$	12
	$0.5 < y_{\gamma\gamma} < 1.0$	$p_T^{\gamma\gamma}: 0, 45, 120, 350$	
	$1.0 < y_{\gamma\gamma} < 1.5$	$p_T^{\gamma\gamma}: 0, 45, 120, 350$	
	$1.5 < y_{\gamma\gamma} < 2.5$	$p_T^{\gamma\gamma}: 0, 45, 120, 350$	
$(p_T^{\gamma 1} + p_T^{\gamma 2})/m_{\gamma\gamma}$ vs $(p_T^{\gamma 1} - p_T^{\gamma 2})/m_{\gamma\gamma}$	$0.6 < (p_T^{\gamma 1} + p_T^{\gamma 2})/m_{\gamma\gamma} \leq 0.8$	$(p_T^{\gamma 1} - p_T^{\gamma 2})/m_{\gamma\gamma}: 0, 0.3$	8
	$0.8 < (p_T^{\gamma 1} + p_T^{\gamma 2})/m_{\gamma\gamma} \leq 1.1$	$(p_T^{\gamma 1} - p_T^{\gamma 2})/m_{\gamma\gamma}: 0, 0.05, 0.1, 0.2, 0.8$	
	$1.1 < (p_T^{\gamma 1} + p_T^{\gamma 2})/m_{\gamma\gamma} \leq 4$	$(p_T^{\gamma 1} - p_T^{\gamma 2})/m_{\gamma\gamma}: 0, 0.3, 0.6, 4$	
$p_T^{\gamma\gamma}$ vs $p_T^{\gamma\gamma j}$	$N_{\text{jets}} = 0$	$p_T^{\gamma\gamma}: 0, 350$	9
	$0 < p_T^{\gamma\gamma j} \leq 30$	$p_T^{\gamma\gamma}: 0, 100, 350$	
	$30 < p_T^{\gamma\gamma j} \leq 60$	$p_T^{\gamma\gamma}: 0, 45, 120, 350$	
	$60 < p_T^{\gamma\gamma j} \leq 350$	$p_T^{\gamma\gamma}: 0, 80, 250, 450$	
$p_T^{\gamma\gamma}$ vs $\tau_{C,j1}$	$N_{\text{jets}} = 0$	$p_T^{\gamma\gamma}: 0, 350$	9
	$0 < \tau_{C,j1} \leq 15$	$p_T^{\gamma\gamma}: 0, 100, 350$	
	$15 < \tau_{C,j1} \leq 25$	$p_T^{\gamma\gamma}: 0, 120, 350$	
	$25 < \tau_{C,j1} \leq 40$	$p_T^{\gamma\gamma}: 0, 200, 350$	
	$40 < \tau_{C,j1} \leq 400$	$p_T^{\gamma\gamma}: 0, 250, 650$	
VBF-enhanced: $p_T^{j_1}$ vs $\Delta\phi_{jj}$	$-\pi < \Delta\phi_{jj} < 0$	$p_T^{j_1}: 30, 120, 500$	4
	$0 < \Delta\phi_{jj} < \pi$	$p_T^{j_1}: 30, 120, 500$	

S+B fits to $m_{\gamma\gamma}$ in different fiducial regions





χ^2 test is used to evaluate the p-value compatibility between the measurements and SM predictions

Fiducial region	Measured [fb]			SM prediction [fb]		95% CL _s upper limit [fb]	p-value
		± stat	± sys				
Diphoton	67	± 5	± 4	64	± 4	-	69%
VBF-enhanced	1.8	± 0.5	± 0.3	1.53	± 0.10	-	64%
$N_{\text{lepton}} \geq 1$	0.81	± 0.23	± 0.06	0.59	± 0.03	-	36%
High E_T^{miss}	0.28	± 0.27	± 0.07	0.302	± 0.017	0.85	93%
$t\bar{t}H$ -enhanced	0.53	± 0.27	± 0.06	0.60	± 0.05	1.13	79%
Total	132	± 10	± 8	126	± 7	-	69%

χ^2 computation includes full uncertainties in fitted cross-section and theory uncertainties from SM predictions

Response matrices for $p_T^{\gamma\gamma}$ and N_{jets} bins



p-values: measured/predicted cross-sections

Variable	p-value										
	default	RadISH NNLOJET	NNLOJet	STWZ BLPTW	MATRIX	SHERPA	GoSAM	SCET _{LIB}	TAUC	ResBos2	PROVBF
$p_T^{\gamma 1}/m_{\gamma\gamma}$	56%	—	—	—	—	—	—	58%	—	32%	—
$p_T^{\gamma 2}/m_{\gamma\gamma}$	93%	—	—	—	—	—	—	49%	—	2%	—
$p_T^{\gamma\gamma}$	86%	68%	—	—	—	—	—	78%	—	54%	—
$ y_{\gamma\gamma} $	76%	—	—	—	—	—	—	78%	—	66%	—
p_T^{j1}	78%	77%	—	—	—	47%	—	48%	—	38%	—
N_{jets}	95%	—	90%	56%	—	59%	84%	—	—	—	—
$N_{b\text{-jets}}$	60%	—	—	—	—	—	—	—	—	—	—
$p_T^{\gamma\gamma j}$	81%	—	—	—	—	68%	—	—	—	78%	—
$m_{\gamma\gamma j}$	95%	—	—	—	—	95%	—	—	—	—	—
$\tau_{C,j1}$	27%	—	—	—	—	—	—	—	11%	13%	—
$\sum \tau_{C,j}$	39%	—	—	—	—	—	—	—	—	—	—
H_T	46%	—	—	—	—	51%	—	—	—	—	—
m_{jj}	79%	—	—	—	—	81%	—	—	—	—	—
$\Delta\phi_{jj}$	91%	—	—	—	—	95%	—	—	—	—	—
$ \Delta\phi_{\gamma\gamma,jj} $	83%	—	—	—	—	88%	—	—	—	—	—
$p_{T,\gamma\gamma jj}$	99%	—	—	—	—	100%	—	—	—	—	—
$p_T^{\gamma\gamma \text{ jetveto } 30 \text{ GeV}}$	84%	—	—	—	83%	—	—	—	—	83%	—
$p_T^{\gamma\gamma \text{ jetveto } 40 \text{ GeV}}$	95%	—	—	—	45%	—	—	—	—	83%	—
$p_T^{\gamma\gamma \text{ jetveto } 50 \text{ GeV}}$	88%	—	—	—	35%	—	—	—	—	30%	—
$p_T^{\gamma\gamma \text{ jetveto } 60 \text{ GeV}}$	67%	—	—	—	52%	—	—	—	—	42%	—
$p_T^{\gamma\gamma} \text{ vs } y_{\gamma\gamma} $	75%	—	—	—	—	—	—	78%	—	—	—
$p_T^{\gamma\gamma} \text{ vs } \tau_{C,j1}$	39%	—	—	—	—	—	—	—	—	—	—
$p_T^{\gamma\gamma} \text{ vs } p_T^{\gamma\gamma j}$	96%	—	—	—	—	—	—	—	—	—	—
$(p_T^{\gamma 1} - p_T^{\gamma 2})/m_{\gamma\gamma} \text{ vs } (p_T^{\gamma 1} + p_T^{\gamma 2})/m_{\gamma\gamma}$	81%	—	—	—	—	—	—	77%	—	—	—
VBF $ \eta^* $	94%	—	—	—	—	—	—	—	—	—	70%
VBF $\Delta\phi_{jj}$	68%	—	—	—	—	—	—	—	—	—	65%
VBF p_T^{j1}	77%	—	—	—	—	—	—	—	—	—	70%
VBF $p_{T,\gamma\gamma jj}$	89%	—	—	—	—	—	—	—	—	—	74%
VBF $p_T^{j1} \text{ vs } \Delta\phi_{jj}$	76%	—	—	—	—	—	—	—	—	—	74%

Interpretation via Effective Field Theory (EFT)


- Strength and Tensor structure of the interactions of the Higgs boson: Effective Field Theory (addition of new interactions CP-even and CP-odd)
 - Constraints derived for the variables: $p_T^{\gamma\gamma}$, **N_jets**, m_{jj} , $\Delta\phi_{jj}$ and p_T^{j1}
 - BSM contributions probed as non-zero Wilson coefficients
 - Basis of parametrization SMEFT: c_i (CP-even)/ \tilde{c}_i (CP-odd) (Wilson coefficients); O_i/\tilde{O}_i (6d operators that introduce new interactions)

affect VH/VBF

$\mathcal{L}_{\text{eff}}^{\text{SMEFT}} \supset$

$c_{HG}O'_g + c_{HW}O'_{HW} + c_{HB}O'_{HB} + c_{HWB}O'_{HWB}$
 $+ c_{H\tilde{G}}\tilde{O}'_g + c_{H\tilde{W}}\tilde{O}'_{HW} + c_{H\tilde{B}}\tilde{O}'_{HB} + c_{H\tilde{W}B}\tilde{O}'_{HWB}$

affect $\Delta\phi_{jj}$



Statistical interpretation:

$$L = \exp \left[-\frac{1}{2} (\sigma_{\text{obs}} - \sigma_{\text{pred}})^T C^{-1} (\sigma_{\text{obs}} - \sigma_{\text{pred}}) \right]$$

- σ_{pred} and σ_{obs} are the k-dimensional vectors from differential cross-section (k = 34, number of bins of the 5 observables distributions)
- C is the covariance matrix: sum of the statistical, systematic and theoretical covariances

interference-terms only

