

# New Physics at CEPC

Xuai Zhuang (IHEP)

[zhuangxa@ihep.ac.cn](mailto:zhuangxa@ihep.ac.cn)



The 8th China LHC Physics Workshop  
(CLHCP2022)

Nov. 23-27, 2022




中國科學院高能物理研究所  
Institute of High Energy Physics  
Chinese Academy of Sciences

# Brief summary of BSM search @CEPC

- BSM working group formed @ 2021.4 Yangzhou WS
- Big updates presented
  - @ 2021.11 CEPC WS (13 talks)
  - @ 2022.5 CEPC WS (17 talks)
  - @ 2022.8 HEP (4-5 Talks)
  - @2022.10 CEPC WS (8 talks)
- BSM white paper is scheduled and going-on smoothly:
  - Preliminary organizers: Liantao Wang, Bruce Mellado, Xuai Zhuang, Jia Liu, Yu Gao, ...
    - ✓ **More to be invited, volunteers are very welcome!**
  - Timeline (TBD): collect inputs and a very brief white paper draft ready by end of 2022; First paper draft is ready by next Spring?
- BSM prospects at CEPC are included in CEPC snowmass white paper: [arXiv:2205.08553](https://arxiv.org/abs/2205.08553)

# CEPC BSM Physics Program



# BSM Inputs & Status



- BSM Higgs (1709.06103; 1808.02037; 1912.01431; 2008.05492 ; 2011.04540)
- SUSY Searches
  - Direct SUSY Searches (CPC46(2022)013106; 2101.12131; 2203.10580; 2202.11011)
  - Indirect search of SUSY (2010.09782)
- Dark Matter and Dark Sector searches
  - Lepton portal DM (JHEP 06 (2021) 149 )
  - Asymmetric DM (PRD 104(2021)055008 )
  - Dark Sector from exotic Z decay (1712.07237), Dark Sector-photon interactions (2208.08142)
  - DM (Millicharged DM, Vector portal DM, DM with EFT interactions): 1903.1211
  - Mono-gamma (2205.05560) ,
  - Dark Fermion in light of Electron Target Absorption ([Kai Ma's talk](#))
- Long-lived particles (1904.10661, 1911.06576, 2201.08960, Yulei Zhang's [Talk](#), Wei Su's [Talk](#), Cen Mo's [Talk](#); )
- More exotics:
  - Heavy neutrinos (2102.12826, 2201.05831);
  - Axion-like particles (2103.05218, 2204.04702, 2210.09335, [J. Phys. G](#) )
  - Electroweak phase transition (1911.10210, 1911.10206, 2011.04540, 2204.05085)
  - Identify CP-odd component in Higgs (Changlong Xu's [talk](#))
  - .....
- Global fits:
  - Global fit of SUSY (2203.04828, 2203.07883)
  - SMEFT global fit (2206.08326)

**Please let me know if any contribution is missing!**

# BSM Higgs

- A large class of BSM physics (singlet extensions, two Higgs-doublet-models (2HDM), SUSY models, Higgs portals, gauge extensions of the SM, ...) motivates these exotic decay considerations.

Representative topologies of the Higgs exotic decays




The 95% C.L. upper limit on selected Higgs exotic decay BR


- Good sensitivity of exotic Higgs decay from CEPC

# SUSY Searches at CEPC


- Mainly light EWKino and slepton for CEPC
- Lepton collider: discovery in all scenarios up to kinematic limit:  $\sqrt{s}/2$








CEPC/FCCee(240)



# SUSY Searches at CEPC







**Higgsino search**  
**CPC46(2022)013106**







# Dark Matter and Dark Sector searches





# DM search at CEPC

- Exposing Dark Sector via exotic Z-boson decay with Future Z-Factories, Jia Liu, Lian-Tao Wang, Xiao-Ping Wang, Wei Xue, [1712.07237](#), PRD 97, 095044 (2018)
- Four models include: Higgs/Vector portal DM, inelastic dark matter and axion like particles.
- Compared with HL-LHC, the reach for the BR of various exotic Z decay modes at Z-factories is sensitive for many decay modes.




[arXiv: 1712.07237](#)

# Dark Matter/Dark Sector searches


- Exposing Dark Sector-photon interactions at CEPC, Y. Zhang, M. Song and L. Chen, arXiv: 2208.08142, [Yu Zhang's talk](#)

- CEPC can probe low-mass light dark states with electromagnetic form factors via mass-dimension 5 operators.



- Exposing Dark Fermion in light of Electron Target Absorption at CEPC, Shao-Feng Ge and Kai Ma, [Kai Ma's talk](#)


- All the effective four-fermion couplings can be constrained to be well above 1TeV scale



# Long-lived particles (LLP)


## Long lifetimes result from a few simple physical mechanisms:

- Small couplings (ex. RPV SUSY)
- Limited phase space: small mass splitting (ex. compressed SUSY, ...)
- Heavy intermediate states
- .....




# LLP at Far Detector (FD)

- Physics with Far Detectors at Future Lepton Colliders, Zeren Simon Wang, Kechen Wang, [1911.06576](#), PRD 101, 075046 (2020)
- Search for long-lived axions with far detectors at future lepton colliders, Minglun Tian, Kechen Wang, Zeren Simon Wang, [2201.08960](#)
- FD can extend and complement the sensitivity to the LLPs compared with Near Detector




Light Scalars from  
Exotic Higgs Decays

[1911.06576](#)



Light Neutralinos  
from Z-boson Decays




Axion-like Particles





# Axion-like particles (ALP)

- Searching for ALP at future electron-positron colliders, H. Y. Zhang, C.X. Yue, Y.C. Guo, and S. Yang, [2103.05218](#), PRD104 (2021) 096008  
 → CEPC is more sensitive to the ALPs couplings  $g_{a\gamma\gamma}$  with mass 2-8 GeV than LHC and CLIC.
- Searching for ALP via decay  $Z \rightarrow a f \bar{f}$  at future Z factories, [2204.04702](#)
- Axion-like particle solution to muon g-2 and its test at Z-factory, Jia Liu's [talk](#)




2-8 GeV

$M_a$  (GeV)





5-70 GeV





# Axion-like particles (ALP)

- The ALP explanation to muon g-2 and its test at CEPC, J. Liu, X.L. Ma, L.T. Wang, X.P. Wang, arXiv:2210.09335, Xiao-Ping Wang's [talk](#)
- ALP can provide a g-2 solution with couplings  $C_{\mu\mu}$  and  $C_{\gamma\gamma}$ ;
- Tera-Z and Higgs factories, can completely cover the relevant parameter space through searches with final states  $(\gamma\gamma)\gamma$ ,  $(\mu^+\mu^-)\gamma$  and  $(\mu^+\mu^-)\mu^+\mu^-$ .



# Heavy neutrino search



- Monojet Search for Heavy Neutrinos at Future Z-Factories, Y.F. Shen, J.N. Ding, Q. Qin, arXiv: 2201.05831, Yin-Fa Shen' [talk](#)
- The monojet method will be able to fill the gap and has better sensitivity around the mass range between 5-15 GeV.







# EWPT at CEPC

- Electroweak Phase Transition in 2HDM under Higgs, Z-pole, and W precision measurements, H. Song, W. Su, and M. Zhang, *arXiv:2204.05085*, *JHEP* 10 (2022) 048 , H. Song's [talk](#)
- Under current constraints, both Type-I and Type-II 2HDM can explain the strong first order electroweak phase transition (SFOEWPT), Z-pole, Higgs precision measurements and mW precision measurement of CDF-II at same time.




# SUSY global fits with CEPC using GAMBIT

- Study of the impact of the Higgs and electroweak precision measurements at the CEPC with GAMBIT global fits of the SUSY models, such as CMSSM, NUHM1, NUHM2 and pMSSM-7, Yang Zhang etc, [arXiv: 2203.04828](https://arxiv.org/abs/2203.04828)
- CEPC can further test the currently allowed parameter space of these models, advance our understanding of the mass spectrum



# SMEFT global fit

- SMEFT global fit for 4-fermion and CPV operators at future colliders, 2206.08326, Yong Du's [talk](#)
- The sensitivity to new physics from global fit is significantly enhanced thanks to the high energy/ luminosity/beam polarization of future lepton colliders



# Summary and Outlook


- **CEPC has good discovery potential for NP at many scenarios which are challenge for LHC**
- **BSM prospects study at CEPC is going on well, many of the analyses are already public**
- **Plan to organize a workshop by end of this year to collect inputs for CEPC BSM white paper**
- **Please let us know if you would like to contribute to the BSM white paper !**

**Thanks for your attention!**

# Backup

20

# CEPC Physics Programme



# Identify CP-odd component in Higgs



- Use di—higgs production to identify CP-odd component in Higgs boson, Changlong Xu's [talk](#)
- Future electron-positron colliders are more powerful for exploring the ZHH Di-Higgs production

## Di-Higgs in CEPC/ILC/FCC-ee


350 GeV  $e^+e^-$  collider:  $1 \text{ ab}^{-1}$

$b\bar{b}b\bar{b}$  final state

$$\mathcal{L}_{ZhA} \sim i(h\partial_\mu A - A\partial_\mu h) Z^\mu \frac{g}{2\cos\theta_w} \cos(\beta - \alpha)$$



$$\begin{aligned} \text{Type-I: } \kappa_h^f &= \sin(\beta - \alpha) + \frac{\cos(\beta - \alpha)}{\tan \beta} \\ \kappa_A^u &= \frac{1}{\tan \beta} \quad \kappa_A^{d,\ell} = -\frac{1}{\tan \beta} \end{aligned}$$




$$\begin{aligned} \text{Type-II: } \kappa_h^u &= \sin(\beta - \alpha) + \frac{\cos(\beta - \alpha)}{\tan \beta} \\ \kappa_h^{d,\ell} &= \sin(\beta - \alpha) - \cos(\beta - \alpha) \tan \beta \\ \kappa_A^u &= \frac{1}{\tan \beta} \quad \kappa_A^{d,\ell} = \tan \beta \end{aligned}$$






# Wino & higgsino

- Prospects for chargino pair production at CEPC, Jia-Rong Yuan, Hua-Jie Cheng, Xu-Ai Zhuang, [arXiv:2105.06135](https://arxiv.org/abs/2105.06135).



Chargino pair via  
on(off)-shell W decay



Signature:  
2 lepton + MET



Discovery in  
all scenarios  
up to  
kinematic  
limit:  $\sqrt{s}/2$


# Bino NLSP at CEPC

- Probing bino NLSP at lepton colliders with Gravitino DM, Junmou Chen, Chengcheng Han, Jin Min Yang, Mengchao Zhang, [arXiv:2101.12131](https://arxiv.org/abs/2101.12131).







# Slepton search

- Prospects for slepton pair production at CEPC, Jia-Rong Yuan, Hua-Jie Cheng, Xu-Ai Zhuang, [arXiv: 2203.10580](https://arxiv.org/abs/2203.10580)




Signature:  
2 lepton + MET




# Heavy selectron search

- Probing relatively heavier right-handed selectron in the GmSUGRA, by Waqas Ahmed, Imtiaz Khan, Tianjun Li, Shabbar Raza and Wenxing Zhang, [arXiv: 2202.11011](#)
- There two types of light neutralinos that achieve the correct relic density by Z-resonance and h-resonance.

Higgs-pole  $\rightarrow m_{\tilde{\chi}_1^0} \approx \frac{1}{2}m_h$  and Z-pole  $\rightarrow m_{\tilde{\chi}_1^0} \approx \frac{1}{2}m_Z$ .




$$e^+e^- \rightarrow \tilde{\chi}_1^0(bino) + \tilde{\chi}_1^0(bino) + \gamma$$



# DM search at CEPC

- Searching for lepton portal dark matter with colliders and interplay with the gravitational wave (GW) astronomy, Jia Liu, Xiao-Ping Wang, KePan Xie, [2104.06421](#), JHEP 06 (2021) 149
- The phase transition GWs can also be a probe of the model.

$$e^+e^- \rightarrow S^{\pm(*)}S^\mp \rightarrow \ell^+\chi\ell'^-\chi \quad h/Z \rightarrow S^{\pm(*)}S^\mp \rightarrow \ell^+\chi\ell'^-\chi \text{ and } h \rightarrow \chi\chi:$$





[arXiv: 2104.06421](#)

FIG. 10. Figure from Ref. [168], the interplay between GW detection and future  $e^+e^-$  collider searches. The gray shaded region is the LISA detectable parameter space. From left to right, the sensitivities for  $\lambda_{HS}$  are shown from future CEPC precision measurements, in which the region above a given  $m_\chi$  (corresponding to a colored line) can be probed.

# DM search at CEPC

- Searching for asymmetric Dark Matter (ADM) at CEPC, Mengchao Zhang, [2104.06988](#), PRD 104, 055008 (2021)
- It is possible to generate dark quark pair through a t-channel process, dark quark  $q'$  will be a jet-like object in detector.



- The mass of mediator can be excluded up to O(10) TeV, better than LHC

# DM search at CEPC



## ■ Probing DM particles at CEPC (Millicharged DM, Vector portal DM, DM with EFT interactions): ZL, Yong-Heng Xu, Yu Zhang ,[1903.1211](#)

- ✓ CEPC can probe **millicharged DM** that is currently unexplored
- ✓ CEPC can probe the parameter space of vector-portal DM models and EFT DM models that are unconstrained by DMDD

## ■ Mono- $\gamma$ Production of a Vector Dark Matter at CEPC, K Ma, [2205.05560](#)

[ZL, Y.-H. Xu, and Y. Zhang, [1903.12114](#)]

new physics process:  $e^+e^- \rightarrow \bar{\chi}\chi\gamma$





# LL Dark Hadrons



- A theory of Dark Pions, Hsin-Chia Cheng, Lingfeng Li, Ennio Salvioni, [2110.10691](#), JHEP 01 (2022) 122, see Lingfeng's [talk](#)
- The dark quarks couple to the SM via irrelevant Z- and Higgs-portal operators. The dark pions, behave as either composite axion-like particles (ALPs) mixing with Z or h

[arXiv: 2110.10691](#)

$$\mathcal{L}_{\text{EFT}} = \frac{1}{2} \bar{\psi}_R Y^\dagger M^{-2} Y [ |H|^2 i \not{D} + i \gamma^\mu H^\dagger D_\mu H ] \psi_R + \text{h.c.} \\ + \frac{1}{2} \bar{\psi}_L \tilde{Y}^\dagger M^{-2} \tilde{Y} [ |H|^2 i \not{D} + i \gamma^\mu H^\dagger D_\mu H ] \psi_L + \text{h.c.} \\ - \bar{\psi}_L \omega \psi_R + \boxed{\bar{\psi}_L \tilde{Y}^\dagger M^{-1} Y \psi_R |H|^2} + \text{h.c.},$$

Dimension-6 Z portal couplings


Dimension-5 Higgs portal coupling



- $m_\pi < m_{\eta'}$  : dimuon mode dominates
- $m_\pi > m_{\eta'}$  : PPP modes (mostly SM  $\pi^+ \pi^- \pi^0$ )

# LLP at near Detector (ND)

- Long-lived light neutralinos at future Z-factories (RPV SUSY), Zeren Simon Wang, Kechen Wang, [1904.10661](#), PRD 101, 115018 (2020)
- The model parameter  $\lambda'_{112}/m_{\tilde{f}}^2$  can be discovered down to as low as  $\sim 1.5 \times 10^{-14}$  ( $3.9 \times 10^{-14}$ )  $\text{GeV}^{-2}$  at the FCC-ee (CEPC)



[arXiv: 1904.10661](#)




# Muon Anomalous Magnetic Moment

Present status: Discrepancy between Theory and Experiment at more than three Standard Deviation level

$$\delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{theory}} = 268(63)(43) \rightarrow 10^{-11}$$

3.6 $\sigma$  Discrepancy

New Physics at the Weak scale can fix this discrepancy. Relevant example : Supersymmetry



$$\delta a_\mu \sim \frac{e}{8\pi^2} \frac{m_\mu^2}{\tilde{m}^2} \text{Sgn}(\mu M_2) \tan \beta \sim 130 \rightarrow 10^{-11} \frac{100 \text{ GeV}}{\tilde{m}}^2 \text{Sgn}(\mu M_2) \tan \beta$$

Grifols, Mendez'85, T. Moroi'95,  
Giudice, Carena, C.W.'95, Martin and Wells'00 ....


Here  $\tilde{m}$  represents the weakly interacting supersymmetric particle masses.

For  $\tan \beta \sim 10$  (50), values of  $\tilde{m} \sim 230$  (510) GeV would be preferred.

Masses of the order of the weak scale lead to a natural explanation of the observed anomaly !

# Axion-like particles (ALP)

- The ALP explanation to muon g-2 and its test at CEPC, J. Liu, X.L. Ma, L.T. Wang, X.P. Wang, arXiv:2210.09335, Xiao-Ping Wang's [talk](#)
- ALP can provide a solution with couplings  $C_{\mu\mu}$  and  $C_{\gamma\gamma}$ ;



- In g-2 solution region, mostly decay to  $a \rightarrow \mu^+ \mu^-$
- The inclusion of  $Z$  diagram makes some difference for large  $m_a$
- Exotic  $Z$  decay should happen