



Bottomonium results at Belle II


Sen Jia (Fudan University)
on behalf of the Belle II Collaboration

e^+e^- Collisions From Φ to Ψ 2022
Aug. 15 – 19, 2022




Bottomonium

Rev. Mod. Phys. 90, 015003 (2018)



- Conventional quarkonium
- Unconventional charged states


Only two bottomonium-like states $Z_b(10610)$ and $Z_b(10650)$ [PRL 108, 122001 (2012)]:



The study of bottomonium states:

- Rich the bottomonium spectroscopy
- Test QCD nonrelativistic effective field theories and lattice QCD [Rev. Mod. Phys. 77, 1423 (2005)]
- Understand the properties of perturbative and non-perturbative QCD [EPJC 71, 1534 (2011)]
- Understand the strong interaction in the color-confinement regime [Phys. Rept. 873 (2020) 1-154]

Belle II detector






- Asymmetric energy e^+e^- collider operating near $\Upsilon(4S)$ mass peak
- Much higher luminosity than predecessor
- Upgraded detectors (better vertex and particle identification performances)
- Achieved peak luminosity: $4.7 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- Current integrated luminosity: $\sim 424/\text{fb}$ ($\sim \text{Babar} \sim 0.5 \text{ Belle}$)

Bottomonium(-like) prospects at Belle II

Four ways to access bottomonia:

- Direct production from e^+e^- : $J^{PC} = 1^{--}$: $\Upsilon(nS)$
- ISR production: $J^{PC} = 1^{--}$: $\Upsilon(nS)$
- Hadronic transitions from $\Upsilon(nS)$ through η , $\pi\pi$, etc
 $J^{PC} = 0^{-+}, 1^{--}, 1^{+-} \dots : \Upsilon(nS), \eta_b(nS), h_b(nS), \dots$
- Radiative transitions from $\Upsilon(nS)$
 $J^{PC} = 0^{-+}, 0^{++}, 1^{++}, 2^{++} : \eta_b(nS), \chi_b$




Bottomonium(-like) prospects at Belle II

Run at $Y(6S)$ and $Y(5S)$ and high energy scan:

- Search for new missing bottomonia $\eta_b(3S)$, $h_b(3P)$, $Y(D)$, exotic states Y_b , Z_b , etc
- Improve precision of already known processes and states, e.g., Z_b
- Measure the effect of the coupled channel contribution
- Study $B^{(*)}\bar{B}^{(**)}$ and $B_s^{(*)}B_s^{(**)}$ threshold regions (challenging for Super-KEKB)

Run at $Y(3S)$ and $Y(2S)$:

- Search for missing $\pi\pi/\eta$ Transitions in inclusive decays to constrain further models
- Search for new physics: LFV, LFU, light Higgs, ...



New result

**Observation of $e^+e^- \rightarrow \omega\chi_{bJ}$ and
search for $X_b \rightarrow \omega Y(1S)$ at \sqrt{s} near 10.75 GeV**

Motivation: $\Upsilon(10753)$

- Belle: several $\sim 1\text{fb}^{-1}$ scan points below $\Upsilon(5S)$
- New structure observed in $\pi^+\pi^-\Upsilon(nS)$ transitions

	$\Upsilon(10860)$	$\Upsilon(11020)$	New structure
M (MeV/c ²)	$10885.3 \pm 1.5^{+2.2}_{-0.9}$	$11000.0^{+4.0}_{-4.5} {}^{+1.0}_{-1.3}$	$10752.7 \pm 5.9^{+0.7}_{-1.1}$
Γ (MeV)	$36.6^{+4.5}_{-3.9} {}^{+0.5}_{-1.1}$	$23.8^{+8.0}_{-6.8} {}^{+0.7}_{-1.8}$	$35.5^{+17.6}_{-11.3} {}^{+3.9}_{-3.3}$

- Theoretical interpretations


Conventional D- or S-D mixed bottomonium:

PRD 105, 074007 (2022), PRD 104, 034036 (2021)
 EPJC 80, 59 (2020), PRD 101, 397 014020 (2020)
 PRD 102, 014036399 (2020), EPJP 137, 357 (2022)
 PRD 105, 114041 (2022), PLB 803, 135340 (2020)
 arXiv:2204.11915, Prog. Part. Nucl. Phys. 117, 103845 (2021)

A tetraquark:


PLB 802, 135217 (2020)
 PRD 103, 074507 (2021)
 arXiv:2205.11475
 Chin. Phys. C 43, 123102 (2019)

- Interpretations as an admixture of the conventional 4S and 3D state predict comparable branching fractions of 10^{-3} for $\Upsilon(10753) \rightarrow \pi^+\pi^-\Upsilon(nS)$ and $\Upsilon(10753) \rightarrow \omega\chi_{bJ}$ [PRD 104, 034036 (2021), PRD 105, 074007 (2022)].



X_b : Bottomonium counterpart of $X(3872)$?


- Two close peaks observed in the cross sections for $e^+e^- \rightarrow \pi^+\pi^-J/\psi$ by BESIII^[1] and $e^+e^- \rightarrow \pi^+\pi^-\Upsilon(nS)$ by Belle^[2], respectively, may suggest similar nature.
- $\Upsilon(4220) \rightarrow \gamma X(3872)$ ^[3] and $\omega\chi_{c0}$ ^[4] observed by BESIII.
- So expect the $\Upsilon(10753)$ state to decay into γX_b with $X_b \rightarrow \omega\Upsilon(1S)$, as well as a potential resonance in the line shape of $\sigma(e^+e^- \rightarrow \omega\chi_{bJ})$.



[1] PRL 118, 092001 (2017); [2] JHEP 10, 220 (2019); [3] PRL 122, 232002 (2019); [4] PRD 99, 091103 (2019)

Unique scan data near $\sqrt{s} = 10.75$ GeV

- In November 2021, Belle II collected 19 fb⁻¹ of unique data at energies above the $\Upsilon(4S)$: four energy scan points around 10.75 GeV.
- Physics goal: understand the nature of the $\Upsilon(10753)$ energy region.



Analysis goals



$e^+e^- \rightarrow \omega\chi_{bJ}$:

- Determine the Born cross section for $e^+e^- \rightarrow \omega\chi_{bJ}$ using unique scan data samples at $\sqrt{s} = 10.701, 10.745$ and 10.805 GeV.
- Study the energy dependence of Born cross section for $e^+e^- \rightarrow \omega\chi_{bJ}$ by combining with Belle data at $\sqrt{s} = 10.867$ GeV [PRL 113, 142001 (2014)].

$e^+e^- \rightarrow \gamma X_b$:

- Search for the X_b using unique scan data samples at $\sqrt{s} = 10.653, 10.701, 10.745$ and 10.805 GeV.

Analysis overview





- Event selection criteria

- 4 or 5 charged particles
- standard Belle II PID: 90-95% efficiency with 1-5% misID
- χ_{bJ} photon energy > 50 MeV
- $0.105 < M(\gamma\gamma) < 0.150$ GeV/ c^2 for π^0 (90% efficiency)
- constrained kinematic fit to $\pi^+\pi^-\pi^0\gamma e^+e^-/\mu^+\mu^-$ final states
- best candidate based on fit quality


- Data driven corrections and systematics from control samples

Mass distributions



- Red box contains 95% of signals

Observation of $e^+e^- \rightarrow \omega\chi_{bJ}$





Two dimensional unbinned maximum likelihood fits to the $M(\gamma Y(1S))$ and $M(\pi^+\pi^-\pi^0)$ distributions.

Channel	\sqrt{s} (GeV)	N_{sig}	$\sigma_{\text{Born}}^{(\text{UL})}$ (pb)
$\omega\chi_{b1}$	10.745	$68.9^{+13.7}_{-13.5}$	$3.6^{+0.7}_{-0.7} \pm 0.4$
$\omega\chi_{b2}$		$27.6^{+11.6}_{-10.0}$	$2.8^{+1.2}_{-1.0} \pm 0.5$
$\omega\chi_{b1}$	10.805	$15.0^{+6.8}_{-6.2}$	1.6 @90% C.L.
$\omega\chi_{b2}$		$3.3^{+5.3}_{-3.8}$	1.5 @90% C.L.

The total χ_{bJ} signal significances are 11.1σ and 4.5σ at $\sqrt{s} = 10.745$ and 10.805 GeV.

Note that the $\sigma_{\text{Born}}(e^+e^- \rightarrow \omega\chi_{b1}/\omega\chi_{b2})$ is only $(0.76 \pm 0.16)/(0.29 \pm 0.14)$ pb at $\sqrt{s} = 10.867$ GeV [PRL 113, 142001(2014)].

Observation of $\Upsilon(10753) \rightarrow \omega\chi_{bJ}$



The $e^+e^- \rightarrow \omega\chi_{bJ}$ ($J = 1, 2$) cross sections peak at $\Upsilon(10753)$.

Combine Belle II measurements with Belle measurement^[1] to fit cross section with function:

$$\sigma_{e^+e^- \rightarrow \omega\chi_{b1}}(\sqrt{s}) = |\sqrt{PS_2(\sqrt{s}) + BW(\sqrt{s})e^{i\phi}}|^2, BW(\sqrt{s}) = \frac{\sqrt{12\pi\Gamma_{ee}\mathcal{B}_f\Gamma}}{s - M^2 + iM\Gamma} \sqrt{\frac{PS_2(\sqrt{s})}{PS_2(M)}}$$

M and Γ of $\Upsilon(10753)$ are fixed according to Ref. [2].


$\Gamma_{ee}\mathcal{B}_f$	Solution I (constructive interference)	Solution II (destructive interference)
$\Gamma_{ee}\mathcal{B}(\Upsilon(10753) \rightarrow \omega\chi_{b1})$	$(0.63 \pm 0.39 \pm 0.20)$ eV	$(2.01 \pm 0.38 \pm 0.76)$ eV
$\Gamma_{ee}\mathcal{B}(\Upsilon(10753) \rightarrow \omega\chi_{b2})$	$(0.53 \pm 0.46 \pm 0.15)$ eV	$(1.32 \pm 0.44 \pm 0.55)$ eV

- $\frac{\Gamma_{ee}\mathcal{B}(\Upsilon(10753) \rightarrow \omega\chi_{b1})}{\Gamma_{ee}\mathcal{B}(\Upsilon(10753) \rightarrow \omega\chi_{b2})} \sim 1$ agrees with the expectation for NRQCD
- $\frac{\Gamma_{ee}\mathcal{B}(\omega\chi_{b1/b2})}{\Gamma_{ee}\mathcal{B}(\pi^+\pi^-\Upsilon(2S))^{[2]}} \sim 1.5$ for $\Upsilon(10753)$ and 0.1 for $\Upsilon(10860)$

[1] PRL 113, 142001(2014); [2] JHEP 10, 220 (2019)

Search for X_b

$$e^+e^- \rightarrow \gamma X_b (\rightarrow \omega Y(1S))$$



- No significant X_b signal is observed.
- The peaks are the reflections of $e^+e^- \rightarrow \omega \chi_{bJ}$.

From simulated events with $m(X_b) = 10.6 \text{ GeV}/c^2$
The yield is fixed at the upper limit at 90% C.L.

Upper limits at 90% C.L. on $\sigma_B(e^+e^- \rightarrow \gamma X_b) \cdot$ $\mathcal{B}(X_b \rightarrow \omega Y(1S))$ (pb)	\sqrt{s} (GeV)	10.653	10.701	10.745	10.805
	$m(X_b) = 10.6 \text{ GeV}/c^2$	0.46	0.33	0.10	0.14
	$m(X_b) = (10.45, 10.65) \text{ GeV}/c^2$	(0.14, 0.55)	(0.25, 0.84)	(0.06, 0.14)	(0.08, 0.37)

Belle II potential – 10.75 GeV

Other active ongoing analyses based on unique scan data at Belle II:

Channel
$B\bar{B}$ decomposition
$e^+e^- \rightarrow \gamma X_b$
$e^+e^- \rightarrow \omega \eta_b(1S)$
$e^+e^- \rightarrow Y(1S) + X$
$e^+e^- \rightarrow \pi^+\pi^- h_b(nP)$
$e^+e^- \rightarrow \pi^+\pi^- Y_2(1D)$
$e^+e^- \rightarrow \pi^+\pi^- Y(nS)$
$e^+e^- \rightarrow \eta h_b(1P)$
$e^+e^- \rightarrow \phi \eta_b(1S)$

- Quarkonium spectroscopy (conventional and exotic)
- Hadronic and radiative transitions
- Annihilations in exclusive final states
- Precision study of the vector states using ISR
- New physics in Bottomonium decays
- ...

Summary

Belle II bottomonium program includes

- Early run near 10.75 GeV, Nov. 2021 (Done)
- $Y(3S)$, $Y(5S)$, $Y(6S)$, and scan above $Y(5S)$ in the future [PTEP 2019 (2019) 12, 123C01]


Based on early run data near $\sqrt{s} = 10.75$ GeV

- New decay modes of $Y(10753) \rightarrow \omega\chi_{bJ}$ are observed for the first time.
- The value of $\Gamma_{ee}\mathcal{B}(Y(10753) \rightarrow \omega\chi_{b1}/\omega\chi_{b2})$ is in the range 0.20–2.9/0.05–2.0 eV, which is consistent with $\Gamma_{ee}\mathcal{B}(\pi^+\pi^-Y(nS))$ ($n = 0, 1$, or 2) [JHEP 10, 220(2019)]. But this is quite different from the $Y(10860)$, where the value of $\Gamma_{ee}\mathcal{B}(Y(10860) \rightarrow \omega\chi_{b1}/\omega\chi_{b2})$ is much smaller from PDG.
- No evidence of a X_b signal is obtained with X_b masses between 10.45 and 10.65 GeV/c², and the upper limits at 90% C.L. are set.

Thanks for your attention!

Backup slides

Alternative fit for $\sigma(e^+e^- \rightarrow \omega\chi_{bJ})$



$$\sigma_{\omega\chi_{c1/2}}(\sqrt{s}) = |BW_{Y(10753)} + BW_{Y(10806)}e^{i\phi}|^2$$

M and Γ are fixed referring to Ref. [1][2]

$$\frac{\Gamma_{ee}B(Y(10753) \rightarrow \omega\chi_{b1})}{\Gamma_{ee}B(Y(10753) \rightarrow \omega\chi_{b2})}$$

Solution I: $\frac{1.24 \pm 0.56(\text{stat.})}{0.92 \pm 0.37(\text{stat.})}$

Solution II: $\frac{1.28 \pm 0.57(\text{stat.})}{1.09 \pm 0.40(\text{stat.})}$

$\chi^2/\text{ndf} = 0.4$ and 0.1 for $\omega\chi_{b1}$ and $\omega\chi_{b2}$

Ref: [1]. JHEP 10, 220(2019);

[2]. PDG 2022