

Dark SHINE – Dark Photon fixed-target search experiment at SHINE Facility

粒子物理天问论坛 2022

Jing Chen

on behalf of the Dark SHINE team

2022.08.06



上海交通大学
SHANGHAI JIAO TONG UNIVERSITY




Evidence of dark matter




Gravitational Lensing




The Bullet Cluster



Rotation Curve



Cosmic Microwave Background




Dark matter candidates



Mass scale of dark matter

(not to scale)

arXiv:1904.07915



- **Searching for light DM:**

- Sub-GeV mass range not fully explored yet.
- Dark photon A' :
 1. Dark matter particles may interact with other dark matter particles via a new force mediated by A' .
 2. Collider experiments searching for dark photon: NA64@CERN, BESIII, BEPCII, LDMX, etc.

- Dark matter is the mysterious substance that makes up roughly a quarter of the Universe.
- Dark Matter can exist in **wide mass range**, from Ultralight DM to Primordial Black Holes.
- experimental search:
 - Space experiments: **DAMPE, AMS etc.**
 - Collider experiments: **LHC, BELLE-II , BESIII etc.**
 - Underground experiments: **CDEX, PandaX, LUX, Xenon etc.**


Search for dark photon



- Dark photon is an important portal between the standard model (SM) particles and the dark matter.

not couple directly to SM particles

obtain a small coupling to the EM current due to ϵ



$$L = L_{SM} + \epsilon F^{\mu\nu} F'_{\mu\nu} + \frac{1}{4} F'^{\mu\nu} F'_{\mu\nu} + m_{A'}^2 A'^{\mu} A'_{\mu}$$

Kinetic mixing term

Field strength tensor

Dark photon field

- The minimal dark-photon model with 3 unknown parameters:

ϵ : kinetic mixing between the SM hypercharge and A' field strength tensors.

$m_{A'}$: dark photon mass.


[arXiv:2104.10280](https://arxiv.org/abs/2104.10280)

Decay branching ratio of $A' \rightarrow \chi\chi$ (assumed to be 1 or 0)

Search for dark photon




- A' production:



- Bremsstrahlung, $eZ \rightarrow eZA'$ & $pZ \rightarrow pZA'$, fixed-target experiment
- Annihilation, $e^+e^- \rightarrow A'\gamma$, e^+e^- collider
- Drell-Yan, $q\bar{q} \rightarrow A'$, hadron collider / fixed nuclear target w/ proton-beam
- Meson decay, $\pi^0 \rightarrow A'\gamma$ or $\eta \rightarrow A'\gamma$ (w/ $m_{A'} < m_{\pi,\eta}$), any experiment w/ high meson production rates

- A' decay:



- Visible decay

Two interaction vertices \rightarrow production rate highly suppressed

- Invisible decay


One interaction vertex \rightarrow interaction probability enhanced

Better sensitivity!

The SHINE facility



- Dark SHINE:
 - **Fixed-target** experiment w/ high frequency **single electron beam** provided by Shanghai High Repetition-Rate XFEL and Extreme Light Facility(**SHINE**)
 - Invisible decay: $m_{A'} > 2m_\chi$, **missing energy / missing momentum**
 - Search for A' in $[m_{A'}, \varepsilon]$ parameter space
- The SHINE:
 - Under construction in Zhangjiang area, Shanghai (2018-2026).
 - $E_{beam} = 8GeV$ with frequency 1MHz
 - Beam intensity: 6.25×10^8 electrons per bunch




Science Bulletin 61,
117(2016), 720-727



The SHINE facility



Single electron beam is needed for Dark SHINE.



Dark SHINE Kicker:
1MHz → 10MHz

3×10^{14} electron-on-target(EOTs) per year!

FEL kicker


1300 buckets provided
by 1.3GHz microwave

100pC in one bucket
 6.25×10^8 electrons per bunch

Dark SHINE Kicker

electron beam w/ one
electron per bunch

Dark SHINE detector



- **Tracker:**
 - Tagging tracker + recoil tracker
 - Incident and recoil electron tracks
 - Two silicon strip sensors w/ a small angle (0.1rad)
 - Resolution: $6\mu m$ (horizontal), $60\mu m$ (vertical)


- **ECAL:**
 - Electron & photon
 - Scintillator: LYSO(Ce)
 - high light yield, fast decay time, low electronic noise
 - $20 \times 20 \times 11$ crystals
 - $2.5 \times 2.5 \times 4 cm^3$
 - Energy resolution of LYSO: 10%

- **HCAL:**
 - Veto hadronic background
 - Scintillator w/ steel absorber
 - $4 \times 4 \times 1$ modules
- **Magnet:**
 - 1.5T magnetic field


Signal & background

- Signal signature:

Most of the incident momentums is transferred to A' .



- Major background processes:



Leading background:
 photon bremsstrahlung

Rare processes:
 photon-nuclear, $\gamma \rightarrow \mu\mu$,
 electron-nuclear



上海交通大学
SHANGHAI JIAO TONG UNIVERSITY

李政道研究所
Tsung-Dao Lee Institute



Jing Chen
Page . 10

Event display



Simulation





Signal

- Low momentum of recoil electron
- Recoil electron angle has on average value





Background

- Recoil electron carry most of the incident momentum
- Recoil electron angle small





ECAL energy vs. HCAL energy




Signal region



- Signal region definition:

number of the reconstructed tracks, $N_{\text{trk}}^{\text{tag,rec}} = 1$;
 missing momentum of electron, $p_{\text{tag}} - p_{\text{rec}} > 4 \text{ GeV}$;
 total energy reconstructed in ECAL, $E_{\text{ECAL}}^{\text{total}} < 2.5 \text{ GeV}$;
 total energy reconstructed in HCAL, $E_{\text{HCAL}}^{\text{total}} < 0.1 \text{ GeV}$;
 max. cell energy in HCAL, $E_{\text{HCAL}}^{\text{MaxCell}} < 2 \text{ MeV}$.

- Signal efficiency:



- 25 mass points (1×10^5 events in each mass point) are produced.

- Around 60% signal events survive the cut-flow.

- Efficiency drops in:

Low-mass region of a few MeV: tight energy cuts.

High-mass region above 1 GeV: particles with large incident/recoil angle go into the HCAL directly.

Background estimation



- Background samples:

Process	Generate Events	Branching Ratio	EOTs
Inclusive	2.5×10^9	1.0	2.5×10^9
Bremsstrahlung	1×10^7	6.70×10^{-2}	1.5×10^8
GMM_target	1×10^7	$1.5(\pm 0.5) \times 10^{-8}$	4.3×10^{14}
GMM_ECAL	1×10^7	$1.63(\pm 0.06) \times 10^{-6}$	6.0×10^{12}
PN_target	1×10^7	$1.37(\pm 0.05) \times 10^{-6}$	4.0×10^{12}
PN_ECAL	1×10^8	$2.31(\pm 0.01) \times 10^{-4}$	4.4×10^{11}
EN_target	1×10^8	$5.1(\pm 0.3) \times 10^{-7}$	1.6×10^{12}
EN_ECAL	1×10^7	$3.25(\pm 0.08) \times 10^{-6}$	1.8×10^{12}

Inclusive background:
 2.5×10^9 EOTs

$\Rightarrow > 3 \times 10^{14}$ EOTs

Lack of statistics!


Extrapolate from fit results

- Cut-flow: No background events left after SR selection.

Background estimation



- Fit the fraction of events below energy cutoff as a function of cut values on ECAL energy.
- **Extrapolate from inclusive background simulation.**
- Validation from inclusive background simulation.
- Extrapolate from rare processes simulation.



$y = 10^{-14}$: less than one background event left w/ ECAL energy cut.

Extrapolate from the fit results.

Lack of statistics in low “cut on $E_{\text{ECal}}^{\text{total}}$ ” region.


Event yield (3×10^{14} EOTs):

$$2.53 \times 10^{-3}$$


Background estimation

- Validation from inclusive background simulation.
- Statistics is limited in $E_{beam} = 8\text{ GeV}$ inclusive samples.
- In extrapolation of inclusive background simulation, the fit range is far away from the final $E_{E\text{Cal}}^{\text{total}}$ cut (2.5 GeV).
- Inclusive samples with E_{beam} from 3 – 7.5 GeV are used to estimate events in low $E_{E\text{Cal}}^{\text{total}}$.
- Scale low E_{beam} events to match the shoulder with $E_{beam} = 8\text{ GeV}$ events.
- **Event yield from direct extrapolation(3×10^{14} EOTs):**

$$N_{100,2} = 3 \times 10^{14} \times N_{100,20} \cdot \frac{N_{fit,100,2}}{N_{fit,100,20}} \\ = 9.23 \times 10^{-3}$$



Background estimation




rare processes scaled according to the corresponding branching ratio.

- **Extrapolate from rare processes simulation.**
- Rare processes background samples are produced with larger statistics.
- Fit the fraction of events below energy cutoff in other rare processes (EN(ECAL), EN(target), PN(ECAL), PN(target)).

Background estimation



- **Extrapolate from rare processes simulation.**
- Estimate the number of background events corresponds to 3×10^{14} EOTs.
- Don't need to further extrapolation on:



GMM_target

4.3×10^{14} EOTs > 3×10^{14} EOTs

GMM_ECAL:





6.0×10^{12} EOTs

Energy carried by the muon pair

HCAL requirements can highly suppress these events (fraction of the remaining GMM events $< 10^{-6}$)

Background estimation

- Extrapolation from rare processes simulation



3×10^{14} EOTs

Bkg. Events:
0.015





Most conservative
number

Sensitivity study



- Expected 90% C.L. limit estimated with 3×10^{14} EOTs (running ~1 year), 9×10^{14} EOTs (~3 years), 1.5×10^{15} EOTs (~5 years) and 1×10^{16} EOTs (with Phase-II upgrade).



- Comparing with other experiments, Dark SHINE can provide competitive sensitivity.



Summary



- Dark SHINE: a fixed-target experiment to search for light dark matter.
- Detector R&D ongoing.
- First round of preliminary study has been finished:
 - Good signal efficiency, background well suppressed (~ 0.015 bkg. event expected for 1 year operation).
 - Expecting competitive sensitivity.
 - Submitted to Science China.

The project is officially sponsored by NSFC Original Exploration Project 2021 and Shanghai pilot program for basic research.



Backup



Dark SHINE detector






Table 1 The detector geometry overview.

Node	Centre (mm)	Size (mm)			Arrangement	Comments
	z	x	y	z		
Tagging Tracker	-307.783	200	400	600.216	7 layers	Second layer rotation: 0.1 rad
Target	0	100	200	0.35		
Recoil Tracker	94.032	500	800	172.714	6 layers	Second layer rotation: 0.1 rad
ECAL	408.539	506	506	454.3	20 × 20 × 11 cells	
HCAL	2660.69	4029.51	4029.51	4048.01	4 × 4 × 1 modules	

Tracker design


- Silicon tracker geometry



Tag tracker						
Z position/mm	-607.5	-507.5	-407.5	-307.5	-207.5	-107.5
X size/mm	100	100	100	100	100	100
Y size/mm	200	200	200	200	200	200

Recoil tracker						
Z position/mm	7.5	22.5	38.5	53.5	89.5	180.5
X size/mm	100	100	100	120	180	250
Y size/mm	200	200	200	230	280	400



Z thickness $\sim 100\mu\text{m}$, 0.001λ
Resolution x: $\sim 6\mu\text{m}$, y: $\sim 60\mu\text{m}$








Calorimeter Design (ECAL)

- **Cubic design** of crystal for the electronic calorimeter
 - Z segmentation for 3D shower reconstruction and (potential) PID
 - Potential PFA combined with tracker: location resolution and better track regression
- Readout with SiPM(light sensor) and waveform sampling
 - Wide dynamic range and abundant models for different application
 - Compact size and (relatively) easy to drive
 - High repeated rate and strictly zero integral/dead time with fast ADC&DAQ
- **LYSO crystal** chosen as baseline design with XY=2.5cm Z=4cm
 - High light yield with good linearity
 - Radiation hard and short decay time





HCAL design




Parameter for the whole HCAL

X:100cm


Y:100cm

Z:360cm

xy crossing



Inclusive cross-section



Inclusive cross-section of dark photon bremsstrahlung from electron interacting with W target, assuming $\varepsilon = 1$.



Invisible background




- Neutrino productions:
 - Moller scattering $e^-e^- \rightarrow e^-e^-$ followed by charged-current quasi-elastic (CCQE) reaction $e^-p \rightarrow \nu_e n$.
 - Neutrino pair production $e^-N \rightarrow e^-N\nu\bar{\nu}$.
 - Bremsstrahlung \oplus CCQE and charge-current exchange with exclusive $e^-p \rightarrow \nu n\pi_0$. No recoil electron, track requirement can remove these processes.


Table 6 Expected invisible background production corresponds to 3×10^{14} EOTs, estimated from different irreducible reaction scenarios. The Bremsstrahlung \oplus CCQE and the charge-current exchange productions can be effectively rejected by the one-track requirement.

irreducible reaction	Moller scattering	neutrino pair production
estimated yield	3×10^{-4}	$< 1.8 \times 10^{-5}$
irreducible reaction	Bremsstrahlung \oplus CCQE	charge-current exchange
estimated yield	0.3	0.3

Dark photon search experimental results



arXiv: 1707.04591



MASS OF THE DARK PHOTON

NA64, PRL123, 121801 (2019)