

# 粒子物理和早期宇宙对称性自发破缺产生的随机引力波

**Ligong Bian(边立功)**


**Chongqing University (重庆大学)**

[lgbycl@cqu.edu.cn](mailto:lgbycl@cqu.edu.cn)


**2022/11/05@北京大学高能物理研究中心**

# Contents


- **Motivation**
- **First-order phase transition & EWSB**
- **Cosmic string & Domain wall**
- **Future**








LHC 探测实验



希格斯质量与电弱相变


随机引力波探测开启了探索早期宇宙背后新物理的一个新的窗口

# The Gravitational Wave Spectrum



强一阶  
电弱相变

电弱重子数产生  
WIMP/FIMP暗物质



Electroweak baryogenesis and dark matter from a singlet Higgs #2

James M. Cline (McGill U.), Kimmo Kainulainen (Jyvaskyla U. and Helsinki Inst. of Phys. and Helsinki U.) (Oct, 2012)

Published in: *JCAP* 01 (2013) 012 • e-Print: 1210.4196 [hep-ph]

pdf DOI cite 217 citations

标准模型+实标量场

Gravitational wave, collider and dark matter signals from a scalar singlet electroweak baryogenesis #4

Ankit Beniwal (Adelaide U. and Adelaide U., Sch. Chem. Phys.), Marek Lewicki (Adelaide U. and Warsaw U. and Adelaide U., Sch. Chem. Phys.), James D. Wells (DESY and Michigan U., MCTP), Martin White (Adelaide U., Sch. Chem. Phys. and Adelaide U.), Anthony G. Williams (Adelaide U., Sch. Chem. Phys. and Adelaide U.) (Feb 20, 2017)

Published in: *JHEP* 08 (2017) 108 • e-Print: 1702.06124 [hep-ph]

pdf DOI cite 131 citations

Impact of a complex singlet: Electroweak baryogenesis and dark matter #6

Minyuan Jiang (Beijing, Inst. Theor. Phys. and Beijing, KITPC and Nanjing U.), Ligong Bian (Beijing, Inst. Theor. Phys. and Beijing, KITPC), Weicong Huang (Beijing, Inst. Theor. Phys. and Beijing, KITPC), Jing Shu (Beijing, Inst. Theor. Phys. and Beijing, KITPC) (Feb 26, 2015)

Published in: *Phys.Rev.D* 93 (2016) 6, 065032 • e-Print: 1502.07574 [hep-ph]

pdf DOI cite 标准模型+复标量场 102 citations

Unified explanation for dark matter and electroweak baryogenesis with direct detection and gravitational wave signatures #5


Mikael Chala (DESY), Germano Nardini (U. Bern, AEC), Ivan Sobolev (Lomonosov Moscow State U. and Moscow, INR) (May 27, 2016)

Published in: *Phys.Rev.D* 94 (2016) 5, 055006 • e-Print: 1605.08663 [hep-ph]


pdf DOI cite 复合希格斯 125 citations




## 超出粒子物理标准模型 新物理




## 早期宇宙演化



解释宇宙正  
反物质不对  
称



预言相变引力  
波，检验爱因  
斯坦广义相对  
论



电弱对称性破  
缺，给出基本  
粒子质量起源

# ► Effective potential at zero temperature

Action 
$$S[\phi] = \int d^4x \mathcal{L}\{\phi(x)\}$$

The generating functional (vacuum-to-vacuum amplitude):


$$Z[j] = \langle 0_{\text{out}} | 0_{\text{in}} \rangle_j \equiv \int d\phi \exp\{i(S[\phi] + \phi j)\} \quad \phi j \equiv \int d^4x \phi(x) j(x)$$

The connected generating functional  $W[j]$  defined as:

$$Z[j] \equiv \exp\{iW[j]\}$$

The effective action  $\Gamma[\phi]$  as the Legendre transformation:

$$\Gamma[\bar{\phi}] = W[j] - \int d^4x \frac{\delta W[j]}{\delta j(x)} j(x) \quad \bar{\phi}(x) = \frac{\delta W[j]}{\delta j(x)}$$

 
$$\frac{\delta \Gamma[\bar{\phi}]}{\delta \bar{\phi}} = \frac{\delta W[j]}{\delta j} \frac{\delta j}{\delta \bar{\phi}} - j - \bar{\phi} \frac{\delta j}{\delta \bar{\phi}} = -j$$

$$\left. \frac{\delta \Gamma[\bar{\phi}]}{\delta \bar{\phi}} \right|_{j=0} = 0$$

# ► Effective potential at zero temperature

Expand  $Z[j]$  ( $W[j]$ ) in a power series of  $j$ , to obtain its representation in terms of Green functions  $G_{(n)}$  (connected Green functions  $G_{(n)}^c$ )

$$Z[j] = \sum_{n=0}^{\infty} \frac{i^n}{n!} \int d^4x_1 \dots d^4x_n j(x_1) \dots j(x_n) G_{(n)}(x_1, \dots, x_n)$$

$$iW[j] = \sum_{n=0}^{\infty} \frac{i^n}{n!} \int d^4x_1 \dots d^4x_n j(x_1) \dots j(x_n) G_{(n)}^c(x_1, \dots, x_n)$$

The effective action can be expanded as

$$\Gamma[\bar{\phi}] = \sum_{n=0}^{\infty} \frac{1}{n!} \int d^4x_1 \dots d^4x_n \bar{\phi}(x_1) \dots \bar{\phi}(x_n) \Gamma^{(n)}(x_1, \dots, x_n)$$

$\Gamma^{(n)}$  are the one-particle irreducible (1PI) Green functions

Fourier transformation

$$\Gamma^{(n)}(x) = \int \prod_{i=1}^n \left[ \frac{d^4p_i}{(2\pi)^4} \exp\{ip_i x_i\} \right] (2\pi)^4 \delta^{(4)}(p_1 + \dots + p_n) \Gamma^{(n)}(p)$$

$$\tilde{\phi}(p) = \int d^4x e^{-ipx} \bar{\phi}(x)$$

$$\Gamma[\bar{\phi}] = \sum_{n=0}^{\infty} \int \prod_{i=1}^n \left[ \frac{d^4p_i}{(2\pi)^4} \tilde{\phi}(-p_i) \right] (2\pi)^4 \delta^{(4)}(p_1 + \dots + p_n) \Gamma^{(n)}(p_1, \dots, p_n) \quad (1)$$

# ► Effective potential at zero temperature

Translationally invariant theory, with  $\bar{\phi}_c$  being constant  $\bar{\phi}(x) = \phi_c$

Define the effective potential  $V_{\text{eff}}(\phi_c)$  as

$$\Gamma[\phi_c] = - \int d^4x V_{\text{eff}}(\phi_c) \quad (2)$$

Using the definition of Dirac  $\delta$ -function

$$\delta^{(4)}(p) = \int \frac{d^4x}{(2\pi)^4} e^{-ipx}$$

We get

$$\tilde{\phi}_c(p) = (2\pi)^4 \phi_c \delta^{(4)}(p).$$

Inserting into EQ.(1), the effective action for constant field configurations recast the form of

$$\Gamma(\phi_c) = \sum_{n=0}^{\infty} \frac{1}{n!} \phi_c^n (2\pi)^4 \delta^{(4)}(0) \Gamma^{(n)}(p_i = 0) = \sum_{n=0}^{\infty} \frac{1}{n!} \phi_c^n \Gamma^{(n)}(p_i = 0) \int d^4x \quad (3)$$

(2,3)



$$V_{\text{eff}}(\phi_c) = - \sum_{n=0}^{\infty} \frac{1}{n!} \phi_c^n \Gamma^{(n)}(p_i = 0) \quad (4)$$

Expanding in powers of momentum, about the point where all external momenta vanish


$$\Gamma[\bar{\phi}] = \int d^4x \left[ -V_{\text{eff}}(\bar{\phi}) + \frac{1}{2} (\partial_\mu \bar{\phi}(x))^2 Z(\bar{\phi}) + \dots \right]$$

# Effective potential at zero temperature

An example

Tree-level potential

$$V_0 = \frac{1}{2}m^2\phi^2 + \frac{\lambda}{4!}\phi^4$$



In momentum space the scalar field is

$$\phi_c(p) = (2\pi)^4 \phi_c \delta^4(p)$$

Recall EQ.(4), we get one-loop potential:

$$\begin{aligned} V_1(\phi_c) &= i \sum_{n=1}^{\infty} \int \frac{d^4 p}{(2\pi)^4} \frac{1}{2n} \left[ \frac{\lambda \phi_c^2 / 2}{p^2 - m^2 + i\epsilon} \right]^n \\ &= -\frac{i}{2} \int \frac{d^4 p}{(2\pi)^4} \log \left[ 1 - \frac{\lambda \phi_c^2 / 2}{p^2 - m^2 + i\epsilon} \right] \end{aligned}$$

After Wick rotation:

$$V_1(\phi_c) = \frac{1}{2} \int \frac{d^4 p_E}{(2\pi)^4} \log \left[ 1 + \frac{\lambda \phi_c^2 / 2}{p_E^2 + m^2} \right]$$

$$p^0 = ip_E^0, \quad p_E = (-ip^0, \vec{p}), \quad p^2 = (p^0)^2 - \vec{p}^2 = -p_E^2$$

1-loop effective potential is

$$V_1(\phi_c) = \frac{1}{2} \int \frac{d^4 p}{(2\pi)^4} \log [p^2 + m^2(\phi_c)] \quad (5)$$

$$V_{\text{eff}}(\phi_c) = V_0(\phi_c) + V_1(\phi_c)$$

shifted mass :

$$m^2(\phi_c) = m^2 + \frac{1}{2}\lambda\phi_c^2 = \frac{d^2 V_0(\phi_c)}{d\phi_c^2}$$



With dimensional regularization

$$V_1(\phi_c) = \frac{1}{2}(\mu^2)^{2-\frac{n}{2}} \int \frac{d^n p}{(2\pi)^n} \log[p^2 + m^2].$$

We calculate the one-loop correction to the effective potential by first calculating it with respect to the mass and then integrating.

$$\frac{\partial V_1}{\partial m^2} = \frac{1}{2}(\mu^2)^{2-\frac{n}{2}} \int \frac{d^n p}{(2\pi)^n} \frac{1}{p^2 + m^2}$$

The derivative is just a single disconnected bubble.

$$V_1 = \frac{m^4}{64\pi} \left( -\left[ \frac{1}{\epsilon} - \gamma_E + \log 4\pi \right] + \log \frac{m^2}{\mu^2} - \frac{3}{2} \right)$$

Subtracting the  $1/\epsilon - \gamma - \log 4\pi$  term, we get

$$V_1 = \frac{1}{64\pi^2} m^4 \left( \log \frac{m^2}{\mu^2} - \frac{3}{2} \right)$$

$$m^2 = d^2V/d\phi^2$$

# ► Effective potential at finite temperature-imaginary time

Feynman rules for the different fields in the imaginary time formalism:

$$\text{Boson propagator} : \frac{i}{p^2 - m^2}; p^\mu = [2ni\pi\beta^{-1}, \vec{p}]$$

$$\text{Fermion propagator} : \frac{i}{\gamma \cdot p - m}; p^\mu = [(2n + 1)i\pi\beta^{-1}, \vec{p}]$$

$$\text{Loop integral} : \frac{i}{\beta} \sum_{n=-\infty}^{\infty} \int \frac{d^3p}{(2\pi)^3}$$

$$\text{Vertex function} : -i\beta(2\pi)^3 \delta_{\sum \omega_i} \delta^{(3)}\left(\sum_i \vec{p}_i\right)$$

With above FR EQ.(5) becomes

$$V_1^\beta(\phi_c) = \frac{1}{2\beta} \sum_{n=-\infty}^{\infty} \int \frac{d^3p}{(2\pi)^3} \log(\omega_n^2 + \omega^2) \quad (6)$$

$$\text{with } \omega^2 = \vec{p}^2 + m^2(\phi_c)$$

Define

$$v(\omega) = \sum_{n=-\infty}^{\infty} \log(\omega_n^2 + \omega^2)$$

We have

$$v(\omega) = 2\beta \left[ \frac{\omega}{2} + \frac{1}{\beta} \log(1 - e^{-\beta\omega}) \right] + \omega - \text{independent terms}$$

Substituting into EQ.(6) we get

$$V_1^\beta(\phi_c) = \int \frac{d^3p}{(2\pi)^3} \left[ \frac{\omega}{2} + \frac{1}{\beta} \log(1 - e^{-\beta\omega}) \right] \quad (7)$$

$$\frac{1}{2} \int \frac{d^3p}{(2\pi)^3} \omega = \frac{1}{2} \int \frac{d^4p}{(2\pi)^4} \log[p^2 + m^2(\phi_c)] \quad \frac{1}{\beta} \int \frac{d^3p}{(2\pi)^3} \log(1 - e^{-\beta\omega}) = \frac{1}{2\pi^2\beta^4} J_B[m^2(\phi_c)\beta^2]$$

# ► Effective potential at finite temperature-real time

Propagators for scalar fields can be written as

$$G(p) \equiv \begin{pmatrix} G^{(11)}(p) & G^{(12)}(p) \\ G^{(21)}(p) & G^{(22)}(p) \end{pmatrix}$$

$$G^{(11)}(p) = \Delta(p) + 2\pi n_B(\omega_p) \delta(p^2 - m^2)$$

$$G^{(22)}(p) = G^{(11)*}$$

$$G^{(12)} = 2\pi e^{\beta\omega_p/2} n_B(\omega_p) \delta(p^2 - m^2)$$

$$G^{(21)} = G^{(12)}$$

$$n_B(\omega) = \frac{1}{e^{\beta\omega} - 1}$$

$\Delta(p)$  is the boson/fermion propagator at zero temperature

The propagators for fermion fields can be written as

$$S(p)_{\alpha\beta} \equiv \begin{pmatrix} S_{\alpha\beta}^{(11)}(p) & S_{\alpha\beta}^{(12)}(p) \\ S_{\alpha\beta}^{(21)}(p) & S_{\alpha\beta}^{(22)}(p) \end{pmatrix}$$

$$S^{(11)}(p) = (\gamma \cdot p + m) (\Delta(p) - 2\pi n_F(\omega_p) \delta(p^2 - m^2))$$

$$S^{(22)}(p) = S^{(11)*}$$

$$S^{(12)} = -2\pi(\gamma \cdot p + m) [\theta(p^0) - \theta(-p^0)] e^{\beta\omega_p/2} n_F(\omega_p) \delta(p^2 - m^2)$$

$$S^{(21)} = -S^{(12)}$$

$$n_F(\omega) = \frac{1}{e^{\beta\omega} + 1}$$

The main feature of the real time formalism is that the propagators come in two terms:

1. one which is the same as in the zero temperature field theory ( $\Delta(p)$ ), and a second one where all the temperature dependence is contained.
2. (12), (21) and (22) components are unphysical since one of their time arguments has an imaginary component.

# ► Effective potential at finite temperature-real time

Disconnected bubble diagrams

$$\frac{dV_1^\beta}{dm^2(\phi_c)} = \frac{1}{2} \int \frac{d^4p}{(2\pi)^4} \left[ \frac{-i}{-p^2 + m^2(\phi_c) - i\epsilon} + 2\pi n_B(\omega) \delta(p^2 - m^2(\phi_c)) \right] \quad (8)$$

After integration on  $m^2(\phi_c)$ , the first part contributes to the effective potential as

$$-\frac{i}{2} \int \frac{d^4p}{(2\pi)^4} \log(-p^2 + m^2(\phi_c) - i\epsilon)$$

Considering

$$-\frac{i}{2} \int_{-\infty}^{\infty} \frac{dx}{2\pi} \log(-x^2 + \omega^2 - i\epsilon) = \frac{\omega}{2} + \text{constant}$$

Performing the  $p^0$  integral, we get

$$\int \frac{d^3p}{(2\pi)^3} \frac{\omega}{2} \quad (9)$$

Using the identity

$$\delta(p^2 - m^2) = \frac{1}{2\omega_p} [\delta(p^0 + \omega_p) + \delta(p^0 - \omega_p)]$$

Integration over  $p^0$  in the  $\beta$ -dependent of the EQ 8, we get



$$\int \frac{d^3p}{(2\pi)^3} \frac{1}{2\omega} n_B(\omega) \quad (10)$$

Upon integration over  $m^2(\phi_c)$  leads to the second term of EQ (6)

# Thermal effective scalar potential for PT study

$$V_T(\phi, T) = V_0(\phi) + T^4 \left[ \sum_B J_B \left( \frac{M_B}{T} \right) + \sum_F J_F \left( \frac{M_F}{T} \right) \right]$$

all fermions **F** and bosons **B** that are relativistic at temperature **T**



**High-T expansion**


$$m/T \ll 1$$

$$V_T(\phi) = V_0(\phi) + \frac{T^2}{24} \left( \sum_S M_S^2(\phi) + 3 \sum_V M_V^2(\phi) + 2 \sum_F M_F^2(\phi) \right) - \frac{T}{12\pi} \left( \sum_S \left( M_S^2(\phi) \right)^{\frac{3}{2}} + \sum_V \left( M_V^2(\phi) \right)^{\frac{3}{2}} \right) + \text{higher order terms.}$$


**MS, MV, MF are the masses of the scalar fields S, vector fields V and fermionic fields F**

# Phase transition types


## Second order





## First order




# FOEWPT & Higgs physics



Higgs announcement seminar on 4 July 2012




Grojean, Servant, Wells 05, Huang, Jokelar, Li, Wagner 15, Cao, Huang, Xie, & Zhang 17, Zhou, Bian, Guo 19





# BNPC, $v/T$ and EW sphaleron



$$\partial_\mu J_B^\mu = i \frac{N_F}{32\pi^2} \left( -g_2^2 F^{a\mu\nu} \tilde{F}_{\mu\nu}^a + g_1^2 f^{\mu\nu} \tilde{f}_{\mu\nu} \right),$$

$$\Delta B = N_F (\Delta N_{CS} - \Delta n_{CS}),$$


$$N_{CS} = -\frac{g_2^2}{16\pi^2} \int d^3x \, 2\epsilon^{ijk} \text{Tr} \left[ \partial_i A_j A_k + i \frac{2}{3} g_2 A_i A_j A_k \right],$$

$$n_{CS} = -\frac{g_1^2}{16\pi^2} \int d^3x \, \epsilon^{ijk} \partial_i B_j B_k,$$


$$A_i \rightarrow U A_i U^{-1} + \frac{i}{g_2} (\partial_i U) U^{-1},$$

$$\delta N_{CS} = \frac{1}{24\pi^2} \int d^3x \, \text{Tr} \left[ (\partial_i U) U^{-1} (\partial_j U) U^{-1} (\partial_k U) U^{-1} \right] \epsilon^{ijk}.$$

The Standard Model already contains a process that violates B-number. It is known as the electroweak sphaleron ("sphaleron" is Greek for "ready to fall").



Klinkhammer & Manton (1984); Kuzmin, Rubakov, & Shaposhnikov (1985); Harvey & Turner (1990) but also identified earlier by Dashen, Hasslacher, & Neveu (1974) and Boguta (1983)



Lattice result,  $T_C = (159.5 \pm 1.5) \text{GeV}$ , Phys.Rev.Lett,113, 141602 (2014).

$$\Gamma^{\text{sym}} \approx 6 \times (18 \pm 3) \alpha_W^5 T^4, \quad \Gamma^{\text{brok}} \sim T^4 \exp\left(-\frac{E_{\text{sph}}}{T}\right)$$

## Washout avoidance, BNPC

$$\Gamma_{\text{sph}} = A_{\text{sph}}(T) \exp[-E_{\text{sph}}(T)/T] < H(T)$$

$$PT_{\text{sph}} \equiv \frac{E_{\text{sph}}(T)}{T} - 7 \ln \frac{v(T)}{T} + \ln \frac{T}{100 \text{ GeV}} \quad PT_{\text{sph}} > (35.9 - 42.8)$$





$$E_{\text{sph}}(T) \approx E_{\text{sph},0} \frac{v(T)}{v} \quad \frac{v(T)}{T} > (0.973 - 1.16) \left( \frac{E_{\text{sph},0}}{1.916 \times 4\pi v/g} \right)^{-1}$$

Hiren H. Patel and Michael J. Ramsey-Musolf, 15'

Xucheng Gan, Andrew J. Long, Lian-Tao Wang, 17'



# Model classes for one-step FOPT



Chung, Long, Wang, Phys.Rev.D 87 (2013) 2, 023509

# ► Thermal driven Class-I

$$V_{\text{eff}}(h, T) \approx \frac{1}{2}(-\mu^2 + cT^2)h^2 - \frac{eT}{12\pi}(h^2)^{3/2} + \frac{\lambda}{4}h^4$$

$$e \sim \sum_{\text{light bosonic fields}} (\text{degrees of freedom}) \times (\text{coupling to Higgs})^{3/2}.$$

$$\frac{v(T_c)}{T_c} \approx \frac{e}{6\pi\lambda}$$

TABLE I. Examples of models in the Thermally (BEC) Driven class. The expressions for  $e$  are calculated in the limit that the field-independent contributions to  $m_{\text{eff}}^2(h, T)$  are negligible (e.g., the thermal mass tuning has been performed). Here, the symbol  $\tilde{A}_t$  is  $\tilde{A}_t = A_t - \mu/\tan\beta$  and  $g_s$  is the number of real scalar singlet degrees of freedom coupling to the Higgs.

Model	$-\Delta\mathcal{L}$	$c$	$e$
SM [43]		$c_{\text{SM}} = \frac{6m_t^2 + 6m_W^2 + 3m_Z^2 + \frac{3}{2}m_H^2}{12v^2}$	$e_{\text{SM}} = \frac{6m_W^3 + 3m_Z^3}{v^3}$
MSSM [41]		$c_{\text{SM}} + \frac{6m_t^2}{12v^2} \left(1 - \frac{\tilde{A}_t^2}{m_Q^2}\right)$	$e_{\text{SM}} + \frac{6m_t^3}{v^3} \left(1 - \frac{\tilde{A}_t^2}{m_Q^2}\right)^{3/2}$
Colored scalar [20]	$M_X^2 X ^2 + \frac{K}{6} X ^4 + Q H ^2 X ^2$	$c_{\text{SM}} + \frac{6}{24} \frac{Q}{2}$	$e_{\text{SM}} + 6\left(\frac{Q}{2}\right)^{3/2}$
Singlet scalar [43,44]	$M^2 S ^2 + \lambda_S S ^4 + 2\zeta^2 H ^2 S ^2$	$c_{\text{SM}} + \frac{g_S}{24} \zeta^2$	$e_{\text{SM}} + g_S \zeta^3$
Singlet Majoron [45]	$\mu_s^2 S ^2 + \lambda_s S ^4 + \lambda_{hs} H ^2 S ^2 + \frac{1}{2}y_i S\nu_i\nu_i + \text{H.c.}$	$c_{\text{SM}} + \frac{2}{24} \frac{\lambda_{hs}}{2}$	$e_{\text{SM}} + 2\left(\frac{\lambda_{hs}}{2}\right)^{3/2}$
Two-Higgs doublets [46]	$\mu_D^2 D^\dagger D + \lambda_D(D^\dagger D)^2 + \lambda_3 H^\dagger H D^\dagger D + \lambda_4  H^\dagger D ^2 + (\lambda_5/2)[(H^\dagger D)^2 + \text{H.c.}]$	$c_{\text{SM}} + \frac{2\lambda_3 + \lambda_4}{12}$	$e_{\text{SM}} + 2\left(\frac{\lambda_3}{2}\right)^{3/2} + \left(\frac{\lambda_3 + \lambda_4 - \lambda_5}{2}\right)^{3/2} + \left(\frac{\lambda_3 + \lambda_4 + \lambda_5}{2}\right)^{3/2}$

# ► Tree driven-Class IIA

$$V_{\text{eff}}(\varphi, T) \approx \frac{1}{2}(m^2 + cT^2)\varphi^2 - \mathcal{E}\varphi^3 + \frac{\lambda}{4}\varphi^4$$

$$T_c \approx \sqrt{\frac{m^2}{c}} \sqrt{\frac{2\mathcal{E}^2}{\lambda m^2} - 1}, \quad \frac{v(T_c)}{T_c} \approx \sqrt{\frac{2c}{\lambda}} \frac{1}{\sqrt{1 - \frac{\lambda m^2}{2\mathcal{E}^2}}} \cos\alpha.$$

TABLE II. Examples of models that fall into Class IIA. For the non-SUSY models, corrections to the SM Lagrangian are shown, whereas for the SUSY models only the superpotential corrections are given.

Model	$\Delta\mathcal{L}$
xSM [53–56]	$\frac{1}{2}(\partial S)^2 - [\frac{b_2}{2}S^2 + \frac{b_3}{3}S^3 + \frac{b_4}{4}S^4 + \frac{a_1}{2}H^\dagger HS^2 + \frac{a_2}{2}H^\dagger HS^2]$
$\mathbb{Z}_2$ xSM [14,57]	$\frac{1}{2}(\partial S)^2 - [\frac{b_2}{2}S^2 + \frac{b_4}{4}S^4 + \frac{a_2}{2}H^\dagger HS^2]$
Two-Higgs doublets [58]	$\mu_D^2 D ^2 + \lambda_D D ^4 + \lambda_3 H ^2 D ^2 + \lambda_4 H^\dagger D ^2 + (\lambda_5/2)[(H^\dagger D)^2 + \text{H.c.}]$
Model	$\Delta W$
NMSSM [59–61]	$\lambda H_1 H_2 N - \frac{\kappa}{3}N^3 + rN$
nMSSM [62]	$\lambda H_1 H_2 S + \frac{m_{12}^2}{\lambda} S$
$\mu\nu$ MSSM [63]	$-\lambda_i H_1 H_2 \nu_i^c + \frac{\kappa_{ijk}}{3} \nu_i^c \nu_j^c \nu_k^c + Y_\nu^{ij} H_2 L_i \nu_j^c$



# Class IIA (1) no extra EWSB: xSM

For the “xSM” model, the gauge invariant finite temperature effective potential is found to be:

$$V(h, s, T) = -\frac{1}{2}[\mu^2 - \Pi_h(T)]h^2 - \frac{1}{2}[-b_2 - \Pi_s(T)]s^2 + \frac{1}{4}\lambda h^4 + \frac{1}{4}a_1 h^2 s + \frac{1}{4}a_2 h^2 s^2 + \frac{b_3}{3}s^3 + \frac{b_4}{4}s^4, \quad (C1)$$

with the thermal masses given by

$$\Pi_h(T) = \left( \frac{2m_W^2 + m_Z^2 + 2m_t^2}{4v^2} + \frac{\lambda}{2} + \frac{a_2}{24} \right) T^2, \quad (C2)$$

$$\Pi_s(T) = \left( \frac{a_2}{6} + \frac{b_4}{4} \right) T^2,$$

## PT strength

$$v^{\text{xSM}}/T \equiv \frac{v_h(T)}{T} = \frac{\sqrt{v_h^2(T) + v_s^2(T)} \cos \theta(T)}{T},$$

$$\cos \theta(T) \equiv \frac{v_h(T)}{\sqrt{v_h^2(T) + v_s^2(T)}},$$


□

For small mixing limit between the extra Higgs and the SM Higgs, one have

$$c_4^{\text{xSM}} = -\frac{a_1^2 - 8b_2\lambda}{32b_2} + \frac{\theta^2(a_1^2(6b_2 - \mu^2) - 8a_1b_2b_3 + 8b_2^2(a_2 - 2\lambda))}{32b_2^2} + O(\theta^3)$$

$$c_6^{\text{xSM}} = -\frac{a_1^2(a_1b_3 - 3a_2b_2)}{192b_2^3} - \frac{\theta^2 a_1}{256b_2^4} (a_1^3b_2 + 4a_1^2b_3(\mu^2 - 3b_2) + 4a_1b_2(a_2(11b_2 - 2\mu^2) - 6b_2(b_4 + \lambda) + 4b_3^2) - 32a_2b_2^2b_3) + O(\theta^3)$$

$$c_8^{\text{xSM}} = \frac{a_1^4b_4}{1024b_2^4} + \frac{a_1^3\theta^2}{1024b_2^5} (a_1(a_2b_2 + 4b_4(\mu^2 - 3b_2)) + 16b_2b_3b_4) + O(\theta^3)$$



# Class IIA (1) with extra EWSB: **GM model**

The most general scalar potential  $V(\Phi, \Delta)$  invariant under  $SU(2)_L \times SU(2)_R \times U(1)_Y$  is given by

$$\begin{aligned}
 V(\Phi, \Delta) = & \frac{1}{2}m_1^2 \text{tr}[\Phi^\dagger \Phi] + \frac{1}{2}m_2^2 \text{tr}[\Delta^\dagger \Delta] + \lambda_1 \left( \text{tr}[\Phi^\dagger \Phi] \right)^2 + \lambda_2 \left( \text{tr}[\Delta^\dagger \Delta] \right)^2 + \lambda_3 \text{tr} \left[ \left( \Delta^\dagger \Delta \right)^2 \right] + \lambda_4 \text{tr}[\Phi^\dagger \Phi] \text{tr}[\Delta^\dagger \Delta] \\
 & + \lambda_5 \text{tr} \left[ \Phi^\dagger \frac{\sigma^a}{2} \Phi \frac{\sigma^b}{2} \right] \text{tr}[\Delta^\dagger T^a \Delta T^b] \\
 & + \mu_1 \text{tr} \left[ \Phi^\dagger \frac{\sigma^a}{2} \Phi \frac{\sigma^b}{2} \right] (P^\dagger \Delta P)_{ab} + \mu_2 \text{tr}[\Delta^\dagger T^a \Delta T^b] (P^\dagger \Delta P)_{ab} ,
 \end{aligned} \tag{3}$$

extra EWSB

**Custodial symmetry**

$v_\chi = \sqrt{2}v_\xi$

$$\Phi \equiv (\varepsilon_2 \phi^*, \phi) = \begin{pmatrix} \phi^{0*} & \phi^+ \\ -\phi^{+*} & \phi^0 \end{pmatrix}, \quad \Delta \equiv (\varepsilon_3 \chi^*, \xi, \chi) = \begin{pmatrix} \chi^{0*} & \xi^+ & \chi^{++} \\ -\chi^{+*} & \xi^0 & \chi^+ \\ \chi^{++*} & -\xi^{+*} & \chi^0 \end{pmatrix}, \tag{1}$$

with

$$\varepsilon_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \varepsilon_3 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & -1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \tag{2}$$

where the phase convention for the scalar field components is:  $\chi^{--} = \chi^{++*}$ ,  $\chi^- = \chi^{+*}$ ,  $\xi^- = \xi^{+*}$ ,  $\phi^- = \phi^{+*}$ .  $\Phi$  and  $\Delta$  are transformed under  $SU(2)_L \times SU(2)_R$  as  $\Phi \rightarrow U_{2,L} \Phi U_{2,R}^\dagger$  and  $\Delta \rightarrow U_{3,L} \Delta U_{3,R}^\dagger$  with  $U_{L,R} = \exp(i\theta_{L,R}^a T^a)$  and  $T^a$  being the  $SU(2)$  generators.

where summations over  $a, b = 1, 2, 3$  are understood,  $\sigma$ 's and  $T$ 's are the  $2 \times 2$  (Pauli matrices)  $3 \times 3$  matrix representations of the  $SU(2)$  generators, respectively



$$T_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, T_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -i & 0 \\ i & 0 & -i \\ 0 & i & 0 \end{pmatrix}, T_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix},$$

The  $P$  matrix, which is the similarity transformation relating the generators in the triplet and adjoint representations, is given by

$$P = \frac{1}{\sqrt{2}} \begin{pmatrix} -1 & i & 0 \\ 0 & 0 & \sqrt{2} \\ 1 & i & 0 \end{pmatrix}.$$

xSM: without extra EWSB

GM: with extra EWSB




$$g_{hxx} = \cos \alpha g_{hxx}^{SM}$$

$$g_{hf\bar{f}} = \cos \alpha / \cos \theta_H g_{hf\bar{f}}^{SM}, \quad g_{hVV} = (\cos \alpha \cos \theta_H - \sqrt{\frac{8}{3}} \sin \alpha \sin \theta_H) g_{hf\bar{f}}^{SM},$$

$$g_{Hf\bar{f}} = \sin \alpha / \cos \theta_H g_{hf\bar{f}}^{SM}, \quad g_{HVV} = (\sin \alpha \cos \theta_H + \sqrt{\frac{8}{3}} \cos \alpha \sin \theta_H) g_{hVV}^{SM}.$$

# Collider & GW complementary search

SNR > 10 points for **two-step** and **one-step** SFOEWPT



Circles and the dotted points for the GM and xSM scenarios

$$\delta\kappa_3^{\text{xSM}} = \alpha_H^2 \left[ -\frac{3}{2} + \frac{2m_H^2 - 2b_3 v_s - 4b_4 v_s^2}{m_h^2} \right] + \mathcal{O}(\alpha_H^3),$$

$$\delta\kappa_4^{\text{xSM}} = \alpha_H^2 \left[ -3 + \frac{5m_H^2 - 4b_3 v_s - 8b_4 v_s^2}{m_h^2} \right] + \mathcal{O}(\alpha_H^3).$$

$$\delta\kappa_3^{\text{GM}} = -\alpha_H \frac{\sqrt{3}\mu_1 v}{2m_h^2} + \frac{\alpha_H v^2 (4\alpha_H - \sqrt{6}\theta_H)(2\lambda_4 + \lambda_5)}{2m_h^2}$$

$$- \frac{(3\alpha_H^2 + \theta_H^2)}{2} + \mathcal{O}(\alpha_H^3, \theta_H^3),$$

$$\delta\kappa_4^{\text{GM}} = -2\alpha_H^2 \left( 1 - \frac{2(2\lambda_4 + \lambda_5)v^2}{m_h^2} \right) + \mathcal{O}(\alpha_H^3).$$

# ► Tree-level driven-Class II B

< 0 causes the potential to turn over

$$V_{\text{eff}}(h, T) \approx \frac{1}{2}(\mu^2 + cT^2)h^2 + \frac{\lambda}{4}h^4 + \frac{1}{8\Lambda^2}h^6$$

stabilizes the EW-broken vacuum

$$\begin{aligned} \lambda &= \frac{m_H^2}{2v^2} \left(1 - \frac{\Lambda_{\text{max}}^2}{\Lambda^2}\right), & \Lambda_{\text{max}} &\equiv \sqrt{3}v^2/m_H, & T_c &= \sqrt{\frac{\mu^2}{c}} \sqrt{\frac{\lambda^2 \Lambda^2}{4\mu^2} - 1}, \\ \mu^2 &= \frac{m_H^2}{2} \left(\frac{\Lambda_{\text{max}}^2}{2\Lambda^2} - 1\right), & \Lambda &< \Lambda_{\text{max}}, & \frac{v(T_c)}{T_c} &= \sqrt{-\lambda} \frac{2}{\sqrt{1 - \frac{4\mu^2}{\lambda^2 \Lambda^2}}}, \\ \lambda_{HHH} &\equiv \frac{m_H^2}{v} \left(1 + 2\frac{\Lambda_{\text{min}}^2}{\Lambda^2}\right), & \Lambda_{\text{min}} &= v^2/m_H, \end{aligned}$$

Model	Couplings	Wilson coefficient of $H^6$
$\mathbb{R}$ Singlet	$-\frac{1}{2}\lambda_{HS} H ^2 S^2 - g_{HS}H^\dagger HS$	$-\frac{\lambda_{HS}}{2} \frac{g_{HS}^2}{M^4}$
$\mathbb{C}$ Singlet	$-g_{HS} H ^2\Phi - \frac{\lambda_{H\Phi}}{2} H ^2\Phi^2 - \frac{\lambda'_{H\Phi}}{2}H^\dagger H \Phi ^2 + h.c.$	$-\frac{ g_{HS} ^2 \lambda'_{H\Phi}}{2M^4} - \frac{\text{Re}[g_{HS}^2 \lambda_{H\Phi}]}{M^4}$
2HDM	$-Z_6 H_1 ^2 H_1^\dagger H_2 - Z_6^* H_1 ^2 H_2^\dagger H_1$	$\frac{ Z_6 ^2}{M^2}$
$\mathbb{R}$ triplet	$gH^\dagger \tau^a H \Phi^a - \frac{\lambda_{H\Phi}}{2} H ^2 \Phi^a ^2$	$-\frac{g^2}{M^4} \left(\frac{\lambda_{H\Phi}}{8} - \lambda\right)$
$\mathbb{C}$ triplet	$gH^T i\sigma_2 \tau^a H \Phi^a - \frac{\lambda_{H\Phi}}{2} H ^2 \Phi^a ^2$ $-\frac{\lambda'}{4}H^\dagger \tau^a \tau^b H \Phi^a (\Phi^b)^\dagger + h.c.$	$-\frac{g^2}{M^4} \left(\frac{\lambda_{H\Phi}}{4} + \frac{\lambda'}{8} - 2\lambda\right)$
$\mathbb{C}$ 4-plet	$-\lambda_{H3\Phi} H_i^* H_j^* H_k^* \Phi^{ijk} + h.c.$	$\frac{ \lambda_{H3\Phi} ^2}{M^2}$

1705.02551



**Higgs potential**  $V(H) = -m^2(H^\dagger H) + \lambda (H^\dagger H)^2 + \frac{(H^\dagger H)^3}{\Lambda^2}$

**Finite temperature potential**  $V_T(h, T) = V(h) + \frac{1}{2}c_{hT}h^2$

**Thermal correction**  $c_{hT} = (4y_t^2 + 3g^2 + g'^2 + 8\lambda)T^2/16$

**Electroweak minimum  
being the global one**

$$\Lambda \geq v^2/m_h$$

**Potential barrier requirement**

$$\Lambda < \sqrt{3}v^2/m_h$$

# ► Loop driven-Class III

$$V_{\text{eff}}(h, T) \approx \frac{1}{2}(\mu^2 + cT^2)h^2 + \frac{\lambda}{4}h^4 + \frac{\kappa}{4}h^4 \ln \frac{h^2}{M^2}$$

$$\lambda = \frac{m_H^2}{2v^2} - \kappa \left( \ln \frac{v^2}{M^2} + \frac{3}{2} \right), \quad T_c \approx \frac{m_H}{2\sqrt{c}} \sqrt{\epsilon} \left( 1 + \frac{1}{8}\epsilon + \frac{37}{384}\epsilon^2 + \dots \right), \quad \epsilon = 1 - \kappa v^2 / m_H^2$$

$$\mu^2 = -\frac{m_H^2}{2} + \kappa v^2, \quad \frac{v(T_c)}{T_c} \approx \frac{2v\sqrt{c}}{m_H} \frac{1}{\sqrt{\epsilon}} \left( 1 - \frac{3}{8}\epsilon - \frac{103}{384}\epsilon^2 + \dots \right).$$

TABLE III. Examples of models in the Loop Driven class.

Model	$-\Delta \mathcal{L}$
Singlet scalars [12,72]	$\sum_i^N M^2  S_i ^2 + \lambda_S  S_i ^4 + 2\zeta^2  H ^2  S_i ^2$
Singlet Majoron [73,74]	$\mu_s^2  S ^2 + \lambda_s  S ^4 + \lambda_{hs}  H ^2  S ^2 + \frac{1}{2} y_i S \nu_i \nu_i + \text{H.c.}$
Two-Higgs doublets [75–78]	$\mu_D^2 D^\dagger D + \lambda_D (D^\dagger D)^2 + \lambda_3 H^\dagger H D^\dagger D + \lambda_4  H^\dagger D ^2 + (\lambda_5/2)[(H^\dagger D)^2 + \text{H.c.}]$

$$V(h_1, h_2, T) = V_0(h_1, h_2) + V_{\text{CW}}(h_1, h_2) + V_{\text{CT}}(h_1, h_2) + V_{\text{th}}(h_1, h_2, T) + V_{\text{daisy}}(h_1, h_2, T)$$

Tree-level

$$V_0(h_1, h_2) = \frac{1}{2} m_{12}^2 t_\beta (h_1 - h_2 t_\beta^{-1})^2 - \frac{v^2}{4} \frac{\lambda_1 h_1^2 + \lambda_2 h_2^2 t_\beta^2}{1 + t_\beta^2} - \frac{v^2}{4} \frac{\lambda_{345} (h_1^2 t_\beta^2 + h_2^2)}{1 + t_\beta^2} + \frac{1}{8} \lambda_1 h_1^4 + \frac{1}{8} \lambda_2 h_2^4 + \frac{1}{4} \lambda_{345} h_1^2 h_2^2$$

One-loop at zero temperature:

$$V_{\text{CW}}(h_1, h_2) = \sum_i (-1)^{2s_i} n_i \frac{\hat{m}_i^4(h_1, h_2)}{64\pi^2} \left[ \ln \left( \frac{\hat{m}_i^2(h_1, h_2)}{Q^2} \right) - C_i \right] \quad [\text{Coleman, Weinberg '73}]$$

One-loop at finite temperature:

$$V_{\text{th}}(h_1, h_2, T) = \frac{T^4}{2\pi^2} \sum_i n_i J_{B,F} \left( \frac{m_i^2(h_1, h_2)}{T^2} \right) \quad [\text{Dolan, Jackiw '74}]$$

$$J_{B,F}(y) = \mp \sum_{l=1}^{\infty} \frac{(\pm 1)^l y}{l^2} K_2(\sqrt{y}l) \quad [\text{Anderson, Halle '92}]$$

$$V_{\text{daisy}}(h_1, h_2, T) = -\frac{T}{12\pi} \sum_i n_i \left[ (M_i^2(h_1, h_2, T))^{\frac{3}{2}} - (m_i^2(h_1, h_2))^{\frac{3}{2}} \right]$$

[Carrington '92; Arnold, Espinosa '93; Delaunay, Grojean, Wells '07]

# Beyond SM models for FOPT

## Higgs&GWs

### SM+Scalar Singlet


Bian, Huang, Shu 15, Cheng, Bian 17, Bian, Tang 18, Chen, Li, Wu, Bian, 19

### SM+Scalar Doublet

Bernon, Bian, Jiang 17, Bian, Liu 18

### SM + Scalar Triplet

Zhou, Cheng, Deng, Bian, Wu 18, Zhou, Bian, Guo, Wu 19, Zhou, Bian, Du, 22



### Composite Higgs

Bian, Wu, Xie 19, Bian, Wu, Xie 20

### NMSSM

Bi, Bian, Huang, Shu, Yin 15, Bian, Guo, Shu 17

### SMEFT



Zhou, Bian, Guo 19

Symmetry breakdown


Symmetry breaking process

Double Higgs Production at Colliders Workshop

$$\Delta\mathcal{L} = -\frac{1}{2} \frac{m_h^2}{v} (1 + \delta\kappa_3) h^3 - \frac{1}{8} \frac{m_h^2}{v^2} (1 + \delta\kappa_4) h^4$$





SNR > 10 for two-step and one-step SFOEWPT



PTA, LISA, TianQin, Taiji, LIGO, ...



# Sphaleron energy and SFOEWPT condition





$$PT_{sph} \equiv \frac{E_{sph}(T)}{T} - 7 \ln \frac{v(T)}{T} + \ln \frac{T}{100 \text{ GeV}}$$

SMEFT


xSM

$$PT_{sph} > (35.9 - 42.8)$$




# Sphaleron Energy & GW

## xSM & SMEFT







Zhou, [Bian\\*](#), Guo\*, Phys.Rev.D 101 (2020) 091903(R)







Scen.	$m_H$ [GeV]	$m_A$ [GeV]	$m_{H^\pm}$ [GeV]	Type	Main $H/A$ decays
A	130 – 300	400 – 600	100 – 300	I	$A \rightarrow W^- H^+$ (60%), $ZH$ (25%)
B	400 – 600	10 – 200	400 – 600	I, II	$H \rightarrow ZA$ (50 – 75%)
C	130 – 200	450 – 800	450 – 800	I, II	$A \rightarrow ZH$ ( $\sim 100\%$ )
D	400 – 600	10 – 250	100 – 250	I	$H \rightarrow W^- H^+$ (60%), $ZA$ (25%)
E	300 – 350	300 – 350	300 – 350	I	$A \rightarrow Zh$ ( $\sim 100\%$ ), $H \rightarrow W^+ W^-$ ( $\gtrsim 40\%$ )



# Class III 2HDM 13 TeV cross sections at the LHC







# Class III 2HDM Triple Higgs coupling











type I

type II



Dashed (solid) line depicts the  $v_w = 0.1(1)$  scenarios.



超出粒子物理模型  
新物理模型

Particle  
physics model


PT parameters

Effective action  $\rightarrow \beta, H_*$

Energy budget  $\rightarrow \alpha, \kappa(\alpha, v_w)$

Bubble wall dynamics  $\rightarrow v_w$

有限温场论计算




GW power spectrum

Numerical simulations  $\rightarrow$   
 $h^2 \Omega_{\text{GW}}(f; H_*, \alpha, \beta, v_w)$

格点场论模拟

重要的引力波源，主要科学目标之一

PTA, LIGO, LISA, 天琴, 太极, ...



LISA sensitivity

Configuration + noise level  $\rightarrow$   
 $h^2 \Omega_{\text{sens}}(f)$


Signal-to-noise ratio

**SNR**



# ► Key Events in the early Universe

Event	time $t$	redshift $z$	temperature $T$
Inflation	$10^{-34}$ s	–	–
Baryogenesis	?	?	?
EW phase transition	20 ps	$10^{15}$	100 GeV
QCD phase transition	$20 \mu\text{s}$	$10^{12}$	150 MeV
Dark matter freeze-out	?	?	?
Neutrino decoupling	1 s	$6 \times 10^9$	1 MeV
Electron-positron annihilation	6 s	$2 \times 10^9$	500 keV
Big Bang nucleosynthesis	3 min	$4 \times 10^8$	100 keV
Matter-radiation equality	60 kyr	3400	0.75 eV
Recombination	260–380 kyr	1100–1400	0.26–0.33 eV
Photon decoupling	380 kyr	1000–1200	0.23–0.28 eV
Reionization	100–400 Myr	11–30	2.6–7.0 meV
Dark energy-matter equality	9 Gyr	0.4	0.33 meV
Present	13.7 Gyr	0	0.24 meV



$$H = \sqrt{\frac{\rho}{3M_{\text{P}}^2}} = 1.66 \times g_*^{1/2} \frac{T^2}{M_{\text{P}}} \quad g_*(T) = \sum_{i=b} g_i \left(\frac{T_i}{T}\right)^4 + \frac{7}{8} \sum_{i=f} g_i \left(\frac{T_i}{T}\right)^4$$

$$\frac{t}{1 \text{ s}} \approx 2.42 g_*^{-1/2} \left(\frac{1 \text{ MeV}}{T}\right)^2$$

Radiation dominate Universe:  $H = (1/2t)$



## Bounce solution

$$S_3(T) = \int 4\pi r^2 dr \left[ \frac{1}{2} \left( \frac{d\phi_b}{dr} \right)^2 + V(\phi_b, T) \right]$$

$$\lim_{r \rightarrow \infty} \phi_b = 0, \quad \left. \frac{d\phi_b}{dr} \right|_{r=0} = 0$$

## Bubble nucleation

$$\Gamma \approx A(T) e^{-S_3/T} \sim 1$$

## PT strength

$$\alpha \equiv \frac{1}{\rho_r} \left( \Delta V_{\text{eff}}(\phi, T) - \frac{T}{4} \Delta \frac{\partial V_{\text{eff}}(\phi, T)}{\partial T} \right)$$

## Phase transition inverse duration

$$\frac{\beta}{H_n} = T \left. \frac{d(S_3(T)/T)}{dT} \right|_{T=T_n}$$

The probability, that a randomly chosen point is still in the false vacuum, given by

$$P(t) = e^{-I(t)} \quad I(t) = \frac{4\pi}{3} \int_{t_c}^t dt' \Gamma(t') a(t')^3 r(t, t')^3$$

The fraction of the space which has already been converted to the broken phase

$$r(t, t') = \int_{t'}^t \frac{v_w(\tilde{t}) d\tilde{t}}{a(\tilde{t})}$$

$r(t, t')$  : the comoving radius of a bubble nucleated at  $t'$  propagated until a subsequent time  $t$

$a(t)$ : the scale factor,  $v_w(t)$ : the wall velocity.

Using temperature  $T$  instead of time variable  $t$ , we have

$$I(T) = \frac{4\pi}{3} \int_T^{T_c} \frac{dT'}{H(T')} \Gamma(T') \frac{r(T, T')^3}{T'^4}$$

The transition completes when  $P(t) \approx 0.7$ , which leads to a percolation temperature  $T_p$  when

$$I(T_p) = 0.34.$$

- Bubble collisions**

$$\Omega_{\text{col}} h^2 = 1.67 \times 10^{-5} \left( \frac{H_*}{\beta} \right)^2 \left( \frac{\kappa \alpha}{1 + \alpha} \right)^2 \left( \frac{100}{g_*} \right)^{1/3} \left( \frac{0.11 v_b^3}{0.42 + v_b^2} \right) \frac{3.8 (f/f_{\text{env}})^{2.8}}{1 + 2.8 (f/f_{\text{env}})^{3.8}}$$

peak frequency:  $f_{\text{env}} = 16.5 \times 10^{-6} \left( \frac{f_*}{H_*} \right) \left( \frac{T_*}{100 \text{ GeV}} \right) \left( \frac{g_*}{100} \right)^{1/6} \text{ Hz}$

- Sound Wave**

$$\Omega h_{\text{sw}}^2(f) = 2.65 \times 10^{-6} (H_* \tau_{\text{sw}}) \left( \frac{\beta}{H} \right)^{-1} v_b \left( \frac{\kappa_\nu \alpha}{1 + \alpha} \right)^2 \left( \frac{g_*}{100} \right)^{-1/3} \left( \frac{f}{f_{\text{sw}}} \right)^3 \left( \frac{7}{4 + 3 (f/f_{\text{sw}})^2} \right)^{7/2}$$

phase transition duration:  $\tau_{\text{sw}} = \min \left[ \frac{1}{H_*}, \frac{R_*}{U_f} \right], H_* R_* = v_b (8\pi)^{1/3} (\beta/H)^{-1}$

Root-mean-square four-velocity of the plasma:

$$\bar{U}_f^2 \approx \frac{3}{4} \frac{\kappa_\nu \alpha}{1 + \alpha}$$

peak frequency:  $f_{\text{sw}} = 1.9 \times 10^{-5} \frac{\beta}{H} \frac{1}{v_b} \frac{T_*}{100} \left( \frac{g_*}{100} \right)^{1/6} \text{ Hz}$

- MHD turbulence**

$$\Omega h_{\text{turb}}^2(f) = 3.35 \times 10^{-4} \left( \frac{\beta}{H} \right)^{-1} \left( \frac{\epsilon \kappa_\nu \alpha}{1 + \alpha} \right)^{3/2} \left( \frac{g_*}{100} \right)^{-1/3} v_b \frac{(f/f_{\text{turb}})^3 (1 + f/f_{\text{turb}})^{-11/3}}{[1 + 8\pi f a_0 / (a_* H_*)]}$$

peak frequency:  $f_{\text{turb}} = 2.7 \times 10^{-5} \frac{\beta}{H} \frac{1}{v_b} \frac{T_*}{100} \left( \frac{g_*}{100} \right)^{1/6} \text{ Hz}$

# GW sources

$$\Omega_{\text{GW}}(f) = \begin{cases} \Omega_{\text{GW}^*} \left(\frac{f}{f_*}\right)^{n_{\text{GW}1}} & \text{for } f < f_*, \\ \Omega_{\text{GW}^*} \left(\frac{f}{f_*}\right)^{n_{\text{GW}2}} & \text{for } f > f_*, \end{cases}$$

**Table 1.** Cosmological GW sources

source	$n_{\text{GW}1}$	$n_{\text{GW}2}$	$f_*$ [Hz]	$\Omega_{\text{GW}}$
Phase transition (bubble collision)	2.8	-2	$\sim 10^{-5} \left(\frac{f_{\text{PT}}}{\beta}\right) \left(\frac{\beta}{H_{\text{PT}}}\right) \left(\frac{T_{\text{PT}}}{100 \text{ GeV}}\right)$	$\sim 10^{-5} \left(\frac{H_{\text{PT}}}{\beta}\right)^2 \left(\frac{\kappa_\phi \alpha}{1+\alpha}\right)^2 \left(\frac{0.11 v_w^3}{0.42 + v_w^2}\right)$
Phase transition (turbulence)	3	-5/3	$\sim 3 \times 10^{-5} \left(\frac{1}{v_w}\right) \left(\frac{\beta}{H_{\text{PT}}}\right) \left(\frac{T_{\text{PT}}}{100 \text{ GeV}}\right)$	$\sim 3 \times 10^{-4} \left(\frac{H_{\text{PT}}}{\beta}\right) \left(\frac{\kappa_{\text{turb}} \alpha}{1+\alpha}\right)^{3/2} v_w$
Phase transition (sound waves)	3	-4	$\sim 2 \times 10^{-5} \left(\frac{1}{v_w}\right) \left(\frac{\beta}{H_{\text{PT}}}\right) \left(\frac{T_{\text{PT}}}{100 \text{ GeV}}\right)$	$\sim 3 \times 10^{-6} \left(\frac{H_{\text{PT}}}{\beta}\right) \left(\frac{\kappa_v \alpha}{1+\alpha}\right)^2 v_w$
Preheating ( $\lambda\phi^4$ )	3	cutoff	$\sim 10^7$	$\sim 10^{-11} \left(\frac{g^2/\lambda}{100}\right)^{-0.5}$
Preheating (hybrid)	2	cutoff	$\sim \frac{g}{\sqrt{\lambda}} \lambda^{1/4} 10^{10.25}$	$\sim 10^{-5} \left(\frac{\lambda}{g^2}\right)^{1.16} \left(\frac{v}{M_{\text{pl}}}\right)^2$
Cosmic strings (loops 1)	[1, 2]	[-1, -0.1]	$\sim 3 \times 10^{-8} \left(\frac{G\mu}{10^{-11}}\right)^{-1}$	$\sim 10^{-9} \left(\frac{G\mu}{10^{-12}}\right) \left(\frac{\alpha_{\text{loop}}}{10^{-1}}\right)^{-1/2}$ (for $\alpha_{\text{loop}} \gg \Gamma G\mu$ )
Cosmic strings (loops 2)	[-1, -0.1]	0	$\sim 3 \times 10^{-8} \left(\frac{G\mu}{10^{-11}}\right)^{-1}$	$\sim 10^{-9.5} \left(\frac{G\mu}{10^{-12}}\right) \left(\frac{\alpha_{\text{loop}}}{10^{-1}}\right)^{-1/2}$ (for $\alpha_{\text{loop}} \gg \Gamma G\mu$ )
Cosmic strings (infinite strings)	[0, 0.2]	[0, 0.2]	—	$\sim 10^{-[11,13]} \left(\frac{G\mu}{10^{-8}}\right)$
Domain walls	3	-1	$\sim 10^{-9} \left(\frac{T_{\text{ann}}}{10^{-2} \text{ GeV}}\right)$	$\sim 10^{-17} \left(\frac{\sigma}{1 \text{ TeV}^3}\right)^2 \left(\frac{T_{\text{ann}}}{10^{-2} \text{ GeV}}\right)^{-4}$
Self-ordering scalar fields	0	0	—	$\sim \frac{511}{N} \Omega_{\text{rad}} \left(\frac{v}{M_{\text{pl}}}\right)^4$
Self-ordering scalar + reheating	0	-2	$\sim 0.4 \left(\frac{T_R}{10^7 \text{ GeV}}\right)$	$\sim \frac{511}{N} \Omega_{\text{rad}} \left(\frac{v}{M_{\text{pl}}}\right)^4$
Magnetic fields	3	$\alpha_B + 1$	$\sim 10^{-6} \left(\frac{T_*}{10^2 \text{ GeV}}\right)$	$\sim 10^{-16} \left(\frac{B}{10^{-10} \text{ G}}\right)$
Inflation+reheating	$\sim 0$	-2	$\sim 0.3 \left(\frac{T_R}{10^7 \text{ GeV}}\right)$	$\sim 2 \times 10^{-17} \left(\frac{r}{0.01}\right)$
Inflation+kination	$\sim 0$	1	$\sim 0.3 \left(\frac{T_R}{10^7 \text{ GeV}}\right)$	$\sim 2 \times 10^{-17} \left(\frac{r}{0.01}\right)$
Particle prod. during inf.	$-2\epsilon$	$-4\epsilon(4\pi\xi - 6)(\epsilon - \eta)$	—	$\sim 2 \times 10^{-17} \left(\frac{r}{0.01}\right)$
2nd-order (inflation)	1	drop-off	$\sim 7 \times 10^5 \left(\frac{T_{\text{reh}}}{10^9 \text{ GeV}}\right)^{1/3} \left(\frac{M_{\text{inf}}}{10^{16} \text{ GeV}}\right)^{2/3}$	$\sim 10^{-12} \left(\frac{T_{\text{reh}}}{10^9 \text{ GeV}}\right)^{-4/3} \left(\frac{M_{\text{inf}}}{10^{16} \text{ GeV}}\right)^{4/3}$
2nd-order (PBHs)	2	drop-off	$\sim 4 \times 10^{-2} \left(\frac{M_{\text{PBH}}}{10^{20} \text{ g}}\right)^{-1/2}$	$\sim 7 \times 10^{-9} \left(\frac{\mathcal{A}^2}{10^{-3}}\right)^2$
Pre-Big-Bang	3	$3 - 2\mu$	—	$\sim 1.4 \times 10^{-6} \left(\frac{H_s}{0.15 M_{\text{pl}}}\right)^4$


## Magnetic Field and Gravitational Waves from the First-Order Phase Transition

Yuefeng Di, Jialong Wang, Ruiyu Zhou, and Ligong Bian<sup>\*</sup>  
*Department of Physics, Chongqing University, Chongqing 401331, China*

Rong-Gen Cai<sup>†</sup>  
*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences,  
P.O. Box 2735, Beijing 100190, China,  
School of Physical Sciences, University of Chinese Academy of Sciences, No. 19A Yuquan Road, Beijing 100049, China,  
and School of Fundamental Physics and Mathematical Sciences, Hangzhou Institute for Advanced Study,  
University of Chinese Academy of Sciences, Hangzhou 310024, China*


Jing Liu<sup>‡</sup>  
*School of Fundamental Physics and Mathematical Sciences, Hangzhou Institute for Advanced Study,  
University of Chinese Academy of Sciences, Hangzhou 310024, China  
and School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China*

### Finite-T Veff




**Finite-T calculation**

### Nucleation



**Lattice Simulation**

### Expansion&Percolation





# Lattice EW field foundation

$\Phi(t, \mathbf{x})$  : Higgs field doublet defined on sites;

$U_i(t, \mathbf{x})$  and  $V_i(t, \mathbf{x})$  : SU(2) and U(1) link fields, defined on the link between the neighboring sites  $\mathbf{x}$  and  $\mathbf{x} + \mathbf{i}$ ,  $\Phi(t, \mathbf{x})$ ,  $U_i(t, \mathbf{x})$  and  $V_i(t, \mathbf{x})$  are defined at time steps  $t + \Delta t, t + 2\Delta t, \dots$ ;


Conjugate momentum fields:  $\Pi(t+\Delta t/2, \mathbf{x})$ ,  $F(t+\Delta t/2, \mathbf{x})$  and  $E(t+\Delta t/2, \mathbf{x})$ , are defined at time steps  $t + \Delta t/2, t + 3\Delta t/2$ .

$$U_i(t, \mathbf{x}) = \exp\left(-\frac{i}{2}g\Delta x\sigma^a W_i^a\right)$$

$$U_0(t, \mathbf{x}) = \exp\left(-\frac{i}{2}g\Delta t\sigma^a W_0^a\right)$$

$$V_i(t, \mathbf{x}) = \exp\left(-\frac{i}{2}g\Delta x B_i\right)$$

$$V_0(t, \mathbf{x}) = \exp\left(-\frac{i}{2}g\Delta t B_0\right).$$



$$D_i\Phi = \frac{1}{\Delta x} [U_i(t, \mathbf{x})V_i(t, \mathbf{x})\Phi(t, \mathbf{x} + \mathbf{i}) - \Phi(t, \mathbf{x})]$$

$$D_0\Phi = \frac{1}{\Delta t} [U_0(t, \mathbf{x})V_0(t, \mathbf{x})\Phi(t + \Delta t, \mathbf{x}) - \Phi(t, \mathbf{x})].$$

$$\Phi(t + \Delta t, \mathbf{x}) = \Phi(t, \mathbf{x}) + \Delta t\Pi(t + \Delta t/2, \mathbf{x})$$

$$V_i(t + \Delta t, \mathbf{x}) = \frac{1}{2}g'\Delta x\Delta t E_i(t + \Delta t/2, \mathbf{x})V_i(t, \mathbf{x})$$


$$U_i(t + \Delta t, \mathbf{x}) = g\Delta x\Delta t F_i(t + \Delta t/2, \mathbf{x})U_i(t, \mathbf{x}),$$

Temporal gauge  
 $U_0(t, \mathbf{x}) = \mathbf{I}_2, V_0(t, \mathbf{x}) = 1$

leapfrog


## Field basis

$$\begin{aligned}\partial_0^2 \Phi &= D_i D_i \Phi - 2\lambda(|\Phi|^2 - \eta^2)\Phi - 3(\Phi^\dagger \Phi)^2 \Phi / \Lambda^2, \\ \partial_0^2 B_i &= -\partial_j B_{ij} + g' \text{Im}[\Phi^\dagger D_i \Phi], \\ \partial_0^2 W_i^a &= -\partial_k W_{ik}^a - g \epsilon^{abc} W_k^b W_{ik}^c + g \text{Im}[\Phi^\dagger \sigma^a D_i \Phi], \\ \partial_0 \partial_j B_j - g' \text{Im}[\Phi^\dagger \partial_0 \Phi] &= 0, \\ \partial_0 \partial_j W_j^a + g \epsilon^{abc} W_j^b \partial_0 W_j^c - g \text{Im}[\Phi^\dagger \sigma^a \partial_0 \Phi] &= 0.\end{aligned}$$



## Lattice implementation

$$\begin{aligned}\Pi(t + \Delta t/2, x) &= \Pi(t - \Delta t/2, x) + \Delta t \left\{ \frac{1}{\Delta x^2} \sum_i [U_i(t, x) V_i(t, x) \Phi(t, x + i) \right. \\ &\quad \left. - 2\Phi(t, x) + U_i^\dagger(t, x - i) V_i^\dagger(t, x - i) \Phi(t, x - i)] - \frac{\partial U}{\partial \Phi^\dagger} \right\} \\ \text{Im}[E_k(t + \Delta t/2, x)] &= \text{Im}[E_k(t - \Delta t/2, x)] + \Delta t \left\{ \frac{g'}{\Delta x} \text{Im}[\Phi^\dagger(t, x + k) U_k^\dagger(t, x) V_k^\dagger(t, x) \Phi(t, x)] \right. \\ &\quad \left. - \frac{2}{g' \Delta x^3} \sum_i \text{Im}[V_k(t, x) V_i(t, x + k) V_k^\dagger(t, x + i) V_i^\dagger(t, x) \right. \\ &\quad \left. + V_i(t, x - i) V_k(t, x) V_i^\dagger(t, x + k - i) V_k^\dagger(t, x - i)] \right\} \\ \text{Tr}[i\sigma^m F_k(t + \Delta t/2, x)] &= \text{Tr}[i\sigma^m F_k(t - \Delta t/2, x)] + \Delta t \left\{ \frac{g}{\Delta x} \text{Re}[\Phi^\dagger(t, x + k) U_k^\dagger(t, x) V_k^\dagger(t, x) i\sigma^m \Phi(t, x)] \right. \\ &\quad \left. - \frac{1}{g \Delta x^3} \sum_i \text{Tr}[i\sigma^m U_k(t, x) U_i(t, x + k) U_k^\dagger(t, x + i) U_i^\dagger(t, x) \right. \\ &\quad \left. + i\sigma^m U_k(t, x) U_i^\dagger(t, x + k - i) U_k^\dagger(t, x - i) U_i(t, x - i)] \right\},\end{aligned}$$



Di, Wang, Zhou, **Bian\***, Cai\*, Liu\*, Phys.Rev.Lett. 126 (2021) 251102



# GW from Bubble collisions

$$\ddot{h}_{ij} - \nabla^2 h_{ij} = 16\pi G T_{ij}^{TT}$$


$$T_{\mu\nu} = \partial_\mu \Phi^\dagger \partial_\nu \Phi - g_{\mu\nu} \frac{1}{2} \text{Re}[(\partial_i \Phi^\dagger \partial^i \Phi)^2]$$

$$\langle \dot{h}_{ij}^{TT}(\mathbf{k}, t) \dot{h}_{ij}^{TT}(\mathbf{k}', t) \rangle = P_h(\mathbf{k}, t) (2\pi)^3 \delta(\mathbf{k} + \mathbf{k}')$$

$$\frac{d\Omega_{\text{gw}}}{d\ln(k)} = \frac{1}{32\pi G \rho_c} \frac{k^3}{2\pi^2} P_h(\mathbf{k}, t)$$



$$\gamma^* = R^* / (2Rc) \quad R^* = \left( \frac{\nu}{N_b} \right)^{1/3}$$



Di, Wang, Zhou, **Bian\***, Cai\*, Liu\*, Phys.Rev.Lett. 126 (2021) 251102



# scalar field + fluid system

The full energy-momentum tensor into two components, one for the fluid  $T^{\mu\nu}$  and one for the Higgs  $T^{\mu\nu}$

$$T_f^{\mu\nu} = (e + p_1)u^\mu u^\nu + p_1 g^{\mu\nu},$$

$$T_\phi^{\mu\nu} = \partial^\mu \phi \partial^\nu \phi - g^{\mu\nu} \left( \frac{1}{2} (\partial \phi)^2 + V_0(\phi) \right),$$

Fluid pressure is the total contribution from all particles

$$p_1(\phi, T) = \frac{\pi^2}{90} g_{\text{eff}} T^4 - V_1(\phi, T) = - \sum_B f_B(m(\phi), T) - \sum_F f_F(m(\phi), T)$$

Thermally corrected potential

$$V_T(\phi) = V_0(\phi) + V_1(\phi, T)$$

$$\partial_\mu T_\phi^{\mu\nu} = + \partial^\nu \phi \frac{dm^2}{d\phi} \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_{\vec{p}}} f(p, x). \quad \Rightarrow \quad \square \phi - V'_T(\phi) = - \frac{dm^2}{d\phi} \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_{\vec{p}}} \delta f(p, x)$$

Boltzmann equation with collisions and external forces

$$f(p, x) = f^{\text{eq}}(p, x) + \delta f(p, x)$$

This leaves us with the equation of motion

$$\square \phi - V'_T(\phi) = -\tilde{\eta} \frac{dm^2}{d\phi} u^\mu \partial_\mu \phi \quad \text{where} \quad \int \frac{d^3 p}{(2\pi)^3} \frac{1}{2E_{\vec{p}}} \delta f(p, x) = \tilde{\eta} u^\mu \partial_\mu \phi,$$

Local equilibrium and perfect fluid

With  $p(\phi, T) = p_1(\phi, T) - V_0(\phi)$ , we have  $T_f^{\mu\nu} = (e + p)u^\mu u^\nu + pg^{\mu\nu}$ ,

The full energy-momentum tensor is conserved  $\partial_\mu (T_f^{\mu\nu} + T_\phi^{\mu\nu}) = 0$

which yields

$$\partial_\mu T_f^{\mu\nu} + V_T'(\phi)\partial^\nu\phi = -\partial^\nu\phi \frac{dm^2}{d\phi} \int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_{\vec{p}}} \delta f(p, x)$$

Consider the scalar product of  $u$  with both sides

$$u_\nu \partial_\mu (wu^\mu u^\nu + pg^{\mu\nu}) + V_T'(\phi)u \cdot \partial\phi = \tilde{\eta}(u \cdot \partial\phi)^2$$

Here,  $w = e + p = Ts$  is the enthalpy density, and  $s = dp/dT$  is the entropy density



Hydrodynamic equations for a single bubble

$$(\xi - v) \frac{\partial_\xi e}{w} = 2 \frac{v}{\xi} + [1 - \gamma^2 v(\xi - v)] \partial_\xi v,$$

$$(1 - v\xi) \frac{\partial_\xi p}{w} = \gamma^2 (\xi - v) \partial_\xi v.$$

$\xi = r/t$ ,  $r$  is the distance from the center of the bubble and  $t$  is the time since nucleation,

$v(\xi_w)$  is the fluid velocity at the location of the bubble wall and  $\xi_w = v_w$  is the wall velocity.

Introducing sound speed  $c_s^2 \equiv (dp/dT)/(de/dT)$ , we have

$$2 \frac{v}{\xi} = \gamma^2 (1 - v\xi) \left[ \frac{\mu^2}{c_s^2} - 1 \right] \partial_\xi v,$$

$$\mu(\xi, v) = \frac{\xi - v}{1 - \xi v}$$

$\mu$  is the fluid velocity at  $\xi$  in a frame that is moving outward at speed  $\xi$

From the conservation of the energy momentum tensor accross the interface, we have the boundary conditions:

$$v_+ v_- = \frac{p_+ - p_-}{e_+ - e_-}, \quad \frac{v_+}{v_-} = \frac{e_- + p_+}{e_+ + p_-},$$

+(-) meaning in front (behind) of the bubble wall (in the rest frame of the bubble wall) .

# Bubble dynamics and fluid @ FOPT


symmetric phase  $p_+ = \frac{1}{3}a_+T_+^4 - \epsilon, \quad e_+ = a_+T_+^4 + \epsilon,$

broken phase  $p_- = \frac{1}{3}a_-T_-^4, \quad e_- = a_-T_-^4,$

Bag equation of state


the false-vacuum energy resulting from the Higgs potential

Different number of light degrees of freedom across the wall, different values  $a_+$  and  $a_-$  (with  $a_+ > a_-$ ) and different temperatures on both sides of the wall



The black circle is the phase interface (bubble wall).

Green region is of non-zero fluid velocity



deflagration walls have a shock wave propagating in front of the wall,  
 detonation walls have a rarefaction wave behind it,  
 hybrid walls have both shock and rarefaction waves.

# 真空泡碰撞、合并、流体演化产生引力波

## 有限温度有效势能

$$V(\phi, T) = \frac{1}{2}\gamma(T^2 - T_0^2)\phi^2 - \frac{1}{3}AT\phi^3 + \frac{1}{4}\lambda\phi^4$$

新物理

## 标量场-相对论流体运动方程

$$-\ddot{\phi} + \nabla^2\phi - \frac{\partial V}{\partial\phi} = \eta W(\dot{\phi} + V^i\partial_i\phi) \quad \eta: \text{粒子和真空泡壁相互作用}$$

$$\begin{aligned} \dot{E} + \partial_i(EV^i) + p[\dot{W} + \partial_i(WV^i)] - \frac{\partial V}{\partial\phi}W(\dot{\phi} + V^i\partial_i\phi) \\ = \eta W^2(\dot{\phi} + V^i\partial_i\phi)^2 \end{aligned}$$



$$\dot{Z}_i + \partial_j(Z_iV^j) + \partial_i p + \frac{\partial V}{\partial\phi}\partial_i\phi = -\eta W(\dot{\phi} + V^j\partial_j\phi)\partial_i\phi$$

equation of state  $\epsilon(T, \phi) = 3aT^4 + V(\phi, T) - T\frac{\partial V}{\partial T},$

$$p(T, \phi) = aT^4 - V(\phi, T)$$

fluid momentum density  $Z_i = W(\epsilon + p)U_i$


fluid energy density  $E = W\epsilon$



$V^i$  is the fluid 3-velocity

$U^i = WV^i$ ,  $W$ : relativistic  $\gamma$ -factor


# 真空泡碰撞、合并、流体演化产生引力波



$$\tau_{ij}^\phi = \partial_i \phi \partial_j \phi, \quad \tau_{ij}^f = W^2 (\epsilon + p) V_i V_j$$

$$(\bar{\epsilon} + \bar{p}) \bar{U}_f^2 = \frac{1}{V} \int_V d^3x \tau_{ii}^f$$

$$(\bar{\epsilon} + \bar{p}) \bar{U}_\phi^2 = \frac{1}{V} \int_V d^3x \tau_{ii}^\phi$$





$$\ddot{u}_{ij} - \nabla^2 u_{ij} = 16\pi G (\tau_{ij}^\phi + \tau_{ij}^f)$$

$$h_{ij}(\mathbf{k}) = \lambda_{ij,lm}(\hat{\mathbf{k}}) u_{lm}(t, \mathbf{k})$$



Bian, Jia, Zhao, Zhu, arxiv:2011.XXXXX

# 正反物质不对称&强一阶电弱相变

## 电弱重子数产生机制



Chup et al, Rev Mod Phys.91.015001



Bian, Liu, Shu, Phys.Rev.Lett. 115 (2015) 021801



# Bubble wall velocity with the EW plasma

Boltzmann equation which dictates the time evolution of the particle distribution

Wall frame

$$\frac{df_a}{dt} = \partial_t f_a + \dot{\vec{x}} \cdot \partial_{\vec{x}} f_a + \dot{\vec{p}} \cdot \partial_{\vec{p}} f_a = C[f_a], \quad (\text{a})$$

The *fluid ansatz* for the distribution function is written as

$$f \approx f_v - f'_v \delta \bar{X} + \delta f_u + \mathcal{O}(\delta f^2), \quad f_v = \frac{1}{e^{\beta\gamma(E-vp_z)} \pm 1}, \quad f'_v \equiv \frac{df_v}{d\beta\gamma E},$$

$$\delta \bar{X} = \mu + \beta\gamma\delta\tau(E - vp_z) \quad \text{perturbations from equilibrium}$$

$\mu$ : chemical potential,  $\delta\tau$ : temperature perturbation,  $\delta f_u$ : the velocity perturbation

The force and group velocity

$$\dot{z} \equiv \frac{\partial\omega}{\partial p_z} = \frac{p_z}{E} + s \frac{m^2\theta'}{2E^2 E_z}, \quad \dot{p}_z \equiv -\frac{\partial\omega}{\partial z} = -\frac{(m^2)'}{2E} + s \frac{(m^2\theta)'}{2E E_z}, \quad (\text{b})$$

$\omega$  is the energy of the WKB wave packet and  $E_z^2 \equiv p_z^2 + m^2$

Inserting the force and group velocity of eq. (b) into the Boltzmann equation (a), we have

$$\left[ \frac{p_z}{E} \partial_z - \frac{(m^2)'}{2E} \partial_{p_z} \right] (f_v - f'_v \delta \bar{X} + \delta f_u) = C[f]$$

# Bubble wall velocity with the EW plasma

With  $q = (\mu, \delta\tau, u)^T$ , the transport equations take the form

$$A_v \vec{q}' + \Gamma \vec{q} = S,$$

$\Gamma$ : the collision term  $C$  in the above equation

$$A_v = \begin{pmatrix} C_v^{1,1} & \gamma v C_0^{-1,0} & D_v^{0,0} \\ C_v^{0,1} & \gamma(C_v^{-1,1} - v C_v^{0,2}) & D_v^{-1,0} \\ C_v^{2,2} & \gamma(C_v^{1,2} - v C_v^{2,3}) & D_v^{1,1} \end{pmatrix},$$

Integrals of the particle distribution functions.

$$S = \gamma v \frac{(m^2)'}{2T^2} \begin{pmatrix} C_v^{1,0} \\ C_v^{0,0} \\ C_v^{2,1} \end{pmatrix},$$

Source term

The Higgs EOM in the presence of out of equilibrium particle populations

$$E_h \equiv \square\phi + \frac{dV_{\text{eff}}(\phi, T)}{d\phi} + \sum_i \frac{dm_i^2}{d\phi} \int \frac{d^3p}{(2\pi)^3} \frac{\delta f_i(p, x)}{2E} = 0,$$

$$M_1 \equiv \int dz E_h h' dz = 0,$$

The total pressure on the wall should be zero

$$M_2 \equiv \int dz E_h h' [2h(z) - h_0] dz = 0.$$

An asymmetry in the total pressure between the front and back of the wall should be zero

$$h(z) = \frac{h_0}{2} \left[ \tanh\left(\frac{z}{L_h}\right) + 1 \right]$$

Bubble profile

# EWBG with the EW plasma

Boltzmann equation

$$(v_g \partial_z + F \partial_{p_z}) f = \mathcal{C}[f]$$

$$v_g = \frac{p_z}{E_w},$$

CP-violating complex mass term

$$\hat{m}(z) = m(z) e^{i\gamma^5 \theta(z)}$$

$$F = -\frac{(m^2)'}{2E_w} + s s_{k_0} \frac{(m^2 \theta)'}{2E_w E_{wz}},$$

$$\mu \equiv \mu_e + s_{k_0} \mu_o,$$

$$\delta f \equiv \delta f_e + s_{k_0} \delta f_o.$$


Transport equations

$$A w' + (m^2)' B w = S + \delta C,$$

collision terms

$$w = (\mu, u)^T$$

$$A = \begin{pmatrix} -D_1 & 1 \\ -D_2 & R \end{pmatrix}, \quad B = \begin{pmatrix} -v_w \gamma_w Q_1 & 0 \\ -v_w \gamma_w Q_2 & \bar{R} \end{pmatrix},$$



Source term  $S = (S_1, S_2)^T$

$$S_{hl}^o = v_w \gamma_w s [(m^2 \theta')' Q_\ell^{8o} - (m^2)' m^2 \theta' Q_\ell^{9o}],$$

chemical potential for left handed baryon number


$$\mu_{B_L} = \frac{1}{2} (1 + 4D_0^t) \mu_{t_L} + \frac{1}{2} (1 + 4D_0^b) \mu_{b_L} + 2D_0^t \mu_{t_R}$$

Baryon asymmetry

$$\eta_B = \frac{405 \Gamma_{\text{sph}}}{4\pi^2 v_w \gamma_w g_* T} \int dz \mu_{B_L} f_{\text{sph}} e^{-45 \Gamma_{\text{sph}} |z| / 4v_w},$$

VEV- insertion source tends to predict a larger baryon asymmetry than the WKB source by a factor of  $\sim 10$ .

# Freeze-in and Freeze-out




$T_{fi} \sim m_{DM}/x_{fi} \sim 100 \text{ GeV}$  for  $m_{DM} \sim \text{weak scale}$

$T_{fo} \sim m_{DM}/x_{fo} \sim O(1) \text{ GeV}$  for  $m_{DM} \sim \text{weak scale}$




## WIMP DM+EWBG

CxSM Jiang, **Bian\***, Huang, Shu 16,  
Chiang, Ramsey-Musolf, Senaha  
18,...




**Two-step FOPT**


## WIMP DM+FOPT




**DM**  
**one-step**  
**two-step**  
**BAU**






Jiang, **Bian\***, Huang, Shu 16



SM+2 real scalars **Chao, Guo, Shu 17,...**







# FIMP DM and Two-step FOPT



1712.03962, Michael J. Baker et al.

**Amplified effects**  
larger thermal  
masses before PT



**Temporarily open**  
of decay channel


**Bian**, Tang, 18

# Collider search for 2step FOPT


## Zh@ILC/CEPC

$$V_0 = -\mu^2|H|^2 + \lambda|H|^4 + \frac{1}{2}\mu_S^2 S^2 + \lambda_{HS}|H|^2 S^2 + \frac{1}{4}\lambda_S S^4$$


$$V_{\text{eff}}(h, T) = V_0(h) + V_0^{CW}(h) + V_T(h, T) + V_r(h, T)$$



Craig, Englert, and McCullough, 1305.5251





Curtin, Meade, Yu, 1409.0005



Huang, Long, and Wang, 1608.06619

# ► Collider search for 2 step FOPT

## ● Off-shell Higgs@LHC



Goncalves, Han, and Mukhopadhyay, 1710.02149

See also: Lee, Park, and Qian, 1812.02679




# LIGO-Virgo search for FOPT

## High-scale PT

Romero, Martinovic, Callister, Guo, et al., Phys.Rev.Lett. 126 (2021) 15, 151301

LIGO-Virgo O3



# ▶ PPTA search for FOPT

## ■ PPTA DR2 dataset constrain low-scale phase transition, dark sector and QCD scale FOPT

PHYSICAL REVIEW LETTERS 127, 251303 (2021)


Editors' Suggestion    Featured in Physics

### Constraining Cosmological Phase Transitions with the Parkes Pulsar Timing Array

Xiao Xue<sup>1,2,3</sup>, Ligong Bian<sup>4,5,\*</sup>, Jing Shu<sup>1,2,6,7,8,†</sup>, Qiang Yuan<sup>9,10,7,‡</sup>, Xingjiang Zhu<sup>11,12,13,§</sup>, N. D. Ramesh Bhat<sup>14</sup>,  
 Shi Dai<sup>15</sup>, Yi Feng<sup>16</sup>, Boris Goncharov<sup>11,12</sup>, George Hobbs<sup>17</sup>, Eric Howard<sup>17,18</sup>, Richard N. Manchester<sup>17</sup>,  
 Christopher J. Russell<sup>19</sup>, Daniel J. Reardon<sup>12,20</sup>, Ryan M. Shannon<sup>12,20</sup>, Renée Spiewak<sup>21,20</sup>,  
 Nithyanandan Thyagarajan<sup>22</sup> and Jingbo Wang<sup>23</sup>


TABLE I: Description of hypotheses tested in this work and the Bayes factors between them.

Hypothesis	Pulsar noise	Common red process	HD process FOPT spectrum	Bayes Factors	Parameter Estimation (median and 1- $\sigma$ interval)	
					$T_*/\text{MeV}, \alpha \times 10^3, \beta/H_*$	$A_{\text{comred}}, \gamma_{\text{comred}}$
H0:Pulsar Noise	yes	no	no			
H1:Common Red	yes	yes	no	$10^{3.5}$ (against H0)		$-14.45^{+0.62}_{-0.64}, 3.31^{+1.36}_{-1.53}$
H2:FOPT	yes	no	yes (full HD)	$10^{1.8}$ (against H0)	$7.4^{+11.9}_{-4.7}, 271^{+165}_{-92}, 9.9^{+11.4}_{-5.4}$	
H3:FOPT1	yes	yes	yes (full HD)	1.04 (against H1)	$9.6^{+232.2}_{-9.2}, 3.8^{+27.9}_{-3.4}, 854^{+9622}_{-782}$	$-14.51^{+0.64}_{-0.68}, 3.36^{+1.39}_{-1.54}$
H4:FOPT2	yes	yes	yes (no-auto HD)	0.96 (against H1)	$10.9^{+290.5}_{-10.6}, 3.2^{+19.9}_{-2.8}, 1053^{+11256}_{-962}$	$-14.45^{+0.62}_{-0.64}, 3.27^{+1.37}_{-1.54}$



# Two-step FOPT potential

## Type-a




$$V_a(\phi, h, T) = \frac{1}{2}(\mu_\phi^2 + c_\phi T^2)\phi^2 + \frac{1}{2}\lambda_{h\phi}h^2\phi^2 + \frac{1}{4}\lambda_\phi\phi^4$$

$$+ \frac{1}{2}(-\mu_h^2 + c_h T^2)h^2 + \frac{1}{4}\lambda_h h^4$$

$$c_\phi = \lambda_\phi/4 + \lambda_{h\phi}/3$$


$$c_h = (2m_W^2 + m_Z^2 + 2m_t^2)/(4v^2) + \lambda_h/2 + \lambda_{h\phi}/12$$



Motivated for DM&EWBG, see:1804.06813,1702.06124,1609.07143, 1605.08663, 1605.08663,etc



# Two-step PT with the second-step being FOPT

**Type-a**



$$h(t=0, r) = \eta_h/2 \left[ 1 - \tanh\left(\frac{r - R_0}{L_w}\right) \right]$$


$$\phi(t=0, r) = \eta_\phi/2 \left[ 1 + \tanh\left(\frac{r - R_0}{L_w}\right) \right]$$





# Two-step PT with the second-step being FOPT


Type-a



# Two-step PT with first-step being FOPT

**Type-b**

Without Global U(1)





$$V_{cct}(\phi, T) = a\phi^4(\log[|\phi|^2/v_\phi^2] - 1/4) + bT^2|\phi|^2$$

$$V_{dt}(\phi, h, T) = \frac{1}{2}c'_h T^2 h^2 + \frac{1}{4}\lambda_h h^4 - \frac{\lambda_p}{4} h^2 \phi^2$$

$$c'_h = (2m_W^2 + m_Z^2 + 2m_t^2)/(4v^2) + \lambda_h/2 + \lambda_p/24$$

$$\langle h \rangle = \sqrt{(\lambda_p \eta^2 - 2c'_h T^2)/(2\lambda_h)}$$


Classical conformal + Dimensional transmutation



# Two-step PT with first-step being FOPT



**Type-b**

Without Global U(1)



$$\phi(t=0, r) = \eta_\phi/2 \left[ 1 - \tanh\left(\frac{r - R_0}{L_w}\right) \right]$$


$$\langle h \rangle = \sqrt{(\lambda_p \eta^2 - 2c'_h T^2)/(2\lambda_h)}$$



# Two-step PT with first-step being FOPT


Type-b

Without Global U(1)




# Cosmic string


## 通常形成于GUTs



## 相变后U(1) 对称性自发破缺




## 宇宙弦




T. W. B. Kibble

## 宇宙弦成圈



Phys.Rev.D 30 (1984) 2036

## 引力波辐射



Yann Gouttenoire et al JCAP07(2020)032



# NANOGrav 12.5-yr dataset & cosmic string


**NANOGrav**

- Data Products
- Events & Meetings
- Outreach
- Publications
- People
- Institutions
- Telescopes
- Governance
- Press

## North American Nanohertz Observatory for Gravitational Waves

### NANOGrav Finds Possible 'First Hints' of Low-Frequency Gravitational Wave Background

Press Images



## The NANOGrav 12.5 yr Data Set: Search for an Isotropic Stochastic Gravitational-wave Background

#1

NANOGrav Collaboration • Zaven Arzoumanian (CRESST, Greenbelt and NASA, Goddard) et al. (Sep 9, 2020)

Published in: *Astrophys.J.Lett.* 905 (2020) 2, L34 • e-Print: [2009.04496](https://arxiv.org/abs/2009.04496) [astro-ph.HE]

[pdf](#) [links](#) [DOI](#) [cite](#)

[↻](#) 411 citations

## On the Evidence for a Common-spectrum Process in the Search for the Nanohertz Gravitational-wave Background with the Parkes Pulsar Timing Array

#2


Boris Goncharov (Swinburne U., Ctr. Astrophys. Supercomput. and Monash U. and GSSI, Aquila and Gran Sasso), R.M. Shannon (Swinburne U., Ctr. Astrophys. Supercomput. and Monash U.), D.J. Reardon (Swinburne U., Ctr. Astrophys. Supercomput. and Monash U.), G. Hobbs (Australia, CSIRO, Epping), A. Zic (Australia, CSIRO, Epping and Macquarie U.) et al. (Jul 26, 2021)

Published in: *Astrophys.J.Lett.* 917 (2021) 2, L19 • e-Print: [2107.12112](https://arxiv.org/abs/2107.12112) [astro-ph.HE]


[pdf](#) [DOI](#) [cite](#)

[↻](#) 94 citations

# NANOGrav 12.5-yr dataset & cosmic string



John Ellis et al, Phys.Rev.Lett.126.041304(2021)



Bian, Cai, Liu, Zhou, Phys.Rev.D 103 (2021) 8, L081301



# ► PPTA dataset & cosmic string

TABLE I: Hypotheses, Bayes factors, and estimated model parameters for the BOS model.

Hypothesis	Pulsar Noise	CPL Process	HD process CS spectrum	Bayes Factors	Parameter Estimation ( $1\sigma$ interval)	
					$\log_{10} G\mu$	$\log_{10} A_{\text{CPL}}, \gamma_{\text{CPL}}$
H0:Pulsar Noise	✓					
H1:CPL	✓	✓		$10^{3.2}$ (/H0)		$-14.48^{+0.62}_{-0.64}, 3.34^{+1.37}_{-1.53}$
H2:CS	✓		✓(full HD)	$10^{3.1}$ (/H0)	$-10.38^{+0.21}_{-0.21}$	
H3:CS1	✓	✓	✓(full HD)	1.96 (/H1)	$< -10.02$ (95% C.L.)	$-15.58^{+1.21}_{-1.64}, 3.11^{+1.95}_{-2.02}$
H4:CS2	✓	✓	✓(no-auto HD)	0.60 (/H1)	$< -10.54$ (95% C.L.)	$-14.61^{+0.58}_{-0.59}, 3.63^{+1.24}_{-1.40}$

TABLE II: Hypotheses, Bayes factors, and estimated model parameters for the LRS model.


Hypothesis	Pulsar Noise	CPL process	HD process CS spectrum	Bayes Factors	Parameter Estimation ( $1\sigma$ interval)	
					$\log_{10} G\mu$	$\log_{10} A_{\text{CPL}}, \gamma_{\text{CPL}}$
H0:Pulsar Noise	✓					
H1:CPL	✓	✓		$10^{3.2}$ (/H0)		$-14.48^{+0.62}_{-0.64}, 3.34^{+1.37}_{-1.53}$
H2:CS	✓		✓(full HD)	$10^{3.3}$ (/H0)	$-10.89^{+0.14}_{-0.17}$	
H3:CS1	✓	✓	✓(full HD)	1.62 (/H1)	$< -10.64$ (95% C.L.)	$-15.44^{+1.18}_{-1.74}, 3.08^{+1.94}_{-1.99}$
H4:CS2	✓	✓	✓(no-auto HD)	0.55 (/H1)	$< -11.04$ (95% C.L.)	$-14.57^{+0.58}_{-0.59}, 3.54^{+1.24}_{-1.41}$




Bian\*, Shu\*, Wang, Yuan\*, Zong, 2205.07293

CS: SSB of U(1) symmetry

The one-dimension topological defects: cosmic string



Potential



T. W. B. Kibble




FIG. 1. Three bubbles of the broken symmetry phase ( $\rho = \eta$ ) colliding. If the phase change of the scalar field around the loop  $\gamma$  is  $\pm 2\pi$ , a string (or antistring) is formed. If the phases  $\alpha_i$  are ordered, then the requirement for a string is  $\alpha_1 + \pi < \alpha_3 < \alpha_2 + \pi$ .

$\xi_{str}$  of the string network is essentially the typical bubble diameter for SFOPT???



Aust. J. Phys., 1997, 50, 697–722

Phys.Rev.D 49 (1994) 1944-1950





**With Global U(1)**

Motivated for strong CP and axion DM



$t/R^* = 0.43$







Zhao, Di, **Bian\***, Cai\*, 2204.04427







# ► Bubbles and vortex&anti-vortex

**Type-b**

**With Global U(1)**



Arrows: phase distribution









# Zombie DM and CS GW

$$\mathcal{L}_{B-L} = \sum_i \bar{\nu}_R^i i \not{D} \nu_R^i - \frac{1}{2} \sum_{i,j} \left( \lambda_R^{ij} \bar{\nu}_R^{i,c} \Phi \nu_R^j + \text{h.c.} \right) - \sum_{i,j} \left( \lambda_D^{ij} \bar{\ell}_L^i \tilde{H} \nu_R^j + \text{h.c.} \right) \\ + D_\mu \Phi^\dagger D^\mu \Phi - \lambda_\phi \left( |\Phi|^2 - \frac{v_\phi^2}{2} \right)^2 - \frac{1}{4} Z'_{\mu\nu} Z'^{\mu\nu},$$

	$\nu_R^{1,2}$	$\nu_R^3$	$\Phi$	$\psi$	$S$
$B-L$	-1	-1	2	-1	0
$\mathbb{Z}_2$	1	-1	1	-1	1

$$\mathcal{L}_{\text{DM}} = \bar{\psi} (i \not{D} - M_\psi) \psi + \frac{1}{2} \partial_\mu S \partial^\mu S - \frac{1}{2} M_S^2 S^2 + \left( \lambda_1 S \bar{\psi} \nu_R + \text{h.c.} \right) + \lambda_2 S \bar{\psi} \psi,$$



zombie collision


$$\nu_R + \psi \rightarrow \psi + \psi, \quad \nu_R + \psi \rightarrow \psi + \bar{\psi},$$

**Entropy dilution**

$$\mathcal{L}_6 = \frac{1}{\Lambda^2} \sum_{i,j,j'} (\bar{\psi}^c u_R^i) (\bar{d}_R^{j,c} d_R^{j'}) + \text{h.c.}$$

Bian, Liu, Xie, 21

# Zombie DM and CS GW



Bian, Liu, Xie, JHEP 11 (2021) 175

$$\Omega_{\text{GW}}(f)h^2 = \frac{8\pi h^2}{3M_{\text{Pl}}^2 H_0^2} G\mu^2 f \sum_{n=1}^{\infty} C_n(f) P_n$$

$$G\mu \sim \frac{v_\phi^2}{M_{\text{Pl}}^2} \sim 10^{-10} \times \left( \frac{M_{\nu R}/\lambda_R}{10^{14} \text{ GeV}} \right)^2$$




$$C_n = \frac{2n}{f^2} \int_0^\infty \frac{dz}{H(z)(1+z)^6} n_{\text{CS}} \left( \frac{2n}{(1+z)f}, t(z) \right)$$

$$n_{\text{CS}}^r(\ell, t) = \frac{0.18}{t^{3/2}(\ell + \Gamma G\mu t)^{5/2}}, \quad (\ell \leq 0.1 t);$$

$$n_{\text{CS}}^{r,m}(\ell, t) = \frac{0.18 t_{\text{eq}}^{1/2}}{t^2(\ell + \Gamma G\mu t)^{5/2}}, \quad (\ell \leq 0.09 t_{\text{eq}} - \Gamma G\mu t);$$




$$n_{\text{CS}}^m(\ell, t) = \frac{0.27 - 0.45(\ell/t)^{0.31}}{t^2(\ell + \Gamma G\mu t)^2}, \quad (\ell \leq 0.18 t)$$

# FIMP DM and CS GW







Bian, Tang, Zhou, 21

# DW & Discrete symmetry



Kibble mechanism



1002.1555



Amplitude:  $\Omega_{GW}^{dw} h^2 (t_0)_{\text{peak}} \simeq 5.20 \times 10^{-20} \times \tilde{\epsilon}_{\text{gw}} \mathcal{A}^4 \left(\frac{10.75}{g_*}\right)^{1/3} \left(\frac{\sigma_{\text{wall}}}{1\text{TeV}^3}\right)^4 \left(\frac{1\text{MeV}^4}{\Delta V}\right)^2$

Area parameter

peak frequency:  $f^{dw} (t_0)_{\text{peak}} = \frac{a(t_{\text{dec}})}{a(t_0)} H(t_{\text{dec}}) \simeq 3.99 \times 10^{-9} \text{Hz} \mathcal{A}^{-1/2} \left(\frac{1\text{TeV}^3}{\sigma_{\text{wall}}}\right)^{1/2} \left(\frac{\Delta V}{1\text{MeV}^4}\right)^{1/2}$

Slope:  $\Omega_{GW}^{dw} h^2 \propto f^3$  when  $f < f_{\text{peak}}$ , and  $\Omega_{GW}^{dw} h^2 \propto f^{-1}$  when  $f \geq f_{\text{peak}}$

Domain wall decay before they overclose Universe

$$\sigma_{\text{wall}} < 2.93 \times 10^4 \text{TeV}^3 \mathcal{A}^{-1} \left(\frac{0.1\text{sec}}{t_{\text{dec}}}\right)$$

Domain wall decay before the BBN

$$t_{\text{dec}} \approx \mathcal{A} \sigma_{\text{wall}} / (\Delta V)$$


$$\Delta V \gtrsim 6.6 \times 10^{-2} \text{MeV}^4 \mathcal{A} \left(\frac{\sigma_{\text{wall}}}{1\text{TeV}^3}\right)$$

1309.5001



# DW GW & DM

$$\mathcal{L} \supset m_{\psi_1} \bar{\psi}_1 \psi_1 + m_{\psi_2} \bar{\psi}_2 \psi_2 + y_s S \psi_1 \bar{\psi}_2 + y_s S^\dagger \bar{\psi}_1 \psi_2$$

$$V = \mu_S^2 |S|^2 + \lambda_S |S|^4 + \frac{\mu_3}{2} (S^3 + S^{\dagger 3})$$



BM1,  $y_s=7.31 \times 10^{-11}$ ,  $m_\chi=30\text{TeV}$ ,  $m_\psi=35\text{TeV}$



$$\Omega_{\text{GW}}^{\text{dw}} h^2(t_0)_{\text{peak}} \simeq 5.20 \times 10^{-20} \times \tilde{e}_{\text{gw}} \mathcal{A}^4 \left( \frac{10.75}{g_*} \right)^{1/3} \times \left( \frac{\sigma_{\text{wall}}}{1 \text{ TeV}^3} \right)^4 \left( \frac{1 \text{ MeV}^4}{\Delta V} \right)^2$$

$$f^{\text{dw}}(t_0)_{\text{peak}} = \frac{a(t_{\text{dec}})}{a(t_0)} H(t_{\text{dec}}) \simeq 3.99 \times 10^{-9} \text{ Hz} \mathcal{A}^{-1/2} \times \left( \frac{1 \text{ TeV}^3}{\sigma_{\text{wall}}} \right)^{1/2} \left( \frac{\Delta V}{1 \text{ MeV}^4} \right)^{1/2}$$

### ❖ Lattice simulation

- **PT GW simulation with holography models**
- **Topological defects: Magnetic monopoles, cosmic strings, domain walls**

### ❖ Pheno

1. EWSB and GW from FOPT
- **Probing the Higgs Potential shape and EWPT patterns with GW production and Colliders complementarily**
2. BAU and GW from FOPT
- **Sphaleron process, bubble velocity**
3. DM and GW from FOPT
- **DM and high/low-scale PT, DM out-of-equilibrium & FOPT, PBH DM&FOPT**

谢谢！