



中国科学院大学
University of Chinese Academy of Sciences




Highlights from the LHCb experiment

何吉波 (Jibo HE)

26th Mini-workshop on the frontier of LHC


Oct 28-30, 2022

Large Hadron Collider



Proton energy: up to 7 TeV (10^{12} eV)
speed: 0.999999991 c

The LHCb experiment



[JINST 3 (2008) S080005]

Vertex Locator

$$\sigma_{PV,x/y} \sim 10 \mu\text{m}, \sigma_{PV,z} \sim 60 \mu\text{m}$$

Tracking (TT, T1-T3)

$$\Delta p/p: 0.4\% \text{ at } 5 \text{ GeV}/c, \text{ to } 0.6\% \text{ at } 100 \text{ GeV}/c$$

RICHs

$$\varepsilon(K \rightarrow K) \sim 95\%, \text{ mis-ID rate } (\pi \rightarrow K) \sim 5\%$$

Muon system (M1-M5)

$$\varepsilon(\mu \rightarrow \mu) \sim 97\%, \text{ mis-ID rate } (\pi \rightarrow \mu) = 1 - 3\%$$

ECAL



$$\sigma_E/E \sim 10\%/\sqrt{E} \oplus 1\% \text{ (} E \text{ in GeV)}$$

HCAL

$$\sigma_E/E \sim 70\%/\sqrt{E} \oplus 10\% \text{ (} E \text{ in GeV)}$$



Indirect search for New Physics

- Precision measurement of heavy hadron decays
 - Flavour-Changing NC
 - Flavour-Changing CC
- Probe New Physics at high energy scale




Indirect search for NP (cont.)

- Overconstrain the CKM triangle



LHCb at 300 fb^{-1} ,
CMS/ATLAS at 3000 fb^{-1} , Belle II at 50 ab^{-1} .

The LHCb trigger (2018)



- L0, Hardware

- $p_T(\mu_1) \times p_T(\mu_2) > (1.5 \text{ GeV})^2$

- $p_T(\mu) > 1.8 \text{ GeV}$

- $E_T(e) > 2.4 \text{ GeV}$

- $E_T(\gamma) > 3.0 \text{ GeV}$


- $E_T(h) > 3.7 \text{ GeV}$

- High Level Trigger

- Stage1, p_T , IP

- Stage2, full selection









LHCb luminosity prospects



LHC era		HL-LHC era		
Run 1 (2010-12)	Run 2 (2015-18)	Run 3 (2022-24)	Run 4 (2027-30)	Run 5+ (2031+)
3 fb ⁻¹	6 fb ⁻¹	23 fb ⁻¹	46 fb ⁻¹	>300 fb ⁻¹ ??
		Phase-1 Upgrade!!	Phase-1b Upgrade!?	Phase-2 Upgrade??

$$B_{(s)}^0 \rightarrow \mu^+ \mu^-$$

- Suppressed in SM, could be enhanced by New Physics



$B_s^0 \rightarrow \mu^+ \mu^-$ effective lifetime

- B_s^0 mixing \Rightarrow effective τ

$$\tau_{\mu^+\mu^-} = \frac{\tau_{B_s}}{1 - y_s^2} \left[\frac{1 + 2A_{\Delta\Gamma}^{\mu^+\mu^-} y_s + y_s^2}{1 + A_{\Delta\Gamma}^{\mu^+\mu^-} y_s} \right]$$

$$A_{\Delta\Gamma}^{\mu^+\mu^-} \equiv \frac{R_H^{\mu^+\mu^-} - R_L^{\mu^+\mu^-}}{R_H^{\mu^+\mu^-} + R_L^{\mu^+\mu^-}} \quad A_{\Delta\Gamma} = 1 \text{ in SM}$$


$$y_s = \frac{\Delta\Gamma_s}{2\Gamma_s}$$

- Measured by LHCb/CMS, not yet sensitive to $A_{\Delta\Gamma}$


$$\tau_{\mu\mu} = 2.07 \pm 0.29 \pm 0.03 \text{ ps}$$

$$1.83^{+0.23}_{-0.20} {}^{+0.04}_{-0.04} \text{ ps}$$

[CMS-PAS-BPH-21-006]





[De Bruyn *et al.*, PRL 109 (2012) 041801]



$$B^0 \rightarrow K^{*0} \mu^+ \mu^-$$








- Rates and angular distributions sensitive to NP



$$\frac{1}{d(\Gamma + \bar{\Gamma})/dq^2} \frac{d^3(\Gamma + \bar{\Gamma})}{d\bar{\Omega}} = \frac{9}{32\pi} \left[\frac{3}{4}(1 - F_L) \sin^2 \theta_K + F_L \cos^2 \theta_K + \frac{1}{4}(1 - F_L) \sin^2 \theta_K \cos 2\theta_\ell \right. \\ \left. - F_L \cos^2 \theta_K \cos 2\theta_\ell + S_3 \sin^2 \theta_K \sin^2 \theta_\ell \cos 2\phi \right. \\ \left. + S_4 \sin 2\theta_K \sin 2\theta_\ell \cos \phi + S_5 \sin 2\theta_K \sin \theta_\ell \cos \phi \right. \\ \left. + \frac{4}{3} A_{FB} \sin^2 \theta_K \cos \theta_\ell + S_7 \sin 2\theta_K \sin \theta_\ell \sin \phi \right. \\ \left. + S_8 \sin 2\theta_K \sin 2\theta_\ell \sin \phi + S_9 \sin^2 \theta_K \sin^2 \theta_\ell \sin 2\phi \right]$$

Branching fraction of $b \rightarrow s \mu^+ \mu^-$

- Pattern of tensions seen, theoretical uncertainty?




P'_5 with $B^0 \rightarrow K^{*0} \mu^+ \mu^-$


- $P'_5 = \frac{S_5}{\sqrt{F_L(1-F_L)}}$, less form-factor dependent

[S. Descotes-Genon, *et al.*, JHEP 01 (2013) 048]

- Also measured by Belle, ATLAS, CMS





[LHCb, JHEP 02 (2016) 104]
 [Belle, PRL 118 (2017) 111801]
 [ATLAS, JHEP 10 (2018) 047]
 [CMS, PLB 781 (2018) 517]



Lepton flavour universality

- In SM, three lepton families (e, μ, τ) have identical couplings to the gauge bosons



– which means, e.g.,

$$R_K = \frac{\mathcal{B}(B^+ \rightarrow K^+ \mu^+ \mu^-)}{\mathcal{B}(B^+ \rightarrow K^+ e^+ e^-)} \cong 1$$

$\mathcal{O}(10^{-4})$ uncertainty

[C. Bobeth *et al.*, JHEP 12 (2007) 040]

$\mathcal{O}(1\%)$ QED correction

[M. Bordone *et al.*, EJP 76 (2016) 440]


- Lepton flavor universality violation? **New Physics!**

Experimental test of LFU

- Well established in SM, e.g. $W \rightarrow \ell \nu$
 - Some tension at LEP,

addressed by ATLAS/CMS

[arXiv:2007.14040, CMS PAS SMP-18-011]



W Leptonic Branching Ratios

ALEPH	10.78 ± 0.29
DELPHI	10.55 ± 0.34
L3	10.78 ± 0.32
OPAL	10.71 ± 0.27

LEP $W \rightarrow e \nu$ 10.71 ± 0.16


ALEPH	10.87 ± 0.26
DELPHI	10.65 ± 0.27
L3	10.03 ± 0.31
OPAL	10.78 ± 0.26

LEP $W \rightarrow \mu \nu$ 10.63 ± 0.15

ALEPH	11.25 ± 0.38
DELPHI	11.46 ± 0.43
L3	11.89 ± 0.45
OPAL	11.14 ± 0.31



LEP $W \rightarrow \tau \nu$ 11.38 ± 0.21
 $\chi^2/\text{ndf} = 6.3 / 9$

LEP $W \rightarrow \ell \nu$ 10.86 ± 0.09
 $\chi^2/\text{ndf} = 15.4 / 11$






LFU in $b \rightarrow c \ell \nu$ decays

- Deviations from SM seen by Babar/Belle/LHCb








$$R(H_c) = \frac{B(H_b \rightarrow H_c \tau^- \bar{\nu}_\tau)}{B(H_b \rightarrow H_c \mu^- \bar{\nu}_\mu)}$$





LFU in $b \rightarrow s \ell^+ \ell^-$ decays

- Deviations from SM seen by LHCb



$$R_X = \frac{\mathcal{B}(H_b \rightarrow X \mu^+ \mu^-)}{\mathcal{B}(H_b \rightarrow X e^+ e^-)}$$



CKM- γ combination






- Simultaneous determination of CKM- γ & charm mixing parameters

– CKM $\gamma = (63.8^{+3.5}_{-3.7})^\circ$

– Charm mixing $x = (0.398^{+0.050}_{-0.049})\%$,




$y = (0.636^{+0.020}_{-0.019})\%$

[LHCb-Conf-2022-003]



CPV in mixing

- $\phi_S = \phi_M - 2\phi_D$,
small in SM
- $B_S^0 \rightarrow J/\psi h^+ h^-$



ΔA_{CP} in charm

$$A_{CP}(f) = \frac{\Gamma(M \rightarrow f) - \Gamma(\bar{M} \rightarrow \bar{f})}{\Gamma(M \rightarrow f) + \Gamma(\bar{M} \rightarrow \bar{f})}$$


$$\Delta A_{CP} \equiv A_{CP}(K^- K^+) - A_{CP}(\pi^- \pi^+)$$

$$\Delta A_{CP}^{\pi\text{-tagged}} = [-18.2 \pm 3.2 (\text{stat.}) \pm 0.9 (\text{syst.})] \times 10^{-4},$$


$$\Delta A_{CP}^{\mu\text{-tagged}} = [-9 \pm 8 (\text{stat.}) \pm 5 (\text{syst.})] \times 10^{-4}.$$

Combined one:

$$\Delta A_{CP} = (-15.4 \pm 2.9) \times 10^{-4}$$





[PRL 122 (2019) 211803]



W mass

- Anti-correlation of PDF at GPD/LHCb
- More measurements required at LHC

[JHEP 01 (2022) 036]



Intrinsic charm?

- Bound to valence quarks, longer time scales
- Z associated with charm













Image: D. Craik




Spectroscopy




- Total fit
- $\bar{D}_2^*(2460) D_s^+$
- $\bar{D}_1^*(2600) D_s^+$
- $\bar{D}_3^*(2750) D_s^+$
- $\bar{D}_1^*(2760) D_s^+$
- $\bar{D}(3000) D_s^+$
- $D^*(2010) - D_s^+$
- $T_{c\bar{s}0}^*(2900) \bar{D}$
- $D\pi$ S-wave D_s^+
- + Data
- Background

$X(6900)$ in Di- J/ψ system ($c\bar{c}c\bar{c}$)




[Sci. Bull. 65 (2020) 032]



No interference, using Breit-Wigner
 $m = 6905 \pm 11 \pm 7$ MeV
 $\Gamma = 80 \pm 19 \pm 33$ MeV







NRSPS interference w/ broad structure
 $m = 6886 \pm 11 \pm 11$ MeV
 $\Gamma = 168 \pm 33 \pm 69$ MeV





Strange pentaquark

($c\bar{c}uds$)



[arXiv:2210.10346]




$$m = 4338.2 \pm 0.7 \pm 0.4 \text{ MeV}$$

$$\Gamma = 7.0 \pm 1.2 \pm 1.3 \text{ MeV}$$

Effective Field Theory of $b \rightarrow sl\ell$

- Integrate out short-distance (high energy) interactions
- Operator production expansion




$$\mathcal{H}_{\text{eff}} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \frac{e^2}{16\pi^2} \sum_i (C_i O_i + C_i' O_i') + h.c.$$

- Wilson coefficients $C_i^{(')}$ encode short-distance physics
- Operators $O_i^{(')}$ describe low-energy QCD (using form factors), which have large theory uncertainties


Global fit

- Different experimental inputs, form factors, assumptions about non-local matrix elements, statistical frameworks

B. Capdevila, M. Fedele, S. Neshatpour, P. Stangl @ LHCb implications 2021 [\[slides\]](#)



global fit









fit to LFU observables + $B_s \rightarrow \mu\mu$

Implications of Flavour anomalies?

G. Isidori @ NJNU

A. Greljo @ LHCb implication 2021 [\[slides\]](#)

Azatov, Garosi, AG, Marzocca, Salko, Trifinopoulos; w.i.p



[CMS, PLB 792 (2019) 345]

Prospects

- LHCb upgrades (2025: 23 fb⁻¹, Upgrade-II: 300 fb⁻¹)

Observable	Current LHCb	LHCb 2025	Belle II	Upgrade II	ATLAS & CMS
EW Penguins					
$R_K (1 < q^2 < 6 \text{ GeV}^2 c^4)$	0.1 [274]	0.025	0.036	0.007	–
$R_{K^*} (1 < q^2 < 6 \text{ GeV}^2 c^4)$	0.1 [275]	0.031	0.032	0.008	–
R_ϕ, R_{pK}, R_π	–	0.08, 0.06, 0.18	–	0.02, 0.02, 0.05	–
CKM tests					
γ , with $B_s^0 \rightarrow D_s^+ K^-$	$(^{+17}_{-22})^\circ$ [136]	4°	–	1°	–
γ , all modes	$(^{+5.0}_{-5.8})^\circ$ [167]	1.5°	1.5°	0.35°	–
$\sin 2\beta$, with $B^0 \rightarrow J/\psi K_s^0$	0.04 [606]	0.011	0.005	0.003	–
ϕ_s , with $B_s^0 \rightarrow J/\psi \phi$	49 mrad [44]	14 mrad	–	4 mrad	22 mrad [607]
ϕ_s , with $B_s^0 \rightarrow D_s^+ D_s^-$	170 mrad [49]	35 mrad	–	9 mrad	–
$\phi_s^{s\bar{s}}$, with $B_s^0 \rightarrow \phi \phi$	154 mrad [94]	39 mrad	–	11 mrad	Under study [608]
a_{sl}^s	33×10^{-4} [211]	10×10^{-4}	–	3×10^{-4}	–
$ V_{ub} / V_{cb} $	6% [201]	3%	1%	1%	–
$B_s^0, B^0 \rightarrow \mu^+ \mu^-$					
$\mathcal{B}(B^0 \rightarrow \mu^+ \mu^-)/\mathcal{B}(B_s^0 \rightarrow \mu^+ \mu^-)$	90% [264]	34%	–	10%	21% [609]
$\tau_{B_s^0 \rightarrow \mu^+ \mu^-}$	22% [264]	8%	–	2%	–
$S_{\mu\mu}$	–	–	–	0.2	–
$b \rightarrow c \ell^- \bar{\nu}_\ell$ LUV studies					
$R(D^*)$	0.026 [215, 217]	0.0072	0.005	0.002	–
$R(J/\psi)$	0.24 [220]	0.071	–	0.02	–
Charm					
$\Delta A_{CP}(KK - \pi\pi)$	8.5×10^{-4} [610]	1.7×10^{-4}	5.4×10^{-4}	3.0×10^{-5}	–
$A_\Gamma (\approx x \sin \phi)$	2.8×10^{-4} [240]	4.3×10^{-5}	3.5×10^{-4}	1.0×10^{-5}	–
$x \sin \phi$ from $D^0 \rightarrow K^+ \pi^-$	13×10^{-4} [228]	3.2×10^{-4}	4.6×10^{-4}	8.0×10^{-5}	–
$x \sin \phi$ from multibody decays	–	$(K3\pi) 4.0 \times 10^{-5}$	$(K_s^0 \pi\pi) 1.2 \times 10^{-4}$	$(K3\pi) 8.0 \times 10^{-6}$	–

[CERN-LHCC-2018-027]


Summary

- Highlights from LHCb
 - Flavour anomalies, $b \rightarrow s\mu^+\mu^-$ BR, P'_5 , $\mathcal{R}_{K^{(*)0}}$, \mathcal{R}_{D^*} , to be confirmed or refuted with more data
 - CP violation, CKM triangle, ϕ_s, γ
 - W mass, intrinsic charm
 - Spectroscopy, $X(6900)$, $P_{cS}(4338)$, ...
- Your suggestions are always appreciated!
 - New observables?

中国科学：物理学 力学 天文学

SCIENCE CHINA Physics, Mechanics & Astronomy

SCI
5.203
IMPACT
FACTOR | Q1



- **Editor's Focus: aim at PRL quality, fast channel**
- **Full-text HTML and timely publication (online immediately)**
- **Two-month free access at Springer**
- **Highlighted at EurekaAlert and other public media**



- **Since 1950, in Chinese**
- **100 Outstanding Academic Journals of China**
- **Indexed in Scopus, ESCI, etc.**
- **Special topic is encouraged, published over 70 special topics**



**Scan the QR code
Get the news**

Supervised by



Sponsored by



Published by

