


Probing the isobaric Ru and Zr nuclear structure with the diffractive photoproduction of ρ^0 mesons

Jie Zhao

Aug. 4, 2023

Charge radius

R. Hofstadter, Rev. Mod. Phys. 28 214-254 (1956)




$$\frac{d\sigma}{d\Omega} = \left(\frac{d\sigma}{d\Omega} \right)_{\text{point}} |F(\vec{q})|^2 \quad F(\vec{q}) = \int d\vec{r} e^{i\vec{q}\cdot\vec{r}} \rho(\vec{r})$$

$$F(q^2) = \int d\vec{r} \rho(r) e^{i\vec{q}\cdot\vec{r}} \xrightarrow{qR \ll 1} 1 - \frac{1}{6} q^2 \int dr d\Omega r^4 \rho(r) + \dots$$





$$\dots = 1 - \frac{1}{6} q^2 \langle r^2 \rangle + \dots$$

G.F. Chew, et.al, Phys. Rev. 106, 1345 (1957); R.A. Schrack, et.al, Phys. Rev. 127, 1772 (1962);



➤ Electron scattering measures the form factor, **charge radius**

Mass radius



$$A(\gamma p \rightarrow \pi^0 p) = A(\gamma n \rightarrow \pi^0 n)$$

G.F. Chew, et.al, Phys. Rev. 106, 1345 (1957); R.A. Schrack, et.al, Phys. Rev. 127, 1772 (1962);
C.M. Tarbert, et.al, Phys. Rev. Lett. 112, 242502 (2014)

- Electron scattering measures the form factor, charge **radius**
- Photoproduction of π^0 meson: $\Delta(1232)$, **the mass radius** (1960s)

Strong-Interaction Nuclear Radii

$$\gamma + (Z, A) \rightarrow \rho + (Z, A)$$








FIG. 1. Results on $R(A)$ and the best fit with $R = r_0 A^{1/3}$.

G.F. Chew, et.al, Phys. Rev. 106, 1345 (1957); R.A. Schrack, et.al, Phys. Rev. 127, 1772 (1962);
 C.M. Tarbert, et.al, Phys. Rev. Lett. 112, 242502 (2014)
 F. Bulos, et.al, Phys. Rev. Lett. 22, 490 (1969); L.J. Lanzerotti, et.al, Phys. Rev. 166, 1365 (1968)
 H. Alvensleben et.al, Phys. Rev. Lett. 24, 786 and 792 (1970)

- Electron scattering measures the form factor, charge **radius**
- Photoproduction of π^0 meson: $\Delta(1232)$, **the mass radius** (1960s)
- Photoproduction of ρ^0 meson:
 “**Determination of Strong-Interaction Nuclear Radii**” (1970s)



STAR, PRC 96, 054904 (2017)
 STAR, PRL 89, 272302 (2002)
 Spencer, et.al, PRC 60, 014903, (1999)

F. Bulos, et.al, Phys. Rev. Lett. 22, 490 (1969)
 H. Alvensleben et.al, Phys. Rev. Lett. 24, 786 and 792 (1970)
 H. Mäntysaari, F. Salazar, B. Schenke, arXiv:2207.03712

- Diffractive pattern (minima) of the coherent ρ^0 production are sensitive to the nuclear size.
- “The slopes of the diffraction patterns measure directly the nuclear density distribution. For example, at $t \rightarrow 0$, the diffraction pattern behaves as e^{at} where a is a measure of the nuclear size.”
- Can used to study the nuclear structure of the isobar Ru and Zr


STAR, Phys. Rev. C 105 (2022), 014901
 T. Prithwish (for STAR), QM2022

STAR, Sci. Adv. 9 (2023) 1

Neutron skin & nuclear deformation of isobars


Talk by Haojie Xu (Wed T01-II)
 Posters by Chunjian Zhang (Wed T14_2),
 Jiangyong Jia (Wed T01)

Precision ratios of flow harmonics (v_2, v_3), asymmetric cumulants ($ac\{3\}$), $\langle p_T \rangle$, moments of $\langle p_T \rangle$ fluctuation, multiplicity distribution $P(N_{ch})$ and net-charge multiplicity (ΔQ) measured in isobars



Pioneering new ways to constrain neutron skin & nuclear deformation with heavy ion collisions

STAR overview, P. Tribedy, QM 2022, Krakow, Poland





Tomography of ultra-relativistic nuclei with polarized photon-gluon collisions.

➤ The γ -A interaction may help to understand the structure of the isobar Ru and Zr nuclei ?

Spencer, et.al, PRC 60, 014903, (1999); STAR, PRL 89, 272302 (2002), PRC 96, 054904 (2017)


C.A. Bertulani, S.R. Klein, J. Nystrand
Ann. Rev. Nucl. Part. Sci. 55 (2005) 271

STAR, PRL 89, 272302 (2002)



Soft-Pomeron (gluons) exchange

- Ultra-peripheral heavy-ion collisions (UPC) ($b > 2R_0$)
- Electromagnetic field as quasi-real photons (EPA)
- ρ^0 meson production through the soft-pomeron (gluons) exchange, sensitive to the **Strong-Interaction Nuclear Radius**



- Time Projection Chamber: tracking and particle identification within $|\eta| < 1$
- Time Of Flight: multiplicity trigger, particle identification and pile-up track removal
- Barrel ElectroMagnetic Calorimeter: topology trigger and pile-up track removal
- Beam-Beam Counter: scintillator counters within $2.1 < |\eta| < 5.2$, forward veto
- Zero Degree Calorimeters: detection of very forward neutrons, $|\eta| > 6.6$


System	Year	Trigger (ID)	Vz	Vr
Ru+Ru	2018	UPC-Jpsi	<100cm	<2cm
Zr+Zr	2018	UPC-Jpsi	<100cm	<2cm

Track cuts: (primary track)

$n_{\text{HitFit}} > 15$, $|\eta| < 1$, $dca < 3\text{cm}$

$p_T > 0.2 \ \&\& \ < 4 \text{ GeV}/c$



TOF matched > 0



plot from Jaroslav Adam

- Events are selected with **number of primary tracks = 2**
- Track pair **not** matched to the BEMC topo. clusters with ($E_0 > 0.5$)

PID: remove the e^+e^- contamination





STAR, PRL 127, 052302 (2021)

ALICE, JHEP. 2015, 95 (2015)

➤ $\sigma_{\pi,A}^2 + \sigma_{\pi,B}^2 < 16$,
remove the e^+e^- contamination from $\gamma+\gamma$ interaction

STAR Diffractive photoproduction of ρ^0 in isobar





➤ Systematic uncertainty sources:

dca: 1.0, 2.0 (3.0) cm; nHitsFit: 20 (15); $|V_z|$: 50 (100) cm

➤ Total systemic uncertainty : $\text{RMS}(\sigma(\text{dca})) \otimes \sigma(\text{nHit}) \otimes \sigma(V_z)$

➤ Diffraction pattern (minima) of the coherent ρ^0 production

Ru and Zr nuclear structure





$$A^* e^{-b^*t}, \quad (t \simeq -p_T^2)$$

- Indication of larger Zr size than Ru from the γ -A interaction.
The slope of the dN/dt ratio is $11.0 \pm 2.9 \pm 0.3$ ($\sim 3\sigma$ sigma effect)
- Interference and deformation effects need to be considered

TABLE II. The Woods-Saxon parameters used in the Glauber simulations for the centrality determination.

Nucleus	Case-1 [83]			Case-2 [83]			Case-3 [113]		
	R (fm)	a (fm)	β_2	R (fm)	a (fm)	β_2	R (fm)	a (fm)	β_2
$^{96}_{44}\text{Ru}$	5.085	0.46	0.158	5.085	0.46	0.053	5.067	0.500	0
$^{96}_{40}\text{Zr}$	5.02	0.46	0.08	5.02	0.46	0.217	4.965	0.556	0



- STARLight simulation with case 1 shows negative slope of the ratio, where Ru size is larger than Zr.
- Case 3: simulation shows positive, which indicates larger Zr size.
- Case 3 agrees better with data, but still large difference (slope ~ 1.7)

- Diffractive photoproduction of ρ^0 mesons in isobar
- Indication of larger Zr size than Ru from the γ -A, the slope of the dN/dt ratio is $11.0 \pm 2.9 \pm 0.3$ ($\sim 3\sigma$ effect)
- Comparison with simulation also indicates larger Zr size
- Interference and deformation effects need to be considered using model calculations