


介质中的形状因子与自旋极化现象




Shu Lin

Sun Yat-Sen University

原子核结构与相对论重离子碰撞，大连， Aug 1 – 5, 2023



nucleus structures discussed so far in the workshop




this talk: nucleus (parton) spin structures

Outline


- ◆ Spin polarizations in heavy ion collisions
- ◆ Success and limitation of quantum kinetic theory
- ◆ Form factors description of spin couplings
- ◆ Electromagnetic form factors in vacuum and in medium
- ◆ Gravitational form factors in vacuum and in medium
- ◆ Summary and outlook

global spin polarization in heavy ion collisions




$$L_{ini} \sim 10^5 \hbar \rightarrow S_{final}$$

Liang, Wang, PRL 2005, PLB 2005






STAR collaboration, Nature 2017 $e^{-\beta(H_0 - \mathbf{S} \cdot \boldsymbol{\omega})}$

local spin polarization in heavy ion collisions



STAR collaboration, PRL 2019



Fu, Liu, Pang, Song, Yin, PRL 2021
 Becattini, et al, PRL 2021
 Yi, Pu, Yang, PRC 2021

vorticity+shear

$$\mathcal{P}^i \sim \omega^i \quad \mathcal{P}^i \sim \epsilon^{ijk} \hat{p}_j \hat{p}_l \sigma_{kl}$$

talk by S. Pu

Quantum kinetic theory for
 collisional contribution etc

SL, Z.y. Wang,
 JHEP 2022




Quantum kinetic theory (QKT): pro and cons

QKT=Boltzmann+spin

Hidaka, Pu, Wang, Yang,
PPNP 2022

Pro: systematic treatment of spin transport at partonic level

Cons: assumes point particle




elementary particle not point-like due to quantum fluctuation

Spin polarization in heavy ion collisions

for $S = \frac{1}{2}$ particle (consider high temperature limit, quark massless)

$$S_i \sim B_i$$


$$S_i \sim \epsilon^{ijk} \hat{p}_j E_k$$



particle in external EM fields

$$S_i \sim \omega_i$$

$$S_i \sim \epsilon^{ijk} \hat{p}_j \hat{p}_l \sigma_{kl}$$



particle in off-equilibrium state:
hydro gradient (mimicked by
metric fields)

Spin polarization and correlation functions

Wigner function

$$S_{\alpha\beta}^{\langle}(X = \frac{x+y}{2}, P) = \int d^4(x-y) e^{iP \cdot (x-y)/\hbar} (-\langle \bar{\psi}_{\beta}(y) \psi_{\alpha}(x) \rangle)$$


- Spin polarization in external electromagnetic fields

$$\langle S^{\langle}(X, P) \rangle_{\text{eq}, A_{\mu}}$$


- Spin polarization in off-equilibrium state: hydro gradient

$$\langle S^{\langle}(X, P) \rangle_{\text{off-eq}} = \langle S^{\langle}(X, P) \rangle_{\text{eq}, h_{\mu\nu}}$$

$$A_{\mu}, h_{\mu\nu} \text{ slow-varying} \quad \partial_X \ll P$$



Electromagnetic form factors





$$Q = P_2 - P_1$$

$$P = \frac{1}{2}(P_1 + P_2)$$

$$\begin{aligned} \langle P_2 | J^\mu(Q) | P_1 \rangle &= \bar{u}(P_2) \left[\gamma^\mu F_1(Q^2) + \frac{i\sigma^{\mu\nu} Q_\nu}{2m} F_2(Q^2) \right] u(P_1) \\ &= \bar{u}(P_2) \left[\frac{P^\mu}{m} G_E(Q^2) + \frac{i\epsilon^{\mu\nu\rho\sigma} Q_\nu P_\rho \gamma_\sigma \gamma^5}{2m^2} G_M(Q^2) \right] u(P_1) \end{aligned}$$


Form factors parameterize interaction based on symmetry

- accessible experimentally
- charge distribution, magnetic moment




PLB 2018,
Ye et al

Gravitational form factors



$$Q = P_2 - P_1$$

$$P = \frac{1}{2}(P_1 + P_2)$$



Nature 2021, Meziani et al

$$\langle P_2 | T^{\mu\nu}(Q) | P_1 \rangle = \bar{u}(P_2) \left[\frac{P^\mu P^\nu}{m} A(Q^2) + \frac{i P^{\{\mu} \sigma^{\nu\} \rho} Q_\rho}{2m} J(Q^2) + \frac{Q^\mu Q^\nu - \eta^{\mu\nu} Q^2}{4m} D(Q^2) \right] u(P_1)$$

Form factors parameterize interaction based on symmetry

- accessible experimentally
- mass distribution, **gravitomagnetic moment**, internal structures

Spin polarization from QKT(CKT)

right-handed fermion
in EM fields

$$S^<(X = \frac{x+y}{2}, P) = \int d^4(x-y) e^{iP \cdot (x-y)/\hbar} U(y, x) \left(-\langle \psi^\dagger(y) \psi(x) \rangle \right)$$

gauge link $U(y, x) = \mathcal{P} \exp[-i \int_x^y dz^\mu A_\mu(z)]$

$$S^< \equiv \bar{\sigma}_\mu S^<\mu$$

$$S^<^0 = -2\pi [\delta(P^2) p_0 f(p_0) + \mathbf{p} \cdot \mathbf{B} \delta'(P^2) f(p_0)]$$

absorb e in E&B fields

$$S^<^i = -2\pi [\delta(P^2) p_i f(p_0) + (p_0 B_i - \epsilon^{ijk} P_j E_k) \delta'(P^2) f(p_0)]$$

$O(\partial^0)$

$\mathbf{E}, \mathbf{B} \sim O(\partial)$

spin-magnetic spin Hall effect
coupling

Hidaka, Pu, Yang, PRD 2018
Gao, Liang, Wang, PRD 2019

Spin polarization from field theory

$$S^{<\mu} = -(S_{ra}^\mu - S_{ar}^\mu) f(P_0)$$

$$S_{ra} = \overline{r} \text{---} a + \overline{r} \text{---} a \begin{array}{c} \text{---} r \\ \text{---} r \\ \text{---} r \\ \otimes \end{array} + \overline{r} \text{---} a \begin{array}{c} \text{---} r \\ \text{---} r \\ \text{---} r \\ \otimes \end{array} \text{---} a \begin{array}{c} \text{---} r \\ \text{---} r \\ \text{---} r \\ \otimes \end{array} \quad q \sim \partial_X$$

resummation to all order in A , and up to $O(q)$ reproduces the CKT results


Two lessons:

- point particle, EM interaction Lorentz invariant
- Electric field can't do work to fermion (implicit in CKT)

$$n^\mu \text{ frame vector} \quad E^\mu = F^{\mu\nu} n_\nu \quad Q \cdot n = 0 \quad \text{static}$$

$$\text{no-work condition} \quad E \cdot P = 0 \quad \longrightarrow \quad P \cdot Q = 0 \quad \text{orthogonal}$$

EM form factors for point particle



$$P^2 = 0 \quad \longrightarrow \quad P_{1,2}^2 \sim O(Q^2)$$

$$P \cdot Q = 0$$

interpretation: scattering of fermions on EM fields

$P_i \cdot \bar{\sigma} = u(P_i)u(P_i)^\dagger$ from massless propagators

$$u(P_2)u^\dagger(P_2)i\sigma^\mu u(P_1)u^\dagger(P_1)A_\mu$$

$$u^\dagger(P_2)i \left(n^\mu + \hat{p}^\mu + \frac{i\epsilon^{\mu\nu\rho\sigma} n_\nu P_\rho Q_\sigma}{2(P \cdot n)} \right) u(P_1) \quad n^\mu \text{ frame vector}$$

no-work condition

$Q \cdot n = 0$ Ward identity satisfied by each structure,

$P \cdot Q = 0$ three form factors (FF) degenerate




What to expect for FF in **medium**?

Effect of quantum fluctuation

- both vertex and fermion states corrected by interaction
- Lift degeneracy of FF


Effect of **medium**

- enhanced fluctuation: large phase space for particles in loop
- breaking of Lorentz invariance, more structures possible
- dissipation effect introduces non-hermiticity, complex form factors



EM form factors in medium

$n^\mu \rightarrow u^\mu$ medium frame vector



$$\sigma^\mu \rightarrow \Gamma^\mu = F_0 u^\mu + F_1 \hat{p}^\mu + F_2 \frac{i\epsilon^{\mu\nu\rho\sigma} u_\nu P_\rho Q_\sigma}{2(P \cdot u)^2}$$

$$S^{<0} = F_2 \left(\vec{p} \cdot \vec{B} \right) 2\pi\delta'(P^2) f(p_0)$$

$$S^{<i} = \left[F_0 \epsilon^{ijk} E_j p_k + F_1 \left(p_0 B^i - (\vec{B} \cdot \vec{p}) \hat{p}^i \right) + F_2 \left(\vec{B} \cdot \vec{p} \right) \hat{p}^i \right] 2\pi\delta'(P^2) f(p_0)$$

spin Hall effect

spin-perpendicular
magnetic coupling

spin-parallel
magnetic coupling

medium interaction can lift the
degeneracy of form factors

Transformation under time-reversal


$$\Gamma^\mu = F_0 u^\mu + F_1 \hat{p}^\mu + F_2 \frac{i\epsilon^{\mu\nu\rho\sigma} u_\nu P_\rho Q_\sigma}{2(P \cdot u)^2}$$

$$\Gamma^0 \quad \text{T-even} \qquad \Gamma^i \quad \text{T-odd}$$

→ All form factors T-even

$$\begin{array}{l} Q \cdot n = 0 \\ P \cdot Q = 0 \end{array} \quad F_i = F_i(p^2, q^2) \quad \rightarrow \quad \text{All form factors real}$$

Example of medium correction to EMFF: vertex




$$-2im_f^2 A_\nu \sigma_\lambda \left(\hat{l}^\lambda \hat{l}^\nu \frac{1}{p^2} \ln \frac{2p}{q} + \hat{p}^\lambda \hat{p}^\nu \frac{1}{p^2} - \hat{p}^\lambda P^\nu \frac{1}{p^3} \ln \frac{2p}{q} \right)$$

$$m_f^2 = \frac{1}{8} g^2 T^2 C_F \quad \hat{l}^i = \frac{1}{pq} \epsilon^{ijk} q_j p_k$$

simplifications

- medium contribution only (HTL)
- leading contributions as $q \rightarrow 0$

Example of medium correction to EMFF: self-energy




$$S^{ra}(P) = \frac{i}{2} \Delta_+(P) (\gamma^0 - \gamma \cdot \hat{p}) + \frac{i}{2} \Delta_-(P) (\gamma^0 + \gamma \cdot \hat{p})$$

chiral symmetry remains

→ $\delta\Gamma^\mu = \delta Z_+ \sigma^\mu$

$$p \gg m_f: \quad \delta\Gamma_{self-energy}^\nu A_\nu = 2 \frac{m_f^2}{2p^2} \left(1 - \ln \frac{2p^2}{m_f^2} \right) \sigma^\nu A_\nu.$$



Le Bellac, thermal field theory

Example of medium correction to EMFF: sum

$$\delta F_0 = \frac{2m_f^2}{p^2} X + \frac{m_f^2}{p^2} \left(1 - \ln \frac{2p^2}{m_f^2} \right), \quad \text{spin Hall effect}$$

$$\delta F_1 = \frac{2m_f^2}{p^2} (X - 1) + \frac{m_f^2}{p^2} \left(1 - \ln \frac{2p^2}{m_f^2} \right), \quad \text{spin-perpendicular magnetic coupling}$$

$$\delta F_2 = \frac{2m_f^2}{p^2} X + \frac{m_f^2}{p^2} \left(1 - \ln \frac{2p^2}{m_f^2} \right), \quad \text{spin-parallel magnetic coupling}$$

$$X = \frac{1}{6} \left(2 \ln \left(\frac{pT}{m_f^2} \right) + \ln \left(\frac{2pT}{m_g^2} \right) - 36 \ln(A) + \ln(16\pi^3) + 3 \right)$$

- all form factors real
- partial lift of the degeneracy $\delta F_1 \neq \delta F_2 = \delta F_0$

Gravitational FF in vacuum

FF for massless case $Q \rightarrow 0$, ignore D-term

$$\langle P_2 | T^{\mu\nu}(Q) | P_1 \rangle = \bar{u}(P_2) \left[A(Q^2) \frac{P^\mu P^\nu}{P \cdot n} \pm B(Q^2) \frac{-i P^{\{\mu} \epsilon^{\nu\} \lambda \sigma \rho} \gamma_\lambda n_\sigma Q_\rho}{P \cdot n} \right] u(P_1)$$

compared to massive case

$$\langle P_2 | T^{\mu\nu}(Q) | P_1 \rangle = \bar{u}(P_2) \left[\frac{P^\mu P^\nu}{m} A(Q^2) + \frac{i P^{\{\mu} \sigma^{\nu\} \rho} Q_\rho}{2m} J(Q^2) + \frac{Q^\mu Q^\nu - \eta^{\mu\nu} Q^2}{4m} D(Q^2) \right] u(P_1)$$

$$T^{\mu\nu} = \frac{i}{2} \bar{\psi} \left(\gamma^{\{\mu} \partial^{\nu\}} - \gamma^{\{\mu} \overleftarrow{\partial}^{\nu\}} \right) \psi. \quad \text{point particle}$$

→ $A = 1 \quad B = -\frac{1}{2}$

metric perturbation $h_{0i}(t, x) = v_i(t, x)$.

$$i\mathcal{M} \sim \bar{u}(P) \sigma_k u(P) i\epsilon^{ijk} q_j v_i \sim \vec{S} \cdot \vec{\omega}$$

spin-vorticity coupling

Gravitational FF in medium

Einstein equivalence principle $B(Q^2 = 0) = -\frac{1}{2}$

spin-vorticity coupling dictated for any $S = \frac{1}{2}$ particle

medium breaks Lorentz invariance,
violating equivalence principle!

Donoghue et al 1984, 1985

Buzzegoli, Kharzeev, PRD 2021

SL, Tian, 2302.12450

$$\Gamma^{\mu\nu} = \gamma \cdot \hat{p} \left(F_0 u^\mu u^\nu + F_1 u^{\{\mu} \hat{p}^{\nu\}} + F_2 \hat{p}^\mu \hat{p}^\nu \right) + \gamma \cdot \hat{l} \left(F_3 \hat{p}^{\{\mu} \hat{l}^{\nu\}} + F_4 u^{\{\mu} \hat{l}^{\nu\}} \right)$$

$$\hat{l}_i = \epsilon^{ijk} \hat{q}_j \hat{p}_k$$

no-work
condition

$$q_0 = 0 \quad P \cdot Q = 0$$

five structures, each satisfies energy-momentum conservation

Gravitational FF in medium: example

vertex correction

$$\delta\Gamma^{\mu\nu} = m_f^2 \left[-\gamma \cdot \hat{p} P^\mu P^\nu \frac{\ln \frac{2p}{q}}{p^3} - \gamma \cdot \hat{l} P^{\{\mu} \hat{l}^{\nu\}} \frac{\ln \frac{2p}{q}}{p^2} + \gamma \cdot \hat{p} \left(2u^\mu u^\nu + u^{\{\mu} \hat{p}^{\nu\}} + \hat{p}^\mu \hat{p}^\nu \right) \frac{1}{p} + 2\gamma \cdot \hat{l} \hat{l}^{\{\mu} \hat{p}^{\nu\}} \right]$$

self-energy

$$\delta\Gamma^{\mu\nu} = \delta Z_+ \gamma^{\{\mu} P^{\nu\}} \quad \delta Z_+ = \frac{m_f^2}{2p^2} \left(1 - \ln \frac{2p^2}{m_f^2} \right)$$

Application: spin-vorticity coupling receives multiplicative renormalization

$$\text{e.g. } p = 500\text{MeV} \quad T = 150\text{MeV} \quad \alpha_s = 0.3$$

7% suppression of spin-vorticity coupling

Summary

- Wigner function from CKT reproduced using field theory, allow for form factors description of spin coupling
- In-medium electromagnetic FF lift degeneracy of spin magnetic coupling and spin Hall effect
- In-medium gravitational FF leads to suppression of spin-vorticity coupling

Outlook

- Examples of composite particle: Λ etc
- Dissipation effect: complex FF
- Applications to spin polarization in heavy ion collisions

Thank you!

IR limit and screening

$$-2im_f^2 A_\nu \sigma_\lambda \left(\hat{l}^\lambda \hat{l}^\nu \frac{1}{p^2} \ln \frac{2p}{q} + \hat{p}^\lambda \hat{p}^\nu \frac{1}{p^2} - \hat{p}^\lambda P^\nu \frac{1}{p^3} \ln \frac{2p}{q} \right)$$

potential IR divergent as $q \rightarrow 0$, divergence cutoff by screening effect

$$\delta\Gamma_{vertex}^\nu A_\nu = 2m_f^2 A_\nu \sigma_\lambda \left[\frac{1}{6p^2} \left(2 \ln \left(\frac{pT}{m_f^2} \right) + \ln \left(\frac{2pT}{m_g^2} \right) - 36 \ln(A) + \ln(16\pi^3) + 3 \right) \right. \\ \left. \times \left(\hat{l}^\lambda \hat{l}^\nu - \hat{p}^\lambda P^\nu \frac{1}{p} \right) + \hat{p}^\lambda \hat{p}^\nu \frac{1}{p^2} \right],$$

$$m_g^2 = \frac{1}{3} g^2 T^2 (C_A + \frac{1}{2} N_f) \quad A \simeq 1.282$$

Spin Hall effect

$$\dot{\mathbf{x}} = \hat{\mathbf{p}} + \dot{\mathbf{p}} \times \mathbf{b};$$

$$\dot{\mathbf{p}} = \mathbf{E} + \dot{\mathbf{x}} \times \mathbf{B}.$$