



Contribution ID: 111

Type: **Parallel or Poster session**

Searching for dark neutrinos through exotic Higgs decays at the ILC

Friday, 1 December 2023 11:45 (15 minutes)

In this study we investigate the feasibility of detecting heavy dark neutrinos (N_d) through exotic Higgs decays at the proposed International Linear Collider (ILC), specifically in the channel of $e^+e^- \rightarrow qq H$ with $H \rightarrow \nu N_d \rightarrow \nu l W \rightarrow \nu l qq$. Analyses based on full detector simulations of the ILD are performed at the center-of-mass energy of 250 GeV for two different beam polarization schemes with a total integrated luminosity of 2 ab^{-1} . A range of dark neutrino masses between the Z boson and Higgs boson masses are studied. The 2σ significance reach for the joint branching ratio of $BR(H \rightarrow \nu N_d) \cdot BR(N_d \rightarrow l W)$ is about 0.1%, nearly independent of the dark neutrino masses, while the 5σ discovery is possible at a branching ratio of 0.3%. Interpreting these results in terms of constraints on the mixing parameters $|\varepsilon_{id}|^2$ between SM neutrinos and the dark neutrino, it is expected to have at least a factor of 10 improvement from current constraints.

You are

non-PhD student

Primary authors: TIAN, Junping (The University of Tokyo); THOR, Simon (KTH Royal Institute of Technology); Prof. ISHINO, Masaya (The University of Tokyo)

Presenter: THOR, Simon (KTH Royal Institute of Technology)

Session Classification: Parallel: Future

Track Classification: Future colliders and experiments (including projections)