


# Double-Aligned 2HDM at the LHC

Michihisa Takeuchi (Sun Yat-sen Univ. (Zhuhai)[中山大学, 珠海])

Phys.Rev.D 105 (2022) 11, 115001

# CP violation beyond the SM required

- Baryon Asymmetry of the Universe by EWBG : too small CPV in the SM  
→ **CPV source of BSM required**
- Consider the possibility: new CPV phases exist in an extended Higgs sector



# Aligned CPV 2HDM and EDM

## Higgs potential (without Z2 sym.)

$$\begin{aligned}
 V = & -\mu_1^2 |\Phi_1|^2 - \mu_2^2 |\Phi_2|^2 - \left\{ \mu_3^2 (\Phi_1^\dagger \Phi_2) + h.c. \right\} \\
 & + \frac{1}{2} \lambda_1 |\Phi_1|^4 + \frac{1}{2} \lambda_2 |\Phi_2|^4 + \lambda_3 |\Phi_1|^2 |\Phi_2|^2 + \lambda_4 |\Phi_2^\dagger \Phi_1|^2 \\
 & + \left\{ \left[ \frac{1}{2} \lambda_5 (\Phi_1^\dagger \Phi_2) + \lambda_6 |\Phi_1|^2 + \lambda_7 |\Phi_2|^2 \right] (\Phi_1^\dagger \Phi_2) + h.c. \right\}
 \end{aligned}$$

(Higgs basis)  
[Davidson, Haber, PRD72, 035004 (2005)]

## Yukawa couplings

$$\begin{aligned}
 \mathcal{L}_{\text{yukawa}} = & -\bar{Q}_L \frac{\sqrt{2} M_u}{v} (\tilde{\Phi}_1 + \zeta_u \tilde{\Phi}_2) u_R \\
 & -\bar{Q}_L \frac{\sqrt{2} M_d}{v} (\Phi_1 + \zeta_d \Phi_2) d_R \\
 & -\bar{L}_L \frac{\sqrt{2} M_e}{v} (\Phi_1 + \zeta_e \Phi_2) e_R \\
 & + h.c.
 \end{aligned}$$

## Higgs basis

$$\Phi_1 = \begin{pmatrix} G^+ \\ \frac{1}{\sqrt{2}}(v + h_1^0 + iG^0) \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} H^+ \\ \frac{1}{\sqrt{2}}(h_2^0 + ih_3^0) \end{pmatrix}$$

## Mass Matrix

$$\mathcal{M}^2 = v^2 \begin{pmatrix} \lambda_1 & \text{Re}[\lambda_6] & -\text{Im}[\lambda_6] \\ \text{Re}[\lambda_6] & \frac{M^2}{v^2} + \frac{1}{2}(\lambda_3 + \lambda_4 + \text{Re}[\lambda_5]) & -\frac{1}{2}\text{Im}[\lambda_5] \\ -\text{Im}[\lambda_6] & -\frac{1}{2}\text{Im}[\lambda_5] & \frac{M^2}{v^2} + \frac{1}{2}(\lambda_3 + \lambda_4 - \text{Re}[\lambda_5]) \end{pmatrix}.$$

Pheno-motivated 2 types of alignments assumed:

**Higgs alignment**  $\lambda_6=0(=\mu_3) \Leftrightarrow$  No mixing among Higgses 125GeV  
Higgs measurements indicate SM like

**Yukawa alignment** to avoid FCNC at tree level


$\rightarrow$  4 complex parameters remain  $\zeta_e, \zeta_d, \zeta_u, \lambda_7$

EDM constraint :

$$-i \frac{d_e}{2} \bar{\psi}_e \sigma^{\mu\nu} \gamma^5 \psi_e F^{\mu\nu} = \sum_f^{t,b,\tau} \frac{\zeta_f}{\zeta_e} \text{ (loop diagram)} + \frac{\lambda_7}{\zeta_e} \text{ (loop diagram)} < 4.1 \times 10^{-30} \text{ ecm}$$

# EW production at LHC

- In 2HDM, always we have the EW productions




Cross section fixed only by the masses

→ No dependence on Yukawa param.

For 6 modes (HA, HH $\pm$ , AH $\pm$ , H+H-)  
 ~ 10-500 fb at 13 TeV LHC  
 (mH ~ 300GeV)

10<sup>3</sup> -10<sup>5</sup> Events at 139fb<sup>-1</sup>



Neutral  $H \rightarrow \tau\tau, bb$

Charged  $H^\pm \rightarrow \tau\nu, tb$

Heavy higgs also decay via

$$H_2 \rightarrow Z^* H_3, H_2 \rightarrow W^{*\pm} H^\mp$$

→ 4  $\tau$  lepton events expected

(BR depends on Yukawa param.)


- Latest LHC 4+ lepton (including taus) searches set very strong constraints



# Current LHC bounds

Various flavor constraints make the parameter space finite

$$m_{H_3} = m_{H^\pm} = 230 \text{ GeV}, m_{H_2} = 280 \text{ GeV}, \quad |\zeta_u| = 0.1$$



At HL-LHC multilepton  
 $BR_\tau \sim 0.2$  reachable

Large  $\tau\tau$  BR  
constrained by LHC  
multi lepton searches

Type X interpretation:  
 $(\zeta_e = \zeta_d^{-1} = \zeta_u^{-1})$

Currently,  
 $\zeta_e = \tan \beta \gtrsim 5$  excluded


At HL-LHC, up to

$\zeta_e = \tan \beta \gtrsim 1.5$   
would be sensitive

# Effects of Charged Higgs spectrum


$$m_{H_3} = m_{H^\pm} \leq m_{H_2}$$

[ light charged H ]



$$m_{H_3} \leq m_{H_2} = m_{H^\pm}$$

[ heavy charged H ]



All 6 modes produced similar in size


If  $H^\pm$  exists below,  $H_2$  decay into  $H^\pm \rightarrow \tau \nu$  : fewer leptons

→ heavier  $H^\pm$  provides stronger constraints ( $H \rightarrow \tau \tau$ ,  $bb$ ,  $H^\pm \rightarrow \tau \nu$ ,  $tb$ )

At  $\Delta M \sim m_W, m_Z$  the situation changes :

difference between light/heavy  $H^\pm$  more significant when open


# Current LHC bounds



Type X interpretation:  $\zeta_e = \tan \beta \gtrsim 2$  excluded at HL-LHC



# Current LHC bounds




Heavier  $H^\pm$  set stronger constraints  
( $H \rightarrow \tau\tau, bb, H^\pm \rightarrow \tau\nu, tb$ )

Type X interpretation:  $\zeta_e = \tan\beta \gtrsim 1$  excluded at HL-LHC

# Current/future reaches in type X-like case

S. Kanemura, M.T., K. Yagyu [Phys.Rev.D 105 (2022) 11, 115001]



Type X-like case, lighter charged Higgs case ( $\Delta m < 0$ ) constrained weaker.  
At HL-LHC almost all parameter space reachable below 2mt.

# Mass measurements at LHC





BR( $\tau\tau$ )  $\sim 1$  already constrained

Can we use  $bb\tau\tau$  mode?

H is heavy enough, collinear approx. valid

$$\vec{p}_{\nu_1} = \alpha_1 \vec{p}_{\tau_{\text{vis}1}}$$

$$\vec{p}_{\nu_2} = \alpha_2 \vec{p}_{\tau_{\text{vis}2}}$$






example )  $m_{H2}, m_{H3}, m_{H^\pm} = 280, 230, 280$  GeV

# Mass measurements at LHC

Multi  $\tau$  is excluded, more bottom BRs

Can we use  $bb\tau\tau$ ? Large  $t\bar{t}$  BG  $\sim 900\text{pb}$

For the masses **230 GeV**, **330 GeV** (signal xs  $\sim 10 - 50$  fb)






Only 1 prong  $\pi^+$  contributions  
 $\tau^\pm \rightarrow \pi^\pm \nu$  (BR  $\sim 10\%$ ) plotted,  
 other modes also usable

top BG reduced by  $\min(m_{b\tau}^{\text{rec}}) > 150$  GeV : efficiency  $\sim 0.04$  vs.  $10^{-4}$

We expect top BG controllable using further 2D cut

# Muon $g-2$ at 2HDMs

S. Iguro, T. Kitahara, M. Lang, M.T. [[arXiv:2304.09887](https://arxiv.org/abs/2304.09887)] [hep-ph]




Light A ( $m_A \sim 30$  GeV) is known as a possible solution to explain muon  $g-2$

Chargino-neutralino, Chargino-chargino searches at LHC in multi-tau SRs already exclude the type-X and aligned 2HDMs to explain muon  $g-2$ .

# Use of Tau-polarization in hLFV

M. Aoki, S. Kanemura, MT, L. Zamakhsyari [Phys.Rev.D 107 (2023) 5, 055037, arXiv: 2302.08489]




**Tau-decays preserve the information on its polarization**



$$-\mathcal{L}_{\text{LFV}} = y_{\tau\mu} h \bar{l}_{L\tau} l_{R\mu} + y_{\mu\tau} h \bar{l}_{L\mu} l_{R\tau} + h.c.$$

ATLAS reports an excess on  $h \rightarrow \tau\mu$  (BR $\sim$  0.1%)  
[arXiv:2302.05225 [hep-ex]]

**Sensitivity for the chirality, which would help to discriminate the UV models**



# Summary

- Baryon Asymmetry of the Univ. — too small CPV phase in the SM, thus CP violation beyond the SM required
- 125GeV Higgs is SM like → **Aligned CPV 2HDM**
- discussed Heavy Higgs discovery, measurements of mass, phases at LHC
- As the first step, we identify the current/future available region by multi-lepton searches at LHC
  - (counter-intuitively) heavier  $H^\pm$  cases stronger constrained  
**S. Kanemura, M.T., K. Yagyu [Phys.Rev.D 105 (2022) 11, 115001]**

At LHC, heavy Higgs mass measurable?

→ possible if they are light.

CPV phases at HL-LHC challenging [requires future ILC?]

- Correlation with 1st order phase transition, EW Baryon number generation
- muon  $g-2$  in 2HDMs, hLFV **S. Iguro, T. Kitahara, M. Lang, MT [arXiv:2304.09887 [hep-ph]]**  
**M. Aoki, S. Kanemura, MT, L. Zamakhsyari [Phys.Rev.D 107 (2023) 5, 055037, arXiv: 2302.08489]**

Backup




# CPV phase measurement



The former study : O(1) phases compatible to EDM constraints with heavy Higgses ~ 300GeV

S. Kanemura, M. Kubota, K. Yagyu [JHEP 08 (2020) 026]

## Angular distribution



Picture by [Jeans, Wilson, PRD98, 013007 (2018)]



At ILC,  $\zeta_e$  phase measurements using azimuthal angle dist. in  $H_2H_3 \rightarrow (bb)(\tau\tau)$

$$\mathcal{M} = \mathcal{M}_{h_1 h_2}^{H \rightarrow \tau^+ \tau^-} \mathcal{M}_{h_1}^{\tau^+} \mathcal{M}_{h_2}^{\tau^-}, \quad \mathcal{M}_h^{\tau^\pm} \sim e^{ih\phi} \quad \text{assuming the heavy higgs masses measured at LHC}$$

→ At LHC, can we discover the heavy higgses? Current reaches?  
Can we measure the masses?




# Current LHC bounds

Various flavor constraints make the parameter space finite

$$m_{H_3} = 230 \text{ GeV}, m_{H_2} = m_{H^\pm} = 280 \text{ GeV}, |\zeta_u| = 0.1$$

[ heavy charged H ]



Large  $\tau\tau$  BR  
constrained by LHC  
multi lepton searches

Type X interpretation:  
Currently,

$$\zeta_e = \tan \beta \gtrsim 3 \text{ excluded}$$

At HL-LHC, up to




$$\zeta_e = \tan \beta \gtrsim 1.5$$

would be sensitive




# Mass measurements at LHC

$m_{H2}, m_{H3}, m_{H\pm}$




280, 230, 280






280, 180, 280




230, 180, 230



280, 180, 180




# Fermion BR



$$R = \frac{1}{1 + r/\zeta^2}, \quad R_\tau = \frac{|\zeta_e|^2}{\zeta^2},$$

$$\zeta^2 = \frac{\sum_f \Gamma(H_3^0 \rightarrow f\bar{f})}{\Gamma_0}, \quad \Gamma_0 = \frac{\sqrt{2}G_F}{8\pi} m_{H_3^0} m_\tau^2,$$

$$\zeta_{50}^2 = r = \begin{cases} \frac{m_{H_3^0}^2}{2m_\tau^2} \sum_{V,\phi} \lambda^{3/2} \left( \frac{m_\phi^2}{m_{H_3^0}^2}, \frac{m_V^2}{m_{H_3^0}^2} \right) & (m_{H_3^0} - m_\phi \geq m_V) \\ \frac{9}{2\sqrt{2}\pi^2} \frac{G_F}{m_\tau^2} \sum_{V,\phi} m_V^4 \delta_V G \left( \frac{m_\phi^2}{m_{H_3^0}^2}, \frac{m_V^2}{m_{H_3^0}^2} \right) & (m_{H_3^0} - m_\phi < m_V), \end{cases}$$




$$R^\pm = \frac{1}{1 + r_\pm/\zeta_\pm^2}, \quad R_\tau^\pm = \frac{|\zeta_e|^2}{\zeta_\pm^2},$$

$$\zeta_\pm^2 = \frac{\sum_f \Gamma(H^\pm \rightarrow f\bar{f}')}{\Gamma_0 |H_3^0 \rightarrow H^\pm|} \simeq |\zeta_e|^2 + 3 \left( 1 - \frac{m_t^2}{m_{H^\pm}^2} \right)^2 \left( \frac{m_t^2}{m_\tau^2} |\zeta_u|^2 + \frac{m_b^2}{m_\tau^2} |\zeta_d|^2 \right),$$


$$r_\pm = \begin{cases} \frac{m_{H^\pm}^2}{2m_\tau^2} \lambda^{3/2} \left( \frac{m_{H_2^0}^2}{m_{H^\pm}^2}, \frac{m_W^2}{m_{H^\pm}^2} \right) & (m_{H^\pm} - m_{H_2^0} \geq m_W) \\ \frac{9}{2\sqrt{2}\pi^2} \frac{G_F}{m_\tau^2} m_W^4 G \left( \frac{m_{H_2^0}^2}{m_{H^\pm}^2}, \frac{m_W^2}{m_{H^\pm}^2} \right) & (m_{H^\pm} - m_{H_2^0} < m_W) \end{cases},$$

# Lepton BR

$$R_\tau = 50\%$$



$$R_\tau^\pm = 50\%$$



$$R_\tau^{-1} \simeq 1 + \frac{3m_b^2}{m_\tau^2} \frac{|\zeta_d|^2}{|\zeta_e|^2} + \left[ \frac{3m_c^2}{m_\tau^2} + \theta_{tt} \frac{3m_t^2}{m_\tau^2} \left( 1 - \frac{4m_t^2}{m_{H_3^0}^2} \right)^{3/2} \right] \frac{|\zeta_u|^2}{|\zeta_e|^2},$$

$$(R_\tau^\pm)^{-1} \simeq 1 + \left[ \frac{3m_s^2}{m_\tau^2} + \theta_{tb} \frac{3m_b^2}{m_\tau^2} \left( 1 - \frac{m_t^2}{m_{H^\pm}^2} \right)^2 \right] \frac{|\zeta_d|^2}{|\zeta_e|^2} + \left[ \frac{3m_c^2}{m_\tau^2} + \theta_{tb} \frac{3m_t^2}{m_\tau^2} \left( 1 - \frac{m_t^2}{m_{H^\pm}^2} \right)^2 \right] \frac{|\zeta_u|^2}{|\zeta_e|^2},$$